wwPDB EM Validation Summary Report

Mar 6, 2026 – 11:22 AM UTC

PDB ID	:	6UVN / pdb_00006uvn
EMDB ID	:	EMD-20908
Title	:	CryoEM structure of VcCascasde-TniQ complex
Authors	:	Chang, L.; Li, Z.; Zhang, H.
Deposited on	:	2019-11-03
Resolution	:	3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev132
MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	53% 39% 9%
2	B	640	48% 30% 22%
3	C	355	54% 32% 13%
3	D	355	56% 39% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	355	
3	F	355	
3	G	355	
3	H	355	
4	I	394	
4	J	394	
5	M	61	
6	N	98	

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 29487 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cas6.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	198	Total 1619	C 1030	N 287	O 295	S 7	0	0

- Molecule 2 is a protein called Cas8/5.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	498	Total 3921	C 2481	N 679	O 743	S 18	0	0

- Molecule 3 is a protein called Cas7.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	337	Total 2695	C 1717	N 463	O 501	S 14	0	0
3	F	337	Total 2695	C 1717	N 463	O 501	S 14	0	0
3	E	337	Total 2695	C 1717	N 463	O 501	S 14	0	0
3	H	337	Total 2695	C 1717	N 463	O 501	S 14	0	0
3	D	337	Total 2695	C 1717	N 463	O 501	S 14	0	0
3	C	308	Total 2495	C 1596	N 429	O 457	S 13	0	0

- Molecule 4 is a protein called TniQ.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	383	Total 3124	C 2004	N 540	O 561	S 19	0	0
4	J	389	Total 3171	C 2034	N 548	O 571	S 18	0	0

- Molecule 5 is DNA/RNA hybrid called crRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	M	61	Total	C	N	O	P	0	0
			1275	569	223	423	60		

- Molecule 6 is a protein called Cas8_HelicalBundle.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	N	81	Total	C	N	O	0	0
			405	243	81	81		

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
7	I	1	Total	Zn	0
			1	1	
7	J	1	Total	Zn	0
			1	1	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cas6

Chain A:



- Molecule 2: Cas8/5

Chain B:





• Molecule 3: Cas7

Chain G:



• Molecule 3: Cas7

Chain F:



• Molecule 3: Cas7

Chain E:





• Molecule 3: Cas7

Chain H:



• Molecule 3: Cas7

Chain D:



• Molecule 3: Cas7

Chain C:



• Molecule 4: TniQ

Chain I:



• Molecule 4: TniQ

Chain J:





• Molecule 5: crRNA

Chain M:



• Molecule 6: Cas8_HelicalBundle

Chain N:



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	582000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	35	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.252	Depositor
Minimum map value	-0.001	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.0215	Depositor
Map size (Å)	268.8, 268.8, 268.8	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.45	0/1656	0.52	0/2233
2	B	0.47	0/4007	0.55	2/5438 (0.0%)
3	C	0.53	0/2563	0.56	1/3481 (0.0%)
3	D	0.47	0/2766	0.49	0/3759
3	E	0.42	0/2766	0.44	0/3759
3	F	0.42	0/2766	0.46	0/3759
3	G	0.42	0/2766	0.48	0/3759
3	H	0.51	0/2766	0.50	0/3759
4	I	0.39	0/3211	0.57	1/4348 (0.0%)
4	J	0.37	0/3260	0.55	0/4416
5	M	0.49	0/1423	0.42	0/2217
All	All	0.45	0/29950	0.51	4/40928 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	391	PRO	N-CA-C	5.97	122.01	113.47
3	C	143	LYS	N-CA-C	-5.37	107.74	114.56
2	B	413	THR	N-CA-C	-5.35	105.56	111.71
4	I	290	PRO	N-CA-C	-5.08	102.01	112.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1619	0	1608	70	0
2	B	3921	0	3892	157	0
3	C	2495	0	2438	102	0
3	D	2695	0	2630	111	0
3	E	2695	0	2630	98	0
3	F	2695	0	2630	87	0
3	G	2695	0	2630	117	0
3	H	2695	0	2630	99	0
4	I	3124	0	3041	168	0
4	J	3171	0	3089	142	0
5	M	1275	0	643	59	0
6	N	405	0	90	4	0
7	I	1	0	0	0	0
7	J	1	0	0	0	0
All	All	29487	0	27951	1099	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 1099 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:289:LEU:HB3	4:I:290:PRO:CD	1.55	1.32
4:I:219:TYR:CD2	4:I:290:PRO:HG2	1.69	1.27
4:I:303:LEU:O	4:I:307:LEU:HD23	1.41	1.17
4:J:219:TYR:CE2	4:J:290:PRO:HG2	1.83	1.13
4:J:290:PRO:CD	4:J:291:GLU:H	1.64	1.08

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/217 (90%)	178 (91%)	18 (9%)	0	100	100
2	B	490/640 (77%)	436 (89%)	52 (11%)	2 (0%)	30	61
3	C	302/355 (85%)	285 (94%)	17 (6%)	0	100	100
3	D	333/355 (94%)	309 (93%)	24 (7%)	0	100	100
3	E	333/355 (94%)	311 (93%)	22 (7%)	0	100	100
3	F	333/355 (94%)	311 (93%)	22 (7%)	0	100	100
3	G	333/355 (94%)	316 (95%)	17 (5%)	0	100	100
3	H	333/355 (94%)	310 (93%)	23 (7%)	0	100	100
4	I	377/394 (96%)	336 (89%)	40 (11%)	1 (0%)	36	67
4	J	383/394 (97%)	340 (89%)	41 (11%)	2 (0%)	24	57
All	All	3413/3775 (90%)	3132 (92%)	276 (8%)	5 (0%)	49	78

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	I	289	LEU
4	J	290	PRO
2	B	15	LEU
4	J	291	GLU
2	B	33	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/198 (91%)	179 (99%)	1 (1%)	78	83
2	B	442/573 (77%)	440 (100%)	2 (0%)	81	85
3	C	276/310 (89%)	272 (99%)	4 (1%)	59	76
3	D	295/310 (95%)	295 (100%)	0	100	100
3	E	295/310 (95%)	293 (99%)	2 (1%)	76	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	F	295/310 (95%)	294 (100%)	1 (0%)	86 87
3	G	295/310 (95%)	293 (99%)	2 (1%)	76 82
3	H	295/310 (95%)	295 (100%)	0	100 100
4	I	345/356 (97%)	343 (99%)	2 (1%)	78 83
4	J	351/356 (99%)	349 (99%)	2 (1%)	78 83
All	All	3069/3343 (92%)	3053 (100%)	16 (0%)	78 85

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	J	289	LEU
4	I	289	LEU
3	C	101	ARG
4	I	181	CYS
3	E	229	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 79 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	141	ASN
4	J	46	HIS
3	C	189	ASN
4	I	46	HIS
4	J	207	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
6	N	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	N	300:UNK	C	304:UNK	N	13.31
1	N	346:UNK	C	355:UNK	N	11.87
1	N	326:UNK	C	333:UNK	N	7.08

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-20908. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices

6.2.1 Primary map



X Index: 128



Y Index: 128



Z Index: 128

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 127



Y Index: 121



Z Index: 95

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.0215. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 271 nm^3; this corresponds to an approximate mass of 245 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.323 Å^{-1}

8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-20908 and PDB model 6UVN. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0215 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0215).

9.4 Atom inclusion (i)



At the recommended contour level, 97% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0215) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9000	0.4860
A	0.8980	0.4790
B	0.8710	0.4710
C	0.9420	0.5160
D	0.9320	0.5150
E	0.9270	0.5150
F	0.9110	0.5100
G	0.9020	0.5010
H	0.9230	0.4970
I	0.8760	0.4600
J	0.8680	0.4440
M	0.9660	0.5060
N	0.5480	0.2250