



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:17 AM UTC

PDB ID : 4UWB / pdb_00004uwb
Title : Fibroblast growth factor receptor 1 kinase in complex with JK-P5
Authors : Beeston, H.; Tucker, J.; Kankanala, J.
Deposited on : 2014-08-11
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	18% 84% 12% ••
1	B	309	18% 83% 13% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	1770	-	-	X	-
4	SO4	B	1769	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FIBROBLAST GROWTH FACTOR RECEPTOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	1	0
			2324	1476	398	433	17			
1	B	299	Total	C	N	O	S	0	0	0
			2334	1487	397	433	17			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	457	GLY	-	expression tag	UNP P11362
A	488	ALA	CYS	engineered mutation	UNP P11362
A	584	SER	CYS	engineered mutation	UNP P11362
B	457	GLY	-	expression tag	UNP P11362
B	488	ALA	CYS	engineered mutation	UNP P11362
B	584	SER	CYS	engineered mutation	UNP P11362

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is N-[4-(4-methylpiperazin-1-yl)phenyl]-1H-indazole-3-carboxamide (CCD ID: JVT) (formula: C₁₉H₂₁N₅O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 25 19 5 1	0	0
3	B	1	Total C N O 25 19 5 1	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	109	Total	O	0	0
			109	109		
5	B	84	Total	O	0	0
			84	84		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1



• Molecule 1: FIBROBLAST GROWTH FACTOR RECEPTOR 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	208.62Å 57.32Å 65.63Å 90.00° 107.58° 90.00°	Depositor
Resolution (Å)	99.44 – 2.31 99.44 – 2.31	Depositor EDS
% Data completeness (in resolution range)	96.0 (99.44-2.31) 96.2 (99.44-2.31)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.229 , 0.272 0.234 , 0.276	Depositor DCC
R_{free} test set	1598 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4931	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SO4, JVT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/2372 (0.0%)	1.00	2/3211 (0.1%)
1	B	0.76	0/2383	0.99	5/3223 (0.2%)
All	All	0.79	1/4755 (0.0%)	1.00	7/6434 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	684	TRP	CA-C	5.47	1.60	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	744	ARG	CA-C-N	-6.68	113.42	120.03
1	B	744	ARG	C-N-CA	-6.68	113.42	120.03
1	B	577	ARG	CA-C-N	6.34	126.91	120.38
1	B	577	ARG	C-N-CA	6.34	126.91	120.38
1	A	473	LEU	CA-C-N	-6.02	113.58	119.78
1	A	473	LEU	C-N-CA	-6.02	113.58	119.78
1	B	727	ASN	N-CA-C	5.46	117.98	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2324	0	2288	34	0
1	B	2334	0	2294	24	0
2	A	12	0	18	2	0
2	B	8	0	12	2	0
3	A	25	0	21	1	0
3	B	25	0	21	0	0
4	A	5	0	0	2	0
4	B	5	0	0	2	0
5	A	109	0	0	4	0
5	B	84	0	0	2	0
All	All	4931	0	4654	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:576:ARG:HG2	1:A:593:GLU:HG2	1.31	1.11
1:A:576:ARG:CG	1:A:593:GLU:HG2	2.06	0.84
1:A:491:GLN:HG3	1:A:515:MET:HG3	1.65	0.77
1:A:589:HIS:O	1:A:591:PRO:HD3	1.92	0.69
1:A:576:ARG:HG2	1:A:593:GLU:CG	2.18	0.66
1:B:597:SER:O	1:B:601:VAL:HG23	1.97	0.65
1:A:707:GLU:HB2	5:A:2081:HOH:O	2.00	0.62
1:B:756:ARG:O	1:B:760:LEU:HG	2.02	0.60
1:A:504:LYS:HD2	1:A:506:ASN:HB2	1.85	0.59
1:A:667:MET:HG2	5:A:2054:HOH:O	2.04	0.57
1:A:723:SER:HB3	1:B:721:LYS:HZ2	1.71	0.56
1:A:616:SER:OG	5:A:2047:HOH:O	2.18	0.55
1:B:598:LYS:NZ	1:B:761:THR:O	2.34	0.53
1:A:491:GLN:HG3	1:A:515:MET:CG	2.37	0.53
1:A:736:CYS:O	1:A:744:ARG:HD3	2.09	0.52
1:B:547:LEU:H	2:B:1767:EDO:H11	1.75	0.52
1:A:583:TYR:CD1	1:A:583:TYR:C	2.88	0.51
1:A:589:HIS:O	1:A:589:HIS:CG	2.64	0.51
1:B:490:GLY:O	1:B:517:LYS:HD2	2.12	0.50
1:B:656:LYS:HE3	1:B:662:LEU:HD21	1.94	0.49
1:A:689:LEU:HD23	1:A:693:ILE:HD12	1.95	0.49
1:A:579:PRO:HA	1:A:589:HIS:O	2.12	0.49
1:B:478:LEU:HD21	1:B:513:VAL:HG21	1.95	0.48
1:B:569:LEU:O	1:B:573:LEU:HG	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:601:VAL:HG12	1:B:758:VAL:CG2	2.43	0.48
1:A:564:ALA:O	3:A:1769:JVT:H7	2.14	0.48
1:A:689:LEU:HD23	1:A:693:ILE:CD1	2.43	0.47
1:A:496:GLU:HB3	1:A:508:VAL:HG13	1.96	0.47
1:A:684:TRP:CD1	1:A:684:TRP:C	2.93	0.47
1:B:601:VAL:HG12	1:B:758:VAL:HG23	1.96	0.47
1:B:477:ARG:NH2	1:B:499:GLY:O	2.47	0.47
1:A:573:LEU:HD13	1:A:693:ILE:HA	1.97	0.46
1:B:661:ARG:HD2	4:B:1769:SO4:O4	2.15	0.46
1:A:576:ARG:CG	1:A:593:GLU:CG	2.87	0.46
1:A:601:VAL:HG12	1:A:758:VAL:CG2	2.46	0.45
1:A:653:TYR:HB3	1:A:664:VAL:HG11	1.98	0.45
1:A:661:ARG:HD2	4:A:1770:SO4:S	2.57	0.45
1:B:627:ARG:NH1	4:B:1769:SO4:O1	2.50	0.44
1:B:546:ASN:HA	2:B:1767:EDO:H11	1.98	0.44
1:B:656:LYS:HG2	1:B:662:LEU:HD23	1.99	0.44
1:A:477:ARG:NE	1:A:477:ARG:HA	2.33	0.43
1:A:573:LEU:CD1	1:A:693:ILE:HA	2.48	0.43
1:B:684:TRP:C	1:B:684:TRP:CD1	2.95	0.43
1:B:688:VAL:HG13	1:B:719:MET:HE1	1.99	0.43
1:B:494:LEU:HD23	5:B:2015:HOH:O	2.18	0.43
1:A:719:MET:H	2:A:1767:EDO:HO2	1.65	0.42
1:A:737:TRP:CD2	2:A:1767:EDO:H22	2.55	0.42
1:A:661:ARG:HD2	4:A:1770:SO4:O2	2.19	0.41
1:A:680:GLN:NE2	5:A:2072:HOH:O	2.47	0.41
1:A:601:VAL:HG12	1:A:758:VAL:HG22	2.02	0.41
1:A:667:MET:HE3	1:A:677:TYR:OH	2.19	0.41
1:A:723:SER:HB3	1:B:721:LYS:NZ	2.35	0.41
1:B:621:HIS:O	1:B:622:ARG:HB2	2.21	0.41
1:B:717:HIS:HA	5:B:2074:HOH:O	2.20	0.41
1:B:732:MET:O	1:B:735:ASP:HB2	2.21	0.40
1:B:622:ARG:HD2	1:B:646:ARG:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/309 (95%)	280 (95%)	11 (4%)	3 (1%)	12	14
1	B	291/309 (94%)	281 (97%)	10 (3%)	0	100	100
All	All	585/618 (95%)	561 (96%)	21 (4%)	3 (0%)	24	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	646	ARG
1	A	487	GLY
1	A	582	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/269 (92%)	241 (98%)	6 (2%)	43	60
1	B	247/269 (92%)	241 (98%)	6 (2%)	43	60
All	All	494/538 (92%)	482 (98%)	12 (2%)	43	60

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	508	VAL
1	A	518	SER
1	A	521	THR
1	A	582	GLU
1	A	644	LEU
1	A	723	SER
1	B	509	THR
1	B	534	MET
1	B	584	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	644	LEU
1	B	674	ASP
1	B	720	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	679	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	JVT	A	1769	-	28,28,28	0.48	0	37,39,39	0.49	0
4	SO4	B	1769	-	4,4,4	0.26	0	6,6,6	0.62	0
2	EDO	A	1768	-	3,3,3	0.55	0	2,2,2	0.03	0
4	SO4	A	1770	-	4,4,4	0.44	0	6,6,6	0.73	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	JVT	B	1768	-	28,28,28	0.41	0	37,39,39	0.64	1 (2%)
2	EDO	A	1766	-	3,3,3	0.53	0	2,2,2	0.38	0
2	EDO	A	1767	-	3,3,3	0.54	0	2,2,2	0.38	0
2	EDO	B	1767	-	3,3,3	0.76	0	2,2,2	0.36	0
2	EDO	B	1766	-	3,3,3	0.53	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JVT	A	1769	-	-	0/12/22/22	0/4/4/4
2	EDO	A	1768	-	-	0/1/1/1	-
3	JVT	B	1768	-	-	2/12/22/22	0/4/4/4
2	EDO	A	1766	-	-	1/1/1/1	-
2	EDO	A	1767	-	-	1/1/1/1	-
2	EDO	B	1767	-	-	1/1/1/1	-
2	EDO	B	1766	-	-	1/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1768	JVT	C13-C12-N4	-2.44	109.25	111.17

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1767	EDO	O1-C1-C2-O2
2	B	1766	EDO	O1-C1-C2-O2
3	B	1768	JVT	C10-C5-N1-C2
2	A	1766	EDO	O1-C1-C2-O2
3	B	1768	JVT	C6-C5-N1-C2
2	B	1767	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1769	JVT	1	0
4	B	1769	SO4	2	0
4	A	1770	SO4	2	0
2	A	1767	EDO	2	0
2	B	1767	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/309 (96%)	1.06	56 (18%) 3 4	18, 49, 91, 110	1 (0%)
1	B	299/309 (96%)	1.21	55 (18%) 3 4	37, 53, 87, 104	0
All	All	598/618 (96%)	1.14	111 (18%) 3 4	18, 51, 88, 110	1 (0%)

All (111) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	583	TYR	5.4
1	B	760	LEU	5.3
1	B	581	LEU	5.2
1	B	502	LYS	4.9
1	B	673	PHE	4.7
1	B	585	TYR	4.7
1	A	587	PRO	4.6
1	B	583	TYR	4.6
1	A	653	TYR	4.6
1	A	486	GLU	4.6
1	A	581	LEU	4.4
1	B	460	VAL	4.4
1	A	649	HIS	4.3
1	B	505	PRO	4.2
1	B	459	GLY	4.0
1	A	504	LYS	4.0
1	B	580	GLY	3.9
1	A	485	GLY	3.9
1	A	519	ASP	3.8
1	B	727	ASN	3.8
1	A	650	HIS	3.7
1	B	579	PRO	3.6
1	B	463	TYR	3.5
1	A	646	ARG	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	584	SER	3.5
1	A	644	LEU	3.5
1	A	490	GLY	3.4
1	B	501	ASP	3.4
1	A	588	SER	3.4
1	A	489	PHE	3.3
1	B	584	SER	3.3
1	A	488	ALA	3.3
1	B	677	TYR	3.3
1	A	594	GLN	3.2
1	A	582	GLU	3.2
1	A	658	THR	3.2
1	B	646	ARG	3.2
1	B	601	VAL	3.2
1	A	651	ILE	3.2
1	B	591	PRO	3.1
1	A	652	ASP	3.1
1	A	765	GLU	3.0
1	A	464	GLU	3.0
1	B	590	ASN	3.0
1	B	659	ASN	3.0
1	B	592	GLU	3.0
1	B	701	TYR	3.0
1	A	724	ASN	2.9
1	B	653	TYR	2.9
1	A	590	ASN	2.9
1	B	645	ALA	2.9
1	B	674	ASP	2.9
1	A	647	ASP	2.8
1	A	521	THR	2.8
1	B	658	THR	2.8
1	B	589	HIS	2.8
1	B	764	GLN	2.8
1	B	763	ASN	2.7
1	B	762	SER	2.7
1	B	489	PHE	2.6
1	A	654	TYR	2.6
1	A	505	PRO	2.6
1	A	673	PHE	2.6
1	A	487	GLY	2.6
1	B	582	GLU	2.6
1	A	589	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	491	GLN	2.5
1	A	465	LEU	2.5
1	A	579	PRO	2.5
1	A	701	TYR	2.5
1	A	526	SER	2.4
1	A	710	PHE	2.4
1	B	461	SER	2.4
1	A	503	ASP	2.4
1	A	522	GLU	2.4
1	B	710	PHE	2.4
1	B	738	HIS	2.4
1	A	592	GLU	2.4
1	A	494	LEU	2.4
1	A	591	PRO	2.3
1	B	488	ALA	2.3
1	B	694	PHE	2.3
1	A	672	LEU	2.3
1	A	483	PRO	2.3
1	B	490	GLY	2.3
1	B	499	GLY	2.3
1	B	675	ARG	2.3
1	A	603	CYS	2.3
1	B	728	GLU	2.3
1	B	676	ILE	2.2
1	A	659	ASN	2.2
1	A	520	ALA	2.2
1	A	657	THR	2.2
1	B	740	VAL	2.2
1	B	652	ASP	2.2
1	A	645	ALA	2.2
1	A	516	LEU	2.2
1	B	731	MET	2.2
1	B	469	PRO	2.2
1	B	578	PRO	2.2
1	B	756	ARG	2.1
1	A	482	LYS	2.1
1	B	672	LEU	2.1
1	B	650	HIS	2.1
1	B	595	LEU	2.1
1	B	757	ILE	2.1
1	A	552	THR	2.1
1	A	731	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	500	LEU	2.1
1	A	725	CYS	2.0
1	B	537	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	1770	5/5	0.86	0.14	57,70,71,76	0
2	EDO	B	1767	4/4	0.88	0.17	48,50,54,63	0
4	SO4	B	1769	5/5	0.88	0.27	57,57,68,70	0
3	JVT	A	1769	25/25	0.90	0.14	39,47,72,76	0
3	JVT	B	1768	25/25	0.90	0.13	42,50,64,67	0
2	EDO	B	1766	4/4	0.92	0.14	46,47,49,50	0
2	EDO	A	1767	4/4	0.92	0.13	42,45,45,48	0
2	EDO	A	1768	4/4	0.92	0.17	40,47,50,53	0
2	EDO	A	1766	4/4	0.93	0.11	41,45,45,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.