



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:57 AM UTC

PDB ID : 1UXA / pdb_00001uxa
Title : ADENOVIRUS AD37 FIBRE HEAD in complex with sialyl-lactose
Authors : Burmeister, W.P.; Guilligay, D.; Cusack, S.; Wadell, G.; Arnberg, N.
Deposited on : 2004-02-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	3% 79% 14% • 5%
1	B	194	5% 85% 9% • 6%
1	C	194	4% 81% 12% • 6%
2	D	2	50% 50%
2	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	2	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FIBER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1467	942	238	283	4			
1	B	183	Total	C	N	O	S	0	0	0
			1447	929	236	278	4			
1	C	183	Total	C	N	O	S	0	0	0
			1447	929	236	278	4			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	173	ALA	TYR	engineered mutation	UNP Q64823
B	173	ALA	TYR	engineered mutation	UNP Q64823
C	173	ALA	TYR	engineered mutation	UNP Q64823
A	174	MET	LEU	engineered mutation	UNP Q64823
B	174	MET	LEU	engineered mutation	UNP Q64823
C	174	MET	LEU	engineered mutation	UNP Q64823
A	175	GLY	VAL	engineered mutation	UNP Q64823
B	175	GLY	VAL	engineered mutation	UNP Q64823
C	175	GLY	VAL	engineered mutation	UNP Q64823
A	176	SER	ALA	engineered mutation	UNP Q64823
B	176	SER	ALA	engineered mutation	UNP Q64823
C	176	SER	ALA	engineered mutation	UNP Q64823

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			32	17	1	14			
2	E	2	Total	C	N	O	0	0	0
			32	17	1	14			

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	2	Total	C	N	O	0	0	0
			32	17	1	14			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		
4	C	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total 4	C 2	O 2	0	0
5	C	1	Total 4	C 2	O 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	168	Total 168	O 168	0	0
6	B	164	Total 164	O 164	0	0
6	C	196	Total 196	O 196	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FIBER PROTEIN



• Molecule 1: FIBER PROTEIN



• Molecule 1: FIBER PROTEIN



• Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



• Molecule 2: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose

Chain F:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.70Å 68.70Å 74.70Å 90.00° 94.90° 90.00°	Depositor
Resolution (Å)	45.18 – 1.50 45.18 – 1.50	Depositor EDS
% Data completeness (in resolution range)	92.0 (45.18-1.50) 92.0 (45.18-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.27 (at 1.50Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.180 , 0.205 0.184 , 0.182	Depositor DCC
R_{free} test set	4616 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4996	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT, SIA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.24	4/1502 (0.3%)	1.29	6/2040 (0.3%)
1	B	1.19	1/1481 (0.1%)	1.28	6/2011 (0.3%)
1	C	1.25	0/1481	1.28	7/2011 (0.3%)
All	All	1.23	5/4464 (0.1%)	1.28	19/6062 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	287	MET	SD-CE	-6.44	1.63	1.79
1	B	334	GLU	CA-C	5.71	1.60	1.52
1	A	354	THR	CA-C	-5.35	1.46	1.52
1	A	304	ARG	CB-CG	-5.23	1.36	1.52
1	A	286	PHE	N-CA	5.06	1.52	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	188	THR	N-CA-C	10.20	125.27	113.02
1	A	265	ALA	N-CA-C	8.84	121.87	111.71
1	B	238	ASN	CA-C-N	8.75	128.48	119.56
1	B	238	ASN	C-N-CA	8.75	128.48	119.56
1	A	354	THR	N-CA-C	7.21	121.24	109.85
1	B	236	LYS	N-CA-C	-7.08	102.98	111.69
1	C	236	LYS	N-CA-C	-7.03	103.05	111.69
1	A	236	LYS	N-CA-C	-6.41	103.55	112.45
1	B	354	THR	N-CA-C	6.21	119.77	110.14
1	C	354	THR	N-CA-C	6.14	119.55	109.85
1	C	353	GLU	N-CA-C	-5.83	99.15	109.06
1	B	265	ALA	N-CA-C	5.77	118.31	111.33
1	C	188	THR	CB-CA-C	-5.30	100.93	110.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	LYS	N-CA-C	5.30	117.97	111.82
1	C	189	THR	CA-C-N	-5.11	113.46	119.84
1	C	189	THR	C-N-CA	-5.11	113.46	119.84
1	B	304	ARG	CG-CD-NE	-5.06	100.87	112.00
1	A	297	SER	CA-C-N	-5.06	113.86	122.56
1	A	297	SER	C-N-CA	-5.06	113.86	122.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1467	0	1453	31	0
1	B	1447	0	1440	10	0
1	C	1447	0	1440	22	0
2	D	32	0	28	0	0
2	E	32	0	28	0	0
3	F	32	0	28	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	B	4	0	3	0	0
5	C	4	0	3	1	0
6	A	168	0	0	2	0
6	B	164	0	0	0	0
6	C	196	0	0	6	0
All	All	4996	0	4423	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:259:ASN:HD22	1:A:259:ASN:H	1.02	1.00
1:A:181:TYR:CA	1:A:184:ARG:HH11	1.84	0.90
1:A:181:TYR:O	1:A:212:LYS:HD3	1.79	0.82
1:C:268:ASN:HD22	1:C:269:PHE:H	1.28	0.81
1:C:183:THR:HB	1:C:269:PHE:CE1	2.18	0.77
1:A:259:ASN:HD22	1:A:259:ASN:N	1.83	0.76
1:A:181:TYR:CB	1:A:184:ARG:HH11	2.01	0.73
1:A:314:GLY:O	1:A:316:LYS:HD2	1.91	0.70
1:A:181:TYR:HA	1:A:184:ARG:HH11	1.56	0.69
1:A:181:TYR:HB3	1:A:286:PHE:HE1	1.57	0.69
1:A:270:ARG:HH11	1:B:215:SER:HB3	1.58	0.69
1:A:181:TYR:CB	1:A:286:PHE:HE1	2.06	0.69
1:C:276:VAL:HG21	6:C:2068:HOH:O	1.92	0.68
1:A:298:ASN:ND2	6:A:2093:HOH:O	2.25	0.68
1:A:259:ASN:H	1:A:259:ASN:ND2	1.81	0.66
1:C:187:TRP:HE1	1:C:275:ASN:ND2	1.95	0.64
1:C:295:LYS:HE2	6:C:2098:HOH:O	1.96	0.64
1:A:181:TYR:HA	1:A:184:ARG:HE	1.62	0.64
1:A:328:ASN:HD22	1:A:336:SER:H	1.42	0.64
1:C:328:ASN:HD22	1:C:336:SER:H	1.46	0.62
1:B:183:THR:CG2	1:B:183:THR:O	2.48	0.61
1:A:181:TYR:HB3	1:A:184:ARG:HH11	1.64	0.61
1:C:318:ASP:OD2	6:C:2131:HOH:O	2.16	0.60
1:C:264:LYS:NZ	6:C:2064:HOH:O	2.36	0.59
1:A:181:TYR:HB3	1:A:184:ARG:NH1	2.18	0.58
1:A:181:TYR:HB2	1:A:286:PHE:CE1	2.39	0.58
1:C:268:ASN:ND2	1:C:269:PHE:H	2.00	0.57
1:C:268:ASN:HD22	1:C:269:PHE:N	1.99	0.57
1:A:181:TYR:CB	1:A:286:PHE:CE1	2.88	0.56
1:B:197:THR:HG23	1:B:200:GLN:C	2.31	0.56
1:C:268:ASN:HB3	1:C:275:ASN:HB3	1.88	0.56
1:A:213:CYS:O	1:C:211:THR:HG21	2.06	0.55
1:C:281:GLU:HG2	6:C:2084:HOH:O	2.05	0.55
1:C:274:SER:C	1:C:275:ASN:HD22	2.15	0.55
1:C:278:THR:HG23	6:C:2078:HOH:O	2.08	0.54
1:C:187:TRP:NE1	1:C:275:ASN:ND2	2.55	0.54
1:C:187:TRP:NE1	1:C:275:ASN:HD21	2.06	0.53
1:A:269:PHE:HB3	6:A:2062:HOH:O	2.08	0.53
1:A:197:THR:HG23	1:A:200:GLN:C	2.34	0.53
1:B:328:ASN:HD22	1:B:336:SER:H	1.55	0.53
1:A:197:THR:HG23	1:A:200:GLN:O	2.12	0.49
1:A:211:THR:HG21	1:B:213:CYS:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:THR:HG23	1:B:200:GLN:O	2.13	0.49
1:B:330:GLU:OE2	1:B:338:THR:HG21	2.13	0.48
1:A:181:TYR:HA	1:A:184:ARG:NH1	2.26	0.47
1:A:314:GLY:C	1:A:316:LYS:HD2	2.39	0.47
1:B:187:TRP:CH2	1:B:190:PRO:HD3	2.50	0.47
1:C:187:TRP:CD1	1:C:275:ASN:HD21	2.35	0.45
1:B:183:THR:O	1:B:183:THR:HG22	2.17	0.44
1:A:251:ASN:HB3	1:A:257:LEU:HD21	2.00	0.44
1:C:330:GLU:OE2	1:C:338:THR:HG21	2.17	0.43
1:C:232:ILE:HD11	5:C:1369:ACT:H2	2.01	0.43
1:B:259:ASN:OD1	1:B:259:ASN:N	2.52	0.43
1:A:328:ASN:ND2	1:A:336:SER:H	2.14	0.42
1:A:187:TRP:CH2	1:A:190:PRO:HD3	2.55	0.42
1:A:316:LYS:HD2	1:A:316:LYS:N	2.35	0.42
1:A:186:LEU:HD11	1:A:280:TYR:CE1	2.55	0.41
1:A:191:ASP:OD1	1:A:191:ASP:C	2.64	0.41
1:C:325:THR:HA	1:C:338:THR:O	2.21	0.41
1:C:183:THR:HB	1:C:269:PHE:CZ	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/194 (94%)	177 (97%)	5 (3%)	1 (0%)	24	8
1	B	181/194 (93%)	176 (97%)	5 (3%)	0	100	100
1	C	181/194 (93%)	171 (94%)	9 (5%)	1 (1%)	21	6
All	All	545/582 (94%)	524 (96%)	19 (4%)	2 (0%)	30	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	189	THR
1	A	182	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/171 (96%)	163 (99%)	2 (1%)	63	38
1	B	163/171 (95%)	161 (99%)	2 (1%)	63	38
1	C	163/171 (95%)	162 (99%)	1 (1%)	78	62
All	All	491/513 (96%)	486 (99%)	5 (1%)	68	45

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	181	TYR
1	A	259	ASN
1	B	183	THR
1	B	277	SER
1	C	190	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	259	ASN
1	A	328	ASN
1	B	298	ASN
1	B	328	ASN
1	C	268	ASN
1	C	275	ASN
1	C	298	ASN
1	C	328	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GAL	D	1	2	12,12,12	0.43	0	17,17,17	0.67	0
2	SIA	D	2	2	20,20,21	1.40	3 (15%)	21,28,31	1.36	4 (19%)
2	GAL	E	1	2	12,12,12	0.43	0	17,17,17	0.52	0
2	SIA	E	2	2	20,20,21	1.26	2 (10%)	21,28,31	1.37	3 (14%)
3	GAL	F	1	3	12,12,12	0.60	0	17,17,17	0.80	0
3	SIA	F	2	3	20,20,21	0.95	0	21,28,31	1.29	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	D	1	2	-	0/2/22/22	0/1/1/1
2	SIA	D	2	2	-	0/18/34/38	0/1/1/1
2	GAL	E	1	2	-	0/2/22/22	0/1/1/1
2	SIA	E	2	2	-	1/18/34/38	0/1/1/1
3	GAL	F	1	3	-	0/2/22/22	0/1/1/1
3	SIA	F	2	3	-	0/18/34/38	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	SIA	C2-C1	3.74	1.56	1.52
2	E	2	SIA	C2-C1	3.48	1.56	1.52
2	D	2	SIA	C11-C10	2.56	1.55	1.50
2	E	2	SIA	C7-C6	2.46	1.56	1.52
2	D	2	SIA	C4-C5	2.38	1.55	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	SIA	O1B-C1-C2	3.54	121.92	112.71
3	F	2	SIA	O1B-C1-C2	3.47	121.74	112.71
2	D	2	SIA	O1B-C1-C2	3.13	120.85	112.71
2	E	2	SIA	O1B-C1-O1A	-2.29	118.88	124.08
2	D	2	SIA	O10-C10-N5	2.21	125.89	121.98
2	E	2	SIA	O10-C10-N5	2.16	125.80	121.98
2	D	2	SIA	C4-C5-N5	-2.10	106.31	110.44
2	D	2	SIA	C8-C7-C6	-2.02	109.25	113.05

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	2	SIA	O7-C7-C8-C9


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



Oligosaccharide Chain E



Oligosaccharide Chain F



5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACT	B	1369	4	3,3,3	1.36	0	3,3,3	1.70	1 (33%)
5	ACT	C	1369	4	3,3,3	1.06	0	3,3,3	1.04	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1369	ACT	O-C-CH3	-2.33	112.99	122.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1369	ACT	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/194 (95%)	0.23	6 (3%) 50 54	12, 19, 34, 51	0
1	B	183/194 (94%)	0.45	10 (5%) 30 33	11, 22, 44, 57	0
1	C	183/194 (94%)	0.07	8 (4%) 39 43	10, 17, 34, 50	0
All	All	551/582 (94%)	0.25	24 (4%) 39 43	10, 19, 38, 57	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	TYR	6.9
1	C	183	THR	6.3
1	C	276	VAL	4.6
1	B	183	THR	4.0
1	B	276	VAL	4.0
1	A	182	ASP	3.4
1	B	258	ASP	3.3
1	C	275	ASN	3.2
1	B	213	CYS	3.0
1	B	279	ALA	2.8
1	A	183	THR	2.5
1	B	281	GLU	2.5
1	B	266	TYR	2.5
1	C	273	ASN	2.5
1	C	278	THR	2.5
1	C	184	ARG	2.5
1	B	278	THR	2.4
1	B	186	LEU	2.4
1	A	298	ASN	2.2
1	A	213	CYS	2.1
1	B	298	ASN	2.1
1	A	266	TYR	2.1
1	C	272	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	274	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GAL	D	1	12/12	0.82	0.17	29,42,45,52	12
2	GAL	E	1	12/12	0.86	0.15	22,36,41,47	12
3	GAL	F	1	12/12	0.89	0.15	11,17,26,31	12
2	SIA	D	2	20/21	0.92	0.17	8,20,31,42	20
2	SIA	E	2	20/21	0.94	0.08	10,16,22,22	20
3	SIA	F	2	20/21	0.95	0.09	8,13,19,29	20

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ACT	B	1369	4/4	0.91	0.13	21,25,30,33	0
5	ACT	C	1369	4/4	0.93	0.14	9,27,31,36	0
4	ZN	B	1367	1/1	0.97	0.07	20,20,20,20	0
4	ZN	A	1367	1/1	0.98	0.08	27,27,27,27	0
4	ZN	C	1367	1/1	0.98	0.06	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.