



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:15 PM UTC

PDB ID : 4UXU / pdb_00004uxu
Title : Crystal Structure of the Extracellular Domain of the Human Alpha9 Nicotinic Acetylcholine Receptor In Complex with Methyllaconitine
Authors : Zouridakis, M.; Giastas, P.; Zarkadas, E.; Tzartos, S.J.
Deposited on : 2014-08-27
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	7% 92% ...
1	B	218	7% 88% 8% .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4168 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NEURONAL ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	2	0
			1720	1096	281	337	6			
1	B	210	Total	C	N	O	S	0	2	0
			1718	1095	280	337	6			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	213	HIS	-	expression tag	UNP Q9UGM1
A	214	HIS	-	expression tag	UNP Q9UGM1
A	215	HIS	-	expression tag	UNP Q9UGM1
A	216	HIS	-	expression tag	UNP Q9UGM1
A	217	HIS	-	expression tag	UNP Q9UGM1
A	218	HIS	-	expression tag	UNP Q9UGM1
B	213	HIS	-	expression tag	UNP Q9UGM1
B	214	HIS	-	expression tag	UNP Q9UGM1
B	215	HIS	-	expression tag	UNP Q9UGM1
B	216	HIS	-	expression tag	UNP Q9UGM1
B	217	HIS	-	expression tag	UNP Q9UGM1
B	218	HIS	-	expression tag	UNP Q9UGM1

- Molecule 2 is METHYLLYCACONITINE (CCD ID: MLK) (formula: C₃₇H₅₀N₂O₁₀).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			49	37	2	10		
2	B	1	Total	C	N	O	0	0
			49	37	2	10		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		
5	B	1	Total	Na	0	0
			1	1		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	264	Total	O	0	0
			264	264		
7	B	238	Total	O	0	0
			238	238		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NEURONAL ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA-9



- Molecule 1: NEURONAL ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA-9



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.43Å 66.25Å 75.19Å 90.00° 102.76° 90.00°	Depositor
Resolution (Å)	41.61 – 1.71 41.61 – 1.71	Depositor EDS
% Data completeness (in resolution range)	96.7 (41.61-1.71) 96.7 (41.61-1.71)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 1.71Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.206 , 0.235 0.214 , 0.234	Depositor DCC
R_{free} test set	3235 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.513	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4168	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, NAG, EDO, NA, MLK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1770	0.77	2/2419 (0.1%)
1	B	0.43	0/1768	0.70	1/2416 (0.0%)
All	All	0.43	0/3538	0.74	3/4835 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	ASP	CA-C-N	8.39	136.80	121.70
1	A	100	ASP	C-N-CA	8.39	136.80	121.70
1	B	167	GLY	N-CA-C	-5.34	105.79	111.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1720	0	1628	10	0
1	B	1718	0	1627	13	0
2	A	49	0	50	1	0
2	B	49	0	50	2	0
3	A	30	0	36	1	0
3	B	30	0	36	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	8	0	12	2	0
4	B	4	0	6	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	28	0	26	0	0
6	B	28	0	26	0	0
7	A	264	0	0	7	2
7	B	238	0	0	4	2
All	All	4168	0	3497	28	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:ASP:HB3	1:A:101:GLU:HB3	1.55	0.88
1:A:4:LYS:N	7:A:3001:HOH:O	2.10	0.82
1:A:107:ASN:ND2	7:A:3150:HOH:O	2.19	0.73
1:A:100:ASP:HB2	1:A:101:GLU:HB2	1.72	0.71
2:B:1213:MLK:O8	7:B:3223:HOH:O	2.08	0.71
1:B:196:SER:N	7:B:3211:HOH:O	2.23	0.66
7:A:3193:HOH:O	1:B:194:CYS:O	2.16	0.59
1:B:25:GLU:H	1:B:25:GLU:CD	2.13	0.56
1:B:197:GLU:H	1:B:197:GLU:CD	2.15	0.54
1:B:35:LEU:HD12	1:B:160:ILE:HD11	1.89	0.54
1:A:25:GLU:H	1:A:25:GLU:CD	2.18	0.51
1:B:97:LYS:NZ	1:B:99:ASP:O	2.35	0.51
1:B:179:GLU:HG3	1:B:209:LYS:HB3	1.95	0.48
1:A:196:SER:N	7:A:3236:HOH:O	2.46	0.48
1:A:100:ASP:OD2	7:A:3146:HOH:O	2.20	0.47
2:A:1213:MLK:H242	2:A:1213:MLK:O27	2.16	0.46
1:B:107:ASN:ND2	7:B:3141:HOH:O	2.50	0.44
1:B:176:VAL:O	1:B:211:ARG:HD2	2.17	0.44
1:B:3:GLY:N	7:B:3001:HOH:O	2.51	0.44
3:B:1214:EPE:H31	3:B:1214:EPE:H81	1.65	0.44
3:A:1215:EPE:H81	7:A:3162:HOH:O	2.18	0.43
1:B:211:ARG:NH2	3:B:1214:EPE:O3S	2.49	0.42
1:A:61:ILE:HD11	4:A:1216:EDO:H12	2.01	0.42
1:B:37:ILE:HD13	1:B:58:ILE:HG12	2.02	0.41
2:B:1213:MLK:O27	2:B:1213:MLK:H242	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99:ASP:OD2	1:B:101:GLU:HB2	2.21	0.41
1:A:154:ASN:HA	7:A:3183:HOH:O	2.20	0.41
1:A:59:ARG:NH1	4:A:1216:EDO:H11	2.35	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:3220:HOH:O	7:B:3046:HOH:O[2_554]	1.98	0.22
7:A:3082:HOH:O	7:B:3042:HOH:O[2_554]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/218 (96%)	204 (97%)	5 (2%)	1 (0%)	24	11
1	B	210/218 (96%)	200 (95%)	10 (5%)	0	100	100
All	All	420/436 (96%)	404 (96%)	15 (4%)	1 (0%)	43	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/200 (97%)	193 (100%)	1 (0%)	81	71
1	B	194/200 (97%)	192 (99%)	2 (1%)	68	53
All	All	388/400 (97%)	385 (99%)	3 (1%)	73	61

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	SER
1	B	27	THR
1	B	196	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	7	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	1217	-	3,3,3	0.40	0	2,2,2	0.36	0
6	NAG	A	2215	1	14,14,15	0.40	0	17,19,21	0.50	0
6	NAG	B	2214	1	14,14,15	0.36	0	17,19,21	0.75	1 (5%)
2	MLK	B	1213	-	56,56,56	1.59	6 (10%)	82,92,92	1.34	11 (13%)
3	EPE	B	1214	-	15,15,15	2.00	1 (6%)	19,20,20	2.71	9 (47%)
3	EPE	A	1215	-	15,15,15	1.96	1 (6%)	19,20,20	2.69	8 (42%)
3	EPE	B	1216	-	15,15,15	1.87	1 (6%)	19,20,20	2.61	8 (42%)
4	EDO	A	1216	-	3,3,3	0.47	0	2,2,2	0.24	0
6	NAG	A	2214	1	14,14,15	0.18	0	17,19,21	1.25	1 (5%)
6	NAG	B	2215	1	14,14,15	0.33	0	17,19,21	0.47	0
3	EPE	A	1214	-	15,15,15	2.02	1 (6%)	19,20,20	2.59	8 (42%)
2	MLK	A	1213	-	56,56,56	1.59	6 (10%)	82,92,92	1.37	14 (17%)
4	EDO	B	1215	-	3,3,3	0.50	0	2,2,2	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1217	-	-	0/1/1/1	-
6	NAG	A	2215	1	-	0/6/23/26	0/1/1/1
6	NAG	B	2214	1	-	0/6/23/26	0/1/1/1
2	MLK	B	1213	-	-	0/24/141/141	0/2/8/8
3	EPE	B	1214	-	-	3/9/19/19	0/1/1/1
3	EPE	A	1215	-	-	7/9/19/19	0/1/1/1
3	EPE	B	1216	-	-	7/9/19/19	0/1/1/1
4	EDO	A	1216	-	-	0/1/1/1	-
6	NAG	A	2214	1	-	0/6/23/26	0/1/1/1
6	NAG	B	2215	1	-	0/6/23/26	0/1/1/1
3	EPE	A	1214	-	-	2/9/19/19	0/1/1/1
2	MLK	A	1213	-	-	2/24/141/141	0/2/8/8
4	EDO	B	1215	-	-	0/1/1/1	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1214	EPE	C10-S	-7.53	1.67	1.77
3	B	1214	EPE	C10-S	-7.40	1.67	1.77
3	A	1215	EPE	C10-S	-7.32	1.67	1.77
3	B	1216	EPE	C10-S	-6.92	1.67	1.77
2	B	1213	MLK	C8-N7	-5.66	1.32	1.40
2	A	1213	MLK	C8-N7	-5.21	1.33	1.40
2	A	1213	MLK	C1-C6	5.11	1.48	1.41
2	A	1213	MLK	C11-N7	-5.03	1.32	1.40
2	B	1213	MLK	C1-C6	4.99	1.48	1.41
2	B	1213	MLK	C11-N7	-4.57	1.33	1.40
2	A	1213	MLK	O14-C13	4.30	1.44	1.33
2	B	1213	MLK	O14-C13	4.29	1.44	1.33
2	B	1213	MLK	C18-C17	-3.36	1.52	1.56
2	A	1213	MLK	C18-C17	-3.29	1.52	1.56
2	A	1213	MLK	C6-N7	-2.91	1.40	1.44
2	B	1213	MLK	C6-N7	-2.74	1.40	1.44

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1215	EPE	C6-N1-C2	5.66	121.03	108.84
3	B	1216	EPE	C6-N1-C2	5.56	120.81	108.84
3	A	1214	EPE	C6-N1-C2	5.53	120.74	108.84
3	B	1216	EPE	C5-N4-C3	5.43	120.53	108.84
3	A	1214	EPE	C5-N4-C3	5.41	120.49	108.84
3	A	1215	EPE	C5-N4-C3	5.38	120.42	108.84
3	B	1214	EPE	C6-N1-C2	5.35	120.37	108.84
3	B	1214	EPE	C5-N4-C3	5.31	120.27	108.84
6	A	2214	NAG	C1-O5-C5	4.78	118.59	112.19
3	B	1216	EPE	C9-N1-C2	3.93	121.70	111.24
3	B	1216	EPE	C7-N4-C3	3.81	121.39	111.24
3	B	1214	EPE	C7-N4-C3	3.76	121.26	111.24
2	A	1213	MLK	C9-C8-N7	3.60	110.88	108.06
2	B	1213	MLK	C21-C16-C17	3.54	114.61	109.91
2	B	1213	MLK	C9-C8-N7	3.52	110.83	108.06
2	A	1213	MLK	O27-C27-C26	3.45	113.63	110.45
3	A	1215	EPE	C7-N4-C5	3.43	120.38	111.24
3	B	1214	EPE	C9-N1-C2	3.40	120.30	111.24
3	A	1214	EPE	C7-N4-C3	3.40	120.30	111.24
2	A	1213	MLK	C10-C11-N7	3.24	110.20	108.24
2	B	1213	MLK	O27-C27-C26	3.21	113.41	110.45
3	A	1214	EPE	C9-N1-C6	3.18	119.71	111.24
3	B	1214	EPE	O2S-S-C10	3.17	111.51	106.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1215	EPE	C9-N1-C2	3.13	119.59	111.24
3	A	1214	EPE	C9-N1-C2	3.12	119.55	111.24
2	B	1213	MLK	C10-C11-N7	3.09	110.11	108.24
3	A	1215	EPE	C9-N1-C6	3.06	119.39	111.24
3	B	1214	EPE	C9-N1-C6	3.03	119.32	111.24
3	A	1215	EPE	C7-N4-C3	2.99	119.19	111.24
3	A	1214	EPE	C7-N4-C5	2.98	119.17	111.24
2	B	1213	MLK	O32-C32-C27	-2.97	103.48	107.97
3	B	1214	EPE	O1S-S-C10	2.90	111.12	106.73
2	A	1213	MLK	O8-C8-C9	-2.86	123.10	127.28
2	A	1213	MLK	O14-C13-C1	2.83	117.79	112.24
2	B	1213	MLK	O8-C8-C9	-2.72	123.30	127.28
3	B	1214	EPE	C7-N4-C5	2.69	118.40	111.24
2	A	1213	MLK	O14-C15-C16	-2.63	104.58	108.39
2	A	1213	MLK	O14-C13-O13	-2.61	118.45	123.67
3	B	1216	EPE	O1S-S-C10	2.59	110.64	106.73
6	B	2214	NAG	C1-O5-C5	2.59	115.66	112.19
3	B	1216	EPE	C7-N4-C5	2.53	117.98	111.24
2	A	1213	MLK	C23-C16-C17	2.47	110.07	108.22
2	A	1213	MLK	C21-C16-C17	2.47	113.18	109.91
2	A	1213	MLK	O32-C32-C27	-2.42	104.33	107.97
3	A	1215	EPE	C5-C6-N1	-2.31	106.00	110.65
2	A	1213	MLK	C32-C31-C30	-2.29	111.34	112.89
3	B	1216	EPE	O2S-S-C10	2.29	110.18	106.73
2	B	1213	MLK	O14-C15-C16	-2.29	105.08	108.39
3	B	1216	EPE	C9-N1-C6	2.28	117.32	111.24
3	A	1215	EPE	O2S-S-C10	2.24	110.12	106.73
2	A	1213	MLK	C16-C23-N23	-2.24	108.95	113.14
2	A	1213	MLK	C21-C16-C23	-2.16	109.46	111.45
2	B	1213	MLK	C16-C23-N23	-2.14	109.14	113.14
3	A	1214	EPE	C3-C2-N1	-2.06	106.49	110.65
2	B	1213	MLK	C32-C31-C30	-2.05	111.50	112.89
3	A	1214	EPE	C2-C3-N4	-2.03	106.55	110.65
2	B	1213	MLK	O14-C13-C1	2.03	116.22	112.24
3	B	1214	EPE	C10-C9-N1	-2.02	104.71	112.36
2	A	1213	MLK	C23-N23-C24	-2.02	107.90	111.25
2	B	1213	MLK	O14-C13-O13	-2.01	119.65	123.67

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1215	EPE	C9-C10-S-O1S
3	B	1214	EPE	C10-C9-N1-C6
3	B	1214	EPE	C8-C7-N4-C3
3	B	1216	EPE	C10-C9-N1-C2
3	B	1216	EPE	C8-C7-N4-C3
3	A	1214	EPE	C8-C7-N4-C3
3	B	1216	EPE	N4-C7-C8-O8
3	A	1215	EPE	C9-C10-S-O3S
3	B	1216	EPE	C9-C10-S-O3S
3	A	1215	EPE	N4-C7-C8-O8
3	B	1216	EPE	S-C10-C9-N1
3	A	1215	EPE	C8-C7-N4-C3
3	A	1215	EPE	C10-C9-N1-C2
3	A	1215	EPE	C9-C10-S-O2S
3	B	1216	EPE	C9-C10-S-O1S
3	B	1216	EPE	C9-C10-S-O2S
3	A	1214	EPE	C10-C9-N1-C2
3	A	1215	EPE	C10-C9-N1-C6
3	B	1214	EPE	N4-C7-C8-O8
2	A	1213	MLK	C34-C35-O35-C37
2	A	1213	MLK	C27-C28-O28-C29


There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1213	MLK	2	0
3	B	1214	EPE	2	0
3	A	1215	EPE	1	0
4	A	1216	EDO	2	0
2	A	1213	MLK	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/218 (96%)	0.38	15 (7%) 22 27	13, 22, 47, 94	2 (0%)
1	B	210/218 (96%)	0.52	15 (7%) 22 27	12, 23, 52, 94	2 (0%)
All	All	420/436 (96%)	0.45	30 (7%) 22 27	12, 23, 51, 94	4 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	PRO	6.4
1	B	105	PRO	5.4
1	B	3	GLY	4.0
1	A	3	GLY	3.9
1	A	194	CYS	3.8
1	B	106	VAL	3.7
1	B	103	SER	3.7
1	A	103	SER	3.5
1	B	195	CYS	3.4
1	A	136	TYR	3.3
1	A	106	VAL	3.3
1	A	100	ASP	3.1
1	B	212	SER	3.1
1	B	194	CYS	3.1
1	B	107	ASN	3.0
1	B	25	GLU	2.8
1	A	212	SER	2.8
1	B	102	SER	2.8
1	B	136	TYR	2.7
1	B	99	ASP	2.7
1	A	16	ASP	2.6
1	A	25	GLU	2.6
1	A	104	GLU	2.5
1	A	195	CYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	100	ASP	2.3
1	A	102	SER	2.3
1	A	98	ALA	2.2
1	A	99	ASP	2.1
1	B	27	THR	2.1
1	B	28	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EPE	B	1216	15/15	0.68	0.23	46,50,85,88	0
4	EDO	B	1215	4/4	0.72	0.23	35,44,49,51	0
4	EDO	A	1216	4/4	0.78	0.18	36,40,42,46	0
3	EPE	B	1214	15/15	0.80	0.21	26,51,64,65	0
3	EPE	A	1215	15/15	0.82	0.17	29,45,72,76	0
6	NAG	A	2214	14/15	0.82	0.14	27,33,40,44	0
4	EDO	A	1217	4/4	0.83	0.16	36,41,43,50	0
3	EPE	A	1214	15/15	0.88	0.15	25,41,52,52	0
6	NAG	B	2214	14/15	0.88	0.11	28,36,39,44	0
2	MLK	B	1213	49/49	0.94	0.09	13,20,37,43	0
6	NAG	A	2215	14/15	0.94	0.08	17,23,33,37	0
2	MLK	A	1213	49/49	0.94	0.09	13,19,35,40	0
6	NAG	B	2215	14/15	0.94	0.08	16,24,35,39	0
5	NA	A	1218	1/1	0.95	0.16	32,32,32,32	0
5	NA	B	1217	1/1	0.98	0.17	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MLK B 1213:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MLK A 1213:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.