



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:33 AM UTC

PDB ID : 5UX6 / pdb_00005ux6
Title : Structure of Human POFUT1 in its apo form
Authors : Xu, X.; McMillan, B.; Blacklow, S.C.
Deposited on : 2017-02-22
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	10% 90% 7% .
1	B	367	7% 89% 7% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11794 atoms, of which 5493 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GDP-fucose protein O-fucosyltransferase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	355	Total	C	H	N	O	S	0	0	0
			5573	1831	2740	483	501	18			
1	B	353	Total	C	H	N	O	S	0	0	0
			5520	1816	2718	477	491	18			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	160	ALA	ASN	engineered mutation	UNP Q9H488
A	385	HIS	-	expression tag	UNP Q9H488
A	386	HIS	-	expression tag	UNP Q9H488
A	387	HIS	-	expression tag	UNP Q9H488
A	388	HIS	-	expression tag	UNP Q9H488
A	389	HIS	-	expression tag	UNP Q9H488
A	390	HIS	-	expression tag	UNP Q9H488
B	160	ALA	ASN	engineered mutation	UNP Q9H488
B	385	HIS	-	expression tag	UNP Q9H488
B	386	HIS	-	expression tag	UNP Q9H488
B	387	HIS	-	expression tag	UNP Q9H488
B	388	HIS	-	expression tag	UNP Q9H488
B	389	HIS	-	expression tag	UNP Q9H488
B	390	HIS	-	expression tag	UNP Q9H488

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	303	Total 303	O 303	0	0
4	B	329	Total 329	O 329	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GDP-fucose protein O-fucosyltransferase 1



- Molecule 1: GDP-fucose protein O-fucosyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.20Å 132.24Å 72.06Å 90.00° 106.44° 90.00°	Depositor
Resolution (Å)	47.78 – 2.09 47.78 – 2.09	Depositor EDS
% Data completeness (in resolution range)	98.3 (47.78-2.09) 98.5 (47.78-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 2.08Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.182 , 0.228 0.184 , 0.230	Depositor DCC
R_{free} test set	1993 reflections (3.73%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11794	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.19	0/2925	0.38	0/3974
1	B	0.21	0/2893	0.38	0/3931
All	All	0.20	0/5818	0.38	0/7905

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2833	2740	2740	13	0
1	B	2802	2718	2718	14	0
2	A	14	13	13	0	0
2	B	14	14	13	0	0
3	A	6	8	8	0	0
4	A	303	0	0	5	0
4	B	329	0	0	5	0
All	All	6301	5493	5492	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:PRO:O	4:A:501:HOH:O	2.02	0.77
1:B:244:ASP:OD1	4:B:501:HOH:O	2.04	0.75
1:A:88:GLN:OE1	4:A:502:HOH:O	2.11	0.68
1:A:286:ASP:OD2	4:A:503:HOH:O	2.13	0.64
1:B:111:LYS:NZ	4:B:505:HOH:O	2.29	0.64
1:B:266:GLN:O	1:B:268:VAL:N	2.31	0.63
1:A:198:GLN:NE2	4:A:509:HOH:O	2.34	0.59
1:A:93:LEU:HD12	4:A:502:HOH:O	2.03	0.58
1:B:136:ASP:OD1	1:B:139:THR:HG23	2.11	0.49
1:B:204:GLU:HG3	4:B:504:HOH:O	2.13	0.48
1:A:301:LEU:HD11	1:A:349:HIS:HB2	1.96	0.48
1:B:266:GLN:C	1:B:267:CYS:SG	2.97	0.47
1:A:134:SER:HB3	1:A:135:PRO:CD	2.44	0.47
1:A:177:GLN:OE1	1:A:177:GLN:N	2.43	0.47
1:B:168:ILE:O	4:B:503:HOH:O	2.20	0.46
1:B:301:LEU:HD11	1:B:349:HIS:HB2	1.97	0.45
1:B:132:GLN:HA	1:B:137:LYS:HA	1.99	0.45
1:A:305:SER:HA	1:A:328:LYS:O	2.18	0.44
1:A:134:SER:HB3	1:A:135:PRO:HD2	2.00	0.43
1:B:59:LYS:NZ	1:B:99:TYR:O	2.47	0.43
1:A:316:VAL:HB	1:A:317:PRO:HD3	1.99	0.43
1:A:88:GLN:HA	1:A:91:PHE:O	2.18	0.43
1:B:181:ARG:NH1	4:B:521:HOH:O	2.51	0.43
1:B:245:TRP:CZ2	1:B:355:VAL:HG23	2.55	0.42
1:B:305:SER:HA	1:B:328:LYS:O	2.20	0.42
1:B:209:GLN:HB3	1:B:368:LEU:CD1	2.51	0.40
1:A:87:TYR:CE1	1:A:91:PHE:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/367 (96%)	340 (97%)	10 (3%)	1 (0%)	36	36
1	B	349/367 (95%)	341 (98%)	6 (2%)	2 (1%)	21	18
All	All	700/734 (95%)	681 (97%)	16 (2%)	3 (0%)	30	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	267	CYS
1	A	312	SER
1	B	312	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/317 (96%)	295 (97%)	8 (3%)	40	46
1	B	297/317 (94%)	292 (98%)	5 (2%)	53	62
All	All	600/634 (95%)	587 (98%)	13 (2%)	45	53

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	THR
1	A	136	ASP
1	A	137	LYS
1	A	164	LEU
1	A	243	SER
1	A	254	ASP
1	A	272	ARG
1	A	359	THR
1	B	173	SER
1	B	186	GLU
1	B	267	CYS
1	B	268	VAL
1	B	274	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	227	HIS
1	A	346	GLN
1	B	146	ASN
1	B	346	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	401	1	14,14,15	0.76	1 (7%)	17,19,21	0.79	1 (5%)
2	NAG	B	401	1	14,14,15	0.99	1 (7%)	17,19,21	0.74	0
3	GOL	A	402	-	5,5,5	0.46	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	401	1	-	2/6/23/26	0/1/1/1
2	NAG	B	401	1	-	2/6/23/26	0/1/1/1
3	GOL	A	402	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAG	C1-C2	3.12	1.56	1.52
2	A	401	NAG	O5-C1	-2.50	1.39	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAG	O4-C4-C3	-2.03	105.58	110.38

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	GOL	O1-C1-C2-C3
2	A	401	NAG	C8-C7-N2-C2
2	A	401	NAG	O7-C7-N2-C2
2	B	401	NAG	C4-C5-C6-O6
2	B	401	NAG	O5-C5-C6-O6
3	A	402	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/367 (96%)	0.03	35 (9%) 13 13	12, 27, 87, 145	0
1	B	353/367 (96%)	0.00	25 (7%) 22 23	13, 28, 79, 143	0
All	All	708/734 (96%)	0.02	60 (8%) 16 17	12, 27, 83, 145	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	267	CYS	6.3
1	A	79	PRO	6.1
1	B	78	PRO	6.0
1	B	383	LYS	5.6
1	A	254	ASP	5.4
1	A	135	PRO	5.3
1	A	80	PHE	5.2
1	B	271	SER	5.1
1	B	74	GLN	5.0
1	B	272	ARG	4.8
1	B	274	THR	4.7
1	B	382	PRO	4.6
1	A	75	HIS	4.5
1	A	139	THR	4.4
1	A	76	HIS	4.4
1	B	179	SER	4.2
1	B	273	SER	4.2
1	B	266	GLN	4.1
1	B	268	VAL	4.1
1	B	79	PRO	4.0
1	B	354	CYS	4.0
1	A	81	THR	3.8
1	A	134	SER	3.8
1	A	133	ARG	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	80	PHE	3.6
1	B	275	ALA	3.5
1	B	217	GLU	3.5
1	B	171	SER	3.4
1	A	138	LYS	3.4
1	A	164	LEU	3.2
1	A	255	GLY	3.1
1	B	83	LEU	3.1
1	B	88	GLN	3.0
1	A	137	LYS	3.0
1	A	175	ARG	2.8
1	A	299	ARG	2.7
1	A	383	LYS	2.7
1	B	270	TYR	2.7
1	A	132	GLN	2.6
1	A	272	ARG	2.6
1	A	136	ASP	2.6
1	A	181	ARG	2.6
1	A	167	GLY	2.5
1	A	83	LEU	2.5
1	A	88	GLN	2.5
1	A	185	LYS	2.5
1	A	74	GLN	2.4
1	A	130	ALA	2.4
1	B	28	SER	2.4
1	A	129	VAL	2.4
1	A	384	LEU	2.3
1	A	110	GLU	2.3
1	A	144	GLU	2.2
1	B	81	THR	2.2
1	A	131	ALA	2.2
1	B	264	SER	2.1
1	B	172	ALA	2.1
1	A	171	SER	2.1
1	A	180	GLN	2.0
1	A	140	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	401	14/15	0.70	0.22	59,80,123,123	0
2	NAG	B	401	14/15	0.73	0.19	63,87,112,112	0
3	GOL	A	402	6/6	0.86	0.38	98,125,147,155	0

6.5 Other polymers [i](#)

There are no such residues in this entry.