



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 29, 2026 – 01:58 AM UTC

PDB ID : 7UX7 / pdb_00007ux7
Title : Crystal structure of MfnG, an L- and D-tyrosine O-methyltransferase from the marformycin biosynthesis pathway of *Streptomyces drozdowiczii*, with SAH bound at 1.2 Å resolution (P212121 - form II)
Authors : Miller, M.D.; Wu, K.-L.; Xu, W.; Xiao, H.; Philips Jr., G.N.
Deposited on : 2022-05-05
Resolution : 1.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1380 (1.16-1.12)
Clashscore	190562	1393 (1.16-1.12)
Ramachandran outliers	187476	1369 (1.16-1.12)
Sidechain outliers	187428	1369 (1.16-1.12)
RSRZ outliers	180081	1379 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	8% 93% 6% .
1	B	384	5% 87% 7% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13317 atoms, of which 6104 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MfnG.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	378	Total	C	H	N	O	S	0	46	0
			6239	1988	3066	569	606	10			
1	B	361	Total	C	H	N	O	S	0	35	0
			6078	1934	3000	563	570	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	cloning artifact	UNP A0A0D4WTP2
A	1	VAL	-	cloning artifact	UNP A0A0D4WTP2
A	376	ALA	-	expression tag	UNP A0A0D4WTP2
A	377	SER	-	expression tag	UNP A0A0D4WTP2
A	378	GLU	-	expression tag	UNP A0A0D4WTP2
A	379	ASN	-	expression tag	UNP A0A0D4WTP2
A	380	LEU	-	expression tag	UNP A0A0D4WTP2
A	381	TYR	-	expression tag	UNP A0A0D4WTP2
A	382	PHE	-	expression tag	UNP A0A0D4WTP2
A	383	GLN	-	expression tag	UNP A0A0D4WTP2
B	0	MET	-	cloning artifact	UNP A0A0D4WTP2
B	1	VAL	-	cloning artifact	UNP A0A0D4WTP2
B	376	ALA	-	expression tag	UNP A0A0D4WTP2
B	377	SER	-	expression tag	UNP A0A0D4WTP2
B	378	GLU	-	expression tag	UNP A0A0D4WTP2
B	379	ASN	-	expression tag	UNP A0A0D4WTP2
B	380	LEU	-	expression tag	UNP A0A0D4WTP2
B	381	TYR	-	expression tag	UNP A0A0D4WTP2
B	382	PHE	-	expression tag	UNP A0A0D4WTP2
B	383	GLN	-	expression tag	UNP A0A0D4WTP2

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 45	C 14	H 19	N 6	O 5	S 1	0	0
2	B	1	Total 45	C 14	H 19	N 6	O 5	S 1	0	0

- Molecule 3 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			9	9		
3	B	1	Total	O	0	0
			9	9		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	443	Total	O	0	39
			461	461		
4	B	418	Total	O	0	37
			431	431		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MfnG



• Molecule 1: MfnG



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.69Å 86.33Å 134.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.18 – 1.14 35.18 – 1.14	Depositor EDS
% Data completeness (in resolution range)	83.4 (35.18-1.14) 83.5 (35.18-1.14)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX 1.20.1	Depositor
R, R_{free}	0.119 , 0.142 0.119 , 0.142	Depositor DCC
R_{free} test set	10679 reflections (4.15%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.7	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	13317	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	1/3243 (0.0%)	0.75	2/4408 (0.0%)
1	B	0.57	0/3150	0.75	0/4283
All	All	0.56	1/6393 (0.0%)	0.75	2/8691 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	358	ARG	CG-CD	-6.87	1.31	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ARG	CG-CD-NE	-5.62	99.62	112.00
1	A	208	SER	CB-CA-C	5.17	120.59	110.67

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	364	ARG	Sidechain
1	B	248[B]	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3173	3066	3042	15	0
1	B	3078	3000	2982	17	0
2	A	26	19	19	0	0
2	B	26	19	19	0	0
3	A	9	0	0	1	0
3	B	9	0	0	1	0
4	A	461	0	0	2	0
4	B	431	0	0	3	0
All	All	7213	6104	6062	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:402:UNL:O1	3:B:402:UNL:O2	1.52	1.23
3:A:402:UNL:O1	3:A:402:UNL:O2	1.53	1.21
1:A:7:VAL:HG11	1:B:34[B]:ARG:HH22	1.57	0.69
1:B:251[A]:ASP:HB3	1:B:254[A]:ARG:HD3	1.78	0.63
1:B:366:THR:HB	4:B:762:HOH:O	1.99	0.62
1:A:194[A]:THR:HG23	1:A:217:ARG:NH2	2.16	0.60
1:A:34[A]:ARG:HG3	1:B:40:ALA:HB1	1.88	0.56
1:A:182:ARG:NH2	4:A:501:HOH:O	2.25	0.54
1:A:7:VAL:HG12	1:A:7:VAL:O	2.08	0.53
1:B:183[A]:LYS:HE3	1:B:352:LEU:HD21	1.91	0.53
1:B:145:ALA:HB2	4:B:586[B]:HOH:O	2.11	0.49
1:A:40:ALA:HB1	1:B:34[B]:ARG:HG3	1.95	0.49
1:B:302:VAL:O	1:B:333:THR:HA	2.13	0.48
1:A:323:MET:HA	1:A:323:MET:HE2	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:PRO:HB2	1:B:314[B]:ARG:HG2	1.97	0.46
1:A:302:VAL:O	1:A:333:THR:HA	2.16	0.46
1:B:137:LEU:O	1:B:140[B]:VAL:HG22	2.15	0.45
1:B:323:MET:HA	1:B:323:MET:HE2	1.98	0.45
1:B:365:ALA:O	1:B:366:THR:HB	2.17	0.45
1:A:82:ASP:OD1	1:A:101[A]:TYR:OH	2.31	0.44
1:A:309[B]:LYS:O	1:A:309[B]:LYS:HG2	2.17	0.44
1:B:291:PRO:HA	4:B:501:HOH:O	2.16	0.44
1:B:139[B]:GLU:OE1	1:B:142:ARG:NH1	2.50	0.44
1:B:294:LEU:HD21	1:B:349:ILE:HD11	2.02	0.42
1:A:182:ARG:NE	4:A:501:HOH:O	2.35	0.42
1:A:271:HIS:ND1	1:A:300[B]:GLN:OE1	2.52	0.42
1:B:271:HIS:ND1	1:B:300[B]:GLN:OE1	2.50	0.42
1:A:82:ASP:OD1	1:A:101[B]:TYR:OH	2.23	0.41
1:B:159:GLN:HB2	1:B:225:VAL:HG21	2.03	0.41
1:A:308:PRO:HB2	1:A:314[B]:ARG:HG2	2.03	0.41
1:A:139[A]:GLU:OE2	1:A:143:HIS:NE2	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/384 (106%)	399 (98%)	6 (2%)	0	100	100
1	B	394/384 (103%)	386 (98%)	8 (2%)	0	100	100
All	All	799/768 (104%)	785 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/299 (107%)	321 (100%)	0	100	100
1	B	308/299 (103%)	308 (100%)	0	100	100
All	All	629/598 (105%)	629 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	159	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are unknown - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SAH	A	401	-	27,28,28	0.39	0	36,40,40	0.47	0
2	SAH	B	401	-	27,28,28	0.24	0	36,40,40	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	A	401	-	-	0/15/31/31	0/3/3/3
2	SAH	B	401	-	-	0/15/31/31	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	378/384 (98%)	-0.09	29 (7%)	19 27	6, 14, 35, 87	46 (12%)
1	B	361/384 (94%)	-0.10	18 (4%)	34 42	6, 15, 43, 114	35 (9%)
All	All	739/768 (96%)	-0.09	47 (6%)	25 34	6, 15, 40, 114	81 (10%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	187	SER	6.7
1	B	186	ALA	6.6
1	B	366	THR	6.0
1	A	368[A]	PRO	5.8
1	A	382[A]	PHE	5.1
1	B	291	PRO	4.1
1	A	148	THR	4.0
1	A	192[A]	ASP	4.0
1	B	254[A]	ARG	4.0
1	A	380[A]	LEU	3.8
1	A	95	ASP	3.7
1	B	16	VAL	3.7
1	A	96	THR	3.7
1	A	151	ALA	3.6
1	A	6	ASN	3.6
1	B	7	VAL	3.6
1	A	383[A]	GLN	3.6
1	B	188	GLY	3.5
1	A	194[A]	THR	3.5
1	A	372[A]	PRO	3.5
1	A	7	VAL	3.4
1	A	150	GLY	3.4
1	B	149	SER	3.3
1	B	59	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	150	GLY	3.1
1	B	148	THR	3.1
1	B	191	GLY	3.0
1	A	146	ARG	3.0
1	A	99	PHE	2.9
1	A	370[A]	ALA	2.7
1	A	253	TRP	2.7
1	B	192	ASP	2.6
1	B	193	ALA	2.5
1	A	94	HIS	2.5
1	A	149	SER	2.4
1	B	6	ASN	2.4
1	B	18	VAL	2.4
1	A	93	ILE	2.3
1	A	376[A]	ALA	2.3
1	A	98	GLY	2.2
1	A	145	ALA	2.2
1	A	193	ALA	2.2
1	A	366	THR	2.2
1	A	381[A]	TYR	2.1
1	A	191	GLY	2.1
1	B	189	ARG	2.1
1	A	367[A]	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	UNL	A	402	9/-	0.96	0.06	10,14,15,16	9
3	UNL	B	402	9/-	0.97	0.05	10,13,14,15	9
2	SAH	A	401	26/26	0.99	0.03	9,11,14,17	0
2	SAH	B	401	26/26	0.99	0.03	8,11,16,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.