



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:46 AM EDT

PDB ID : 2UY5 / pdb_00002uy5
Title : ScCTS1_kinetin crystal structure
Authors : Hurtado-Guerrero, R.; van Aalten, D.M.F.
Deposited on : 2007-04-02
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	12% 85% 8% . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDOCHITINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2179	1380	335	458	6	0	6	1

- Molecule 2 is N-(FURAN-2-YLMETHYL)-7H-PURIN-6-AMINE (CCD ID: H35) (formula: C₁₀H₉N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	16	10	5	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	192	Total	O	0	0
			192	192		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ENDOCHITINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.35Å 111.62Å 37.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.60 20.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.8 (20.00-1.60) 96.7 (20.00-1.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.222 0.217 , 0.247	Depositor DCC
R_{free} test set	825 reflections (2.07%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2387	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: H35

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	10/2245 (0.4%)	0.95	5/3057 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254	ALA	CA-CB	-7.66	1.40	1.53
1	A	262	ASP	N-CA	-6.31	1.38	1.46
1	A	255	ALA	CA-CB	-5.93	1.43	1.53
1	A	269	THR	CA-CB	-5.89	1.44	1.53
1	A	269	THR	CA-C	-5.75	1.45	1.52
1	A	267	GLU	CA-C	-5.54	1.46	1.52
1	A	266	LEU	CA-C	-5.41	1.46	1.52
1	A	271	ALA	C-N	5.37	1.41	1.33
1	A	267	GLU	N-CA	-5.15	1.40	1.46
1	A	255	ALA	C-O	-5.15	1.17	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	SER	N-CA-C	6.31	117.90	108.31
1	A	267	GLU	N-CA-C	-5.93	104.51	110.97
1	A	268	SER	N-CA-C	5.73	117.20	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	SER	CB-CA-C	-5.33	101.26	112.82
1	A	253	SER	N-CA-C	5.01	121.48	110.80

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	251	SER	Peptide
1	A	252	ALA	Peptide
1	A	260	ILE	Mainchain
1	A	263	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2179	0	2028	36	0
2	A	16	0	9	0	0
3	A	192	0	0	0	0
All	All	2387	0	2037	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:ASP:C	1:A:264:SER:HB3	1.49	1.34
1:A:262:ASP:HB3	1:A:264:SER:OG	1.34	1.26
1:A:262:ASP:HB3	1:A:264:SER:CB	1.65	1.25
1:A:265:LEU:H	1:A:265:LEU:HD12	0.95	1.05
1:A:262:ASP:C	1:A:264:SER:CB	2.29	1.04
1:A:262:ASP:CB	1:A:264:SER:CB	2.43	0.95
1:A:265:LEU:HD12	1:A:265:LEU:N	1.74	0.94
1:A:262:ASP:CA	1:A:264:SER:HB3	1.98	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:ASP:HB2	1:A:265:LEU:HD11	1.49	0.93
1:A:262:ASP:CB	1:A:264:SER:HB3	2.00	0.92
1:A:262:ASP:HB3	1:A:264:SER:HB3	1.51	0.86
1:A:262:ASP:O	1:A:264:SER:HB3	1.79	0.81
1:A:262:ASP:HB2	1:A:265:LEU:CD1	2.17	0.73
1:A:262:ASP:CA	1:A:264:SER:CB	2.64	0.69
1:A:265:LEU:H	1:A:265:LEU:CD1	1.79	0.65
1:A:263:THR:N	1:A:264:SER:CB	2.63	0.62
1:A:251:SER:OG	1:A:253:SER:HB3	2.04	0.56
1:A:265:LEU:N	1:A:265:LEU:CD1	2.52	0.53
1:A:117:SER:H	1:A:121:GLN:NE2	2.11	0.48
1:A:226:TRP:NE1	1:A:230:LEU:HD23	2.28	0.48
1:A:263:THR:O	1:A:267:GLU:HB2	2.14	0.46
1:A:265:LEU:O	1:A:269:THR:N	2.40	0.46
1:A:36:ASN:H	1:A:41:GLN:NE2	2.13	0.46
1:A:237:SER:O	1:A:240:LYS:HD3	2.16	0.46
1:A:262:ASP:CB	1:A:264:SER:OG	2.29	0.46
1:A:262:ASP:CB	1:A:265:LEU:CD1	2.91	0.46
1:A:262:ASP:CB	1:A:265:LEU:HD11	2.34	0.44
1:A:251:SER:OG	1:A:252:ALA:N	2.46	0.43
1:A:263:THR:N	1:A:264:SER:HB2	2.33	0.42
1:A:78:THR:HG22	1:A:84:LEU:HD12	2.01	0.42
1:A:269:THR:O	1:A:273:ILE:HG12	2.20	0.42
1:A:61:LEU:HD23	1:A:107:LEU:HG	2.02	0.41
1:A:64:PHE:HB2	1:A:65:PRO:HA	2.01	0.41
1:A:251:SER:OG	1:A:289:GLN:NE2	2.53	0.41
1:A:87:THR:HG22	1:A:145:ASP:OD2	2.22	0.40
1:A:252:ALA:N	1:A:253:SER:HB3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	288/294 (98%)	273 (95%)	14 (5%)	1 (0%)	36 20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	219	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	238/239 (100%)	233 (98%)	5 (2%)	47 23

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	84	LEU
1	A	230	LEU
1	A	265	LEU
1	A	289	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	41	GLN
1	A	121	GLN
1	A	289	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	H35	A	1313	-	18,18,18	1.69	4 (22%)	24,24,24	2.60	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H35	A	1313	-	-	0/5/5/5	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1313	H35	C4-N9	-4.46	1.34	1.38
2	A	1313	H35	CAP-N6	3.59	1.51	1.45
2	A	1313	H35	C5-N7	-2.63	1.33	1.37
2	A	1313	H35	CAP-CAK	2.13	1.53	1.48

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1313	H35	C4-N9-C8	5.83	107.55	103.84
2	A	1313	H35	C5-C4-N3	-5.28	119.62	124.21
2	A	1313	H35	N3-C4-N9	4.06	132.50	125.78
2	A	1313	H35	N3-C2-N1	-3.76	122.89	128.58
2	A	1313	H35	C5-N7-C8	3.33	109.05	106.27
2	A	1313	H35	C5-C4-N9	-3.25	107.87	110.46
2	A	1313	H35	N7-C8-N9	-3.11	109.13	113.87
2	A	1313	H35	C2-N3-C4	3.03	118.82	113.19
2	A	1313	H35	CAK-CAP-N6	-2.65	106.59	112.06
2	A	1313	H35	C6-C5-C4	2.01	119.89	118.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/294 (97%)	1.11	36 (12%) 8 7	16, 26, 36, 47	6 (2%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	GLY	6.0
1	A	311	ALA	4.7
1	A	139	ALA	4.6
1	A	262	ASP	4.1
1	A	265	LEU	3.9
1	A	274	ALA	3.5
1	A	271	ALA	3.3
1	A	27	THR	3.2
1	A	252	ALA	3.2
1	A	310	SER	3.1
1	A	259	TYR	3.0
1	A	275	SER	3.0
1	A	312	SER	3.0
1	A	80	SER	2.9
1	A	138	GLY	2.9
1	A	292	SER	2.9
1	A	309	THR	2.9
1	A	266	LEU	2.7
1	A	122	ALA	2.6
1	A	258	GLY	2.6
1	A	241	ASN	2.5
1	A	272	ASP	2.5
1	A	112	GLY	2.5
1	A	166	ALA	2.5
1	A	263	THR	2.5
1	A	51	SER	2.5
1	A	253	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	116	PHE	2.3
1	A	197	VAL	2.3
1	A	26	ASN	2.3
1	A	124	THR	2.2
1	A	136	GLY	2.2
1	A	307	LEU	2.2
1	A	120	SER	2.1
1	A	156	ILE	2.1
1	A	137	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	H35	A	1313	16/16	0.91	0.08	14,20,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.