



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:11 AM UTC

PDB ID : 2UZP / pdb_00002uzp
Title : Crystal structure of the C2 domain of human protein kinase C gamma.
Authors : Pike, A.C.W.; Amos, A.; Johansson, C.; Sobott, F.; Savitsky, P.; Berridge, G.;
Fedorov, O.; Umeano, C.; Gorrec, F.; Bunkoczi, G.; Debreczeni, J.; von Delft,
F.; Arrowsmith, C.H.; Edwards, A.; Weigelt, J.; Sundstrom, M.; Knapp, S.
Deposited on : 2007-04-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	
1	B	144	
1	C	144	

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 4078 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN KINASE C GAMMA TYPE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	8	0
			1203	757	216	226	4			
1	B	142	Total	C	N	O	S	0	3	0
			1139	722	200	213	4			
1	C	142	Total	C	N	O	S	0	2	0
			1143	725	200	214	4			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	1
			21	9	1	9	2		
2	B	1	Total	C	N	O	P	0	1
			21	9	1	9	2		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is COBALT (II) ION (CCD ID: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Co	0	0
			1	1		
3	B	1	Total	Co	0	0
			1	1		
3	C	1	Total	Co	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Ca	0	0
			3	3		
4	B	3	Total	Ca	0	0
			3	3		
4	C	3	Total	Ca	0	0
			3	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	145	Total O 145 145	0	0
7	B	115	Total O 115 115	0	0
7	C	132	Total O 132 132	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN KINASE C GAMMA TYPE

Chain A: 





- Molecule 1: PROTEIN KINASE C GAMMA TYPE

Chain B: 



- Molecule 1: PROTEIN KINASE C GAMMA TYPE

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	138.50Å 138.50Å 68.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.00) 100.0 (50.00-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.171 , 0.205 (Not available) , 0.269	Depositor DCC
R_{free} test set	2294 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.301	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4078	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CO, EDO, PLP, DSN, SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1239	0.73	0/1684
1	B	0.62	0/1173	0.71	0/1597
1	C	0.61	0/1173	0.70	0/1595
All	All	0.62	0/3585	0.71	0/4876

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	202[B]	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1203	0	1189	23	0
1	B	1139	0	1114	4	0
1	C	1143	0	1123	2	0
2	A	21	0	4	0	0
2	B	21	0	4	0	0
2	C	15	0	6	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
5	A	30	0	0	0	0
5	B	35	0	0	0	0
5	C	35	0	0	0	0
6	A	8	0	12	0	0
6	B	8	0	12	0	0
6	C	16	0	24	0	0
7	A	145	0	0	0	0
7	B	115	0	0	0	0
7	C	132	0	0	0	0
All	All	4078	0	3488	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205[A]:ARG:HB3	1:A:207[A]:LEU:HD11	1.26	1.14
1:A:205[A]:ARG:CB	1:A:207[A]:LEU:HD11	1.80	1.10
1:A:201[ILE]:HG22	1:A:203[B]:ASP:HB3	1.50	0.92
1:A:205[A]:ARG:HB2	1:A:207[A]:LEU:HD11	1.68	0.74
1:A:205[A]:ARG:C	1:A:207[A]:LEU:HG	2.17	0.69
1:A:203[B]:ASP:N	1:A:204[B]:PRO:HD2	2.11	0.66
1:A:205[A]:ARG:O	1:A:207[A]:LEU:HG	1.95	0.65
1:A:203[B]:ASP:N	1:A:204[B]:PRO:CD	2.66	0.58
1:A:199[LYS]:NZ	1:A:203[B]:ASP:HB2	2.19	0.57
1:A:203[B]:ASP:H	1:A:204[B]:PRO:CD	2.17	0.56
1:A:203[B]:ASP:N	1:A:203[B]:ASP:OD2	2.39	0.56
1:A:203[B]:ASP:H	1:A:204[B]:PRO:HD2	1.74	0.53
1:A:205[B]:ARG:NH2	1:A:207[B]:LEU:HD11	2.24	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:0:MET:HE3	1:B:288:VAL:HG13	1.92	0.51
1:A:201:ILE:HA	1:A:202[B]:PRO:C	2.37	0.49
1:A:203[B]:ASP:O	1:A:204[B]:PRO:C	2.56	0.49
1:A:203[B]:ASP:O	1:A:205[B]:ARG:N	2.46	0.48
1:A:205[A]:ARG:HB2	1:A:207[A]:LEU:HD12	1.95	0.48
1:A:199:LYS:HZ1	1:A:203[B]:ASP:HB2	1.79	0.47
1:A:170:ALA:HA	1:A:267:LEU:HD22	1.97	0.46
1:B:157:ARG:HD3	1:C:157:ARG:HD3	1.98	0.46
1:A:169:THR:OG1	1:A:172:GLU:OE2	2.30	0.45
1:A:0:MET:HE3	1:A:288:VAL:HG13	1.99	0.43
1:A:205[B]:ARG:HD3	1:A:207[B]:LEU:HD12	2.00	0.43
1:A:205[B]:ARG:O	1:A:206[B]:ASN:HB2	2.19	0.41
1:B:194:PRO:HA	1:B:245:TRP:O	2.21	0.41
1:C:194:PRO:HA	1:C:245:TRP:O	2.21	0.41
1:B:165:ILE:HG23	1:B:175:VAL:HG22	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/144 (103%)	139 (94%)	6 (4%)	3 (2%)	6	2
1	B	143/144 (99%)	140 (98%)	2 (1%)	1 (1%)	18	14
1	C	142/144 (99%)	138 (97%)	3 (2%)	1 (1%)	18	14
All	All	433/432 (100%)	417 (96%)	11 (2%)	5 (1%)	12	6

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	B	0	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	0	MET
1	A	203[A]	ASP
1	A	203[B]	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/127 (105%)	132 (99%)	1 (1%)	73	80
1	B	123/127 (97%)	122 (99%)	1 (1%)	73	80
1	C	124/127 (98%)	123 (99%)	1 (1%)	73	80
All	All	380/381 (100%)	377 (99%)	3 (1%)	73	80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	HIS
1	B	154	HIS
1	C	154	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	280	GLN
1	A	287	ASN
1	B	230	ASN
1	B	280	GLN
1	B	287	ASN
1	C	182	ASN
1	C	210	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 45 ligands modelled in this entry, 12 are monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	B	1294[B]	-	15,15,16	1.20	2 (13%)	21,22,23	1.15	0
5	SO4	B	1301	-	4,4,4	0.38	0	6,6,6	0.35	0
6	EDO	B	1307	-	3,3,3	0.39	0	2,2,2	0.44	0
2	PLP	A	1294[B]	-	15,15,16	1.19	2 (13%)	21,22,23	1.28	4 (19%)
5	SO4	C	1299	4	4,4,4	0.36	0	6,6,6	0.56	0
5	SO4	A	1301	-	4,4,4	0.31	0	6,6,6	0.81	0
5	SO4	B	1305	-	4,4,4	0.38	0	6,6,6	0.50	0
6	EDO	C	1308	-	3,3,3	0.35	0	2,2,2	0.52	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	C	1300	-	4,4,4	0.15	0	6,6,6	0.42	0
5	SO4	A	1303	-	4,4,4	0.20	0	6,6,6	0.46	0
5	SO4	B	1300	-	4,4,4	0.20	0	6,6,6	0.36	0
2	PLP	B	1294[A]	-	15,15,16	1.19	2 (13%)	21,22,23	1.06	0
2	PLP	A	1294[A]	-	15,15,16	1.18	2 (13%)	21,22,23	1.24	4 (19%)
5	SO4	C	1301	-	4,4,4	0.25	0	6,6,6	0.29	0
5	SO4	C	1302	-	4,4,4	0.27	0	6,6,6	0.57	0
6	EDO	C	1309	-	3,3,3	0.49	0	2,2,2	0.27	0
2	PLP	C	1294	1,3	15,15,16	1.05	2 (13%)	21,22,23	1.15	2 (9%)
6	EDO	B	1306	-	3,3,3	0.41	0	2,2,2	0.11	0
5	SO4	C	1303	-	4,4,4	0.24	0	6,6,6	0.15	0
5	SO4	B	1299	4	4,4,4	0.33	0	6,6,6	0.63	0
5	SO4	B	1303	-	4,4,4	0.28	0	6,6,6	0.40	0
5	SO4	A	1302	-	4,4,4	0.29	0	6,6,6	0.35	0
5	SO4	C	1304	-	4,4,4	0.22	0	6,6,6	0.53	0
5	SO4	C	1305	-	4,4,4	0.20	0	6,6,6	0.39	0
5	SO4	B	1302	-	4,4,4	0.30	0	6,6,6	0.25	0
5	SO4	B	1304	-	4,4,4	0.25	0	6,6,6	0.15	0
6	EDO	A	1305	-	3,3,3	0.33	0	2,2,2	0.47	0
6	EDO	C	1306	-	3,3,3	0.42	0	2,2,2	0.11	0
6	EDO	C	1307	-	3,3,3	0.58	0	2,2,2	0.11	0
5	SO4	A	1299	4	4,4,4	0.35	0	6,6,6	0.45	0
5	SO4	A	1300	-	4,4,4	0.23	0	6,6,6	0.17	0
6	EDO	A	1306	-	3,3,3	0.50	0	2,2,2	0.11	0
5	SO4	A	1304	-	4,4,4	0.23	0	6,6,6	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	C	1308	-	-	0/1/1/1	-
2	PLP	B	1294[B]	-	-	3/6/6/8	0/1/1/1
6	EDO	C	1309	-	-	1/1/1/1	-
2	PLP	C	1294	1,3	-	0/6/6/8	0/1/1/1
6	EDO	B	1306	-	-	0/1/1/1	-
6	EDO	B	1307	-	-	1/1/1/1	-
6	EDO	A	1305	-	-	1/1/1/1	-
6	EDO	C	1306	-	-	0/1/1/1	-
6	EDO	C	1307	-	-	0/1/1/1	-
2	PLP	A	1294[B]	-	-	3/6/6/8	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	1294[A]	-	-	0/6/6/8	0/1/1/1
6	EDO	A	1306	-	-	1/1/1/1	-
2	PLP	A	1294[A]	-	-	2/6/6/8	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1294[A]	PLP	C2-N1	3.06	1.39	1.33
2	A	1294[B]	PLP	C2-N1	3.06	1.39	1.33
2	C	1294	PLP	C2-N1	2.37	1.38	1.33
2	B	1294[A]	PLP	C2-N1	2.27	1.37	1.33
2	B	1294[B]	PLP	C2-N1	2.27	1.37	1.33
2	A	1294[A]	PLP	C6-N1	2.24	1.39	1.34
2	A	1294[B]	PLP	C6-N1	2.24	1.39	1.34
2	C	1294	PLP	C6-N1	2.22	1.38	1.34
2	B	1294[A]	PLP	C6-N1	2.15	1.38	1.34
2	B	1294[B]	PLP	C6-N1	2.15	1.38	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1294[A]	PLP	C2A-C2-C3	2.66	123.91	120.80
2	A	1294[B]	PLP	C2A-C2-C3	2.66	123.91	120.80
2	A	1294[A]	PLP	O3-C3-C2	2.38	122.52	117.58
2	A	1294[B]	PLP	O3-C3-C2	2.38	122.52	117.58
2	C	1294	PLP	C5-C6-N1	-2.26	120.16	123.83
2	A	1294[A]	PLP	C4A-C4-C3	-2.07	117.08	120.52
2	A	1294[B]	PLP	C4A-C4-C3	-2.07	117.08	120.52
2	C	1294	PLP	O3-C3-C2	2.06	121.85	117.58
2	A	1294[A]	PLP	C3-C2-N1	-2.04	118.39	120.96
2	A	1294[B]	PLP	C3-C2-N1	-2.04	118.39	120.96

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1294[A]	PLP	C4-C5-C5A-O4P
2	A	1294[B]	PLP	C5A-O4P-P-O1P
2	A	1294[B]	PLP	C5A-O4P-P-O3P
2	B	1294[B]	PLP	C5A-O4P-P-O1P
2	B	1294[B]	PLP	C5A-O4P-P-O2P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	1294[B]	PLP	C5A-O4P-P-O3P
6	B	1307	EDO	O1-C1-C2-O2
2	A	1294[A]	PLP	C6-C5-C5A-O4P
6	A	1305	EDO	O1-C1-C2-O2
2	A	1294[B]	PLP	C5A-O4P-P-O2P
6	C	1309	EDO	O1-C1-C2-O2
6	A	1306	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/144 (97%)	0.83	9 (6%) 25 24	18, 36, 46, 69	8 (5%)
1	B	141/144 (97%)	1.47	32 (22%) 2 2	23, 37, 51, 79	3 (2%)
1	C	141/144 (97%)	1.15	19 (13%) 7 6	23, 37, 50, 73	2 (1%)
All	All	423/432 (97%)	1.15	60 (14%) 6 5	18, 37, 50, 79	13 (3%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	293	ALA	7.6
1	A	204[A]	PRO	7.4
1	C	293	ALA	7.2
1	A	293	ALA	6.8
1	B	204	PRO	5.5
1	A	205[A]	ARG	4.0
1	A	207[A]	LEU	3.8
1	B	185	PRO	3.7
1	C	204	PRO	3.7
1	A	206[A]	ASN	3.6
1	B	205	ARG	3.6
1	B	188	PRO	3.4
1	B	236	VAL	3.3
1	C	191	LEU	3.1
1	A	203[A]	ASP	3.1
1	B	251	SER	3.1
1	B	286	TYR	3.1
1	C	206	ASN	3.1
1	A	292	ASP	3.0
1	C	292	ASP	2.9
1	C	205	ARG	2.9
1	B	191	LEU	2.8
1	C	252	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	292	ASP	2.8
1	B	271	VAL	2.8
1	B	184	ILE	2.8
1	B	166	ARG	2.7
1	B	252	ARG	2.7
1	B	248	ASP	2.7
1	B	250	THR	2.6
1	B	186	MET	2.6
1	B	206	ASN	2.6
1	C	188	PRO	2.6
1	B	245	TRP	2.4
1	C	250	THR	2.4
1	C	219	LEU	2.4
1	C	189	ASN	2.4
1	B	187	ASP	2.4
1	B	255	PHE	2.4
1	B	173	ILE	2.4
1	B	168	PRO	2.3
1	C	182	ASN	2.3
1	A	286	TYR	2.3
1	B	247	TRP	2.3
1	C	183	LEU	2.3
1	C	190	GLY	2.3
1	C	236	VAL	2.2
1	B	270	PRO	2.2
1	B	264	SER	2.2
1	B	190	GLY	2.1
1	B	266	LEU	2.1
1	B	189	ASN	2.1
1	A	166	ARG	2.1
1	C	251	SER	2.1
1	B	283	GLY	2.1
1	C	185	PRO	2.1
1	B	233	PRO	2.0
1	C	184	ILE	2.0
1	B	183	LEU	2.0
1	C	207	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	DSN	A	-1	6/7	0.84	0.18	38,42,46,54	0
1	DSN	B	-1	6/7	0.85	0.17	35,38,42,48	0
1	DSN	C	-1	6/7	0.87	0.16	36,39,42,48	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	B	1307	4/4	0.72	0.21	57,58,60,67	0
5	SO4	C	1303	5/5	0.75	0.18	70,75,83,86	0
5	SO4	A	1302	5/5	0.77	0.16	50,64,71,74	0
5	SO4	B	1304	5/5	0.78	0.14	78,78,86,86	0
5	SO4	B	1305	5/5	0.78	0.18	47,50,59,60	5
6	EDO	C	1309	4/4	0.82	0.22	37,57,63,64	0
5	SO4	C	1302	5/5	0.85	0.15	45,50,60,63	0
6	EDO	A	1305	4/4	0.85	0.19	55,55,59,66	0
6	EDO	C	1308	4/4	0.86	0.20	50,54,55,68	0
5	SO4	B	1302	5/5	0.87	0.12	35,54,55,62	0
2	PLP	C	1294	15/16	0.87	0.12	27,32,70,78	0
6	EDO	A	1306	4/4	0.87	0.20	30,57,59,59	0
5	SO4	B	1301	5/5	0.88	0.19	29,37,45,50	5
5	SO4	C	1305	5/5	0.89	0.19	34,41,49,52	5
5	SO4	A	1303	5/5	0.90	0.24	42,43,53,62	0
5	SO4	A	1304	5/5	0.90	0.18	26,41,49,53	5
5	SO4	A	1301	5/5	0.90	0.18	37,44,53,63	0
5	SO4	C	1301	5/5	0.90	0.13	38,58,61,67	0
2	PLP	A	1294[A]	15/16	0.92	0.11	23,33,43,47	6
2	PLP	A	1294[B]	15/16	0.92	0.11	23,34,45,58	6
2	PLP	B	1294[B]	15/16	0.93	0.11	26,35,44,57	6
2	PLP	B	1294[A]	15/16	0.93	0.11	29,34,42,42	6
5	SO4	C	1299	5/5	0.93	0.17	31,32,45,50	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	B	1303	5/5	0.93	0.18	34,37,45,51	5
6	EDO	C	1307	4/4	0.94	0.18	24,25,36,38	0
5	SO4	B	1300	5/5	0.94	0.14	46,47,51,52	0
5	SO4	C	1304	5/5	0.94	0.15	31,31,49,52	5
6	EDO	C	1306	4/4	0.96	0.11	24,32,37,40	0
6	EDO	B	1306	4/4	0.96	0.13	26,32,35,38	0
5	SO4	B	1299	5/5	0.97	0.13	24,29,32,34	0
5	SO4	A	1299	5/5	0.97	0.14	24,34,36,37	0
5	SO4	C	1300	5/5	0.97	0.13	30,37,38,44	0
5	SO4	A	1300	5/5	0.97	0.15	32,34,41,48	0
4	CA	C	1297	1/1	0.98	0.16	28,28,28,28	0
4	CA	C	1298	1/1	0.98	0.13	22,22,22,22	0
4	CA	A	1298	1/1	0.98	0.15	27,27,27,27	0
4	CA	B	1298	1/1	0.99	0.17	20,20,20,20	0
4	CA	C	1296	1/1	0.99	0.15	22,22,22,22	0
3	CO	C	1295	1/1	0.99	0.09	22,22,22,22	1
4	CA	A	1296	1/1	0.99	0.12	22,22,22,22	0
4	CA	A	1297	1/1	0.99	0.14	21,21,21,21	0
3	CO	B	1295	1/1	0.99	0.09	25,25,25,25	1
4	CA	B	1296	1/1	0.99	0.14	25,25,25,25	0
4	CA	B	1297	1/1	0.99	0.12	22,22,22,22	0
3	CO	A	1295	1/1	1.00	0.07	22,22,22,22	1

6.5 Other polymers

There are no such residues in this entry.