



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:45 AM UTC

PDB ID : 4V0N / pdb_00004v0n
Title : Crystal structure of BBS1N in complex with ARL6DN, soaked with mercury
Authors : Mourao, A.; Lorentzen, E.
Deposited on : 2014-09-17
Resolution : 3.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2351 (3.18-3.10)
Clashscore	190562	2452 (3.18-3.10)
Ramachandran outliers	187476	2324 (3.18-3.10)
Sidechain outliers	187428	2324 (3.18-3.10)
RSRZ outliers	180081	2351 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	78% 20%
1	C	169	68% 28%
1	E	169	65% 30%
1	G	169	69% 28%
2	B	425	50% 24% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	425	2%46%24%•28%
2	F	425	3%48%22%•28%
2	H	425	%46%26%•24%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARF-LIKE SMALL GTPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	166	Total	C	N	O	S	0	0	0
			1263	807	215	235	6			
1	C	166	Total	C	N	O	S	0	0	0
			1287	820	222	239	6			
1	E	166	Total	C	N	O	S	0	0	0
			1282	819	220	237	6			
1	G	166	Total	C	N	O	S	0	0	0
			1286	822	221	237	6			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	GLY	-	expression tag	UNP A8JF99
A	13	ALA	-	expression tag	UNP A8JF99
A	14	ALA	-	expression tag	UNP A8JF99
A	15	SER	-	expression tag	UNP A8JF99
C	12	GLY	-	expression tag	UNP A8JF99
C	13	ALA	-	expression tag	UNP A8JF99
C	14	ALA	-	expression tag	UNP A8JF99
C	15	SER	-	expression tag	UNP A8JF99
E	12	GLY	-	expression tag	UNP A8JF99
E	13	ALA	-	expression tag	UNP A8JF99
E	14	ALA	-	expression tag	UNP A8JF99
E	15	SER	-	expression tag	UNP A8JF99
G	12	GLY	-	expression tag	UNP A8JF99
G	13	ALA	-	expression tag	UNP A8JF99
G	14	ALA	-	expression tag	UNP A8JF99
G	15	SER	-	expression tag	UNP A8JF99


- Molecule 2 is a protein called BARDET-BIEDL SYNDROME 1 PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	330	Total	C	N	O	S	0	0	0
			2442	1561	423	445	13			
2	D	304	Total	C	N	O	S	0	0	0
			2197	1399	383	403	12			
2	F	304	Total	C	N	O	S	0	0	0
			2200	1405	379	403	13			
2	H	325	Total	C	N	O	S	0	0	0
			2386	1526	410	437	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	37	ARG	LYS	conflict	UNP A8JEA1
D	37	ARG	LYS	conflict	UNP A8JEA1
F	37	ARG	LYS	conflict	UNP A8JEA1
H	37	ARG	LYS	conflict	UNP A8JEA1

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	E	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	G	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Mg 1	0	0
4	C	1	Total 1	Mg 1	0	0
4	E	1	Total 1	Mg 1	0	0
4	G	1	Total 1	Mg 1	0	0


- Molecule 5 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Hg 1	0	0
5	B	13	Total 13	Hg 13	0	0
5	C	1	Total 1	Hg 1	0	0
5	D	12	Total 12	Hg 12	0	0
5	E	1	Total 1	Hg 1	0	0
5	F	14	Total 14	Hg 14	0	0
5	G	1	Total 1	Hg 1	0	0
5	H	14	Total 14	Hg 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ARF-LIKE SMALL GTPASE




- Molecule 1: ARF-LIKE SMALL GTPASE




- Molecule 1: ARF-LIKE SMALL GTPASE



- Molecule 1: ARF-LIKE SMALL GTPASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.75Å 123.75Å 443.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	96.50 – 3.13 96.50 – 3.13	Depositor EDS
% Data completeness (in resolution range)	99.0 (96.50-3.13) 99.2 (96.50-3.13)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 3.13Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.213 , 0.249 0.229 , 0.259	Depositor DCC
R_{free} test set	3506 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	104.1	Xtriage
Anisotropy	0.263	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14532	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, HG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1289	0.96	1/1755 (0.1%)
1	C	0.75	0/1313	1.09	3/1784 (0.2%)
1	E	0.78	0/1308	1.15	8/1777 (0.5%)
1	G	0.57	0/1312	0.98	2/1781 (0.1%)
2	B	0.71	0/2494	1.13	15/3412 (0.4%)
2	D	0.59	0/2240	1.00	7/3066 (0.2%)
2	F	0.59	0/2244	1.07	15/3071 (0.5%)
2	H	0.71	0/2437	1.14	9/3339 (0.3%)
All	All	0.66	0/14637	1.08	60/19985 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	H	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	144	ASN	N-CA-C	8.36	120.28	110.41
2	H	146	GLU	N-CA-C	8.35	122.85	109.24
2	B	341	LEU	CA-C-N	8.17	128.12	119.05
2	B	341	LEU	C-N-CA	8.17	128.12	119.05
2	H	145	THR	N-CA-C	7.49	119.42	108.86

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	226	ILE	Peptide
2	H	145	THR	Peptide
2	H	226	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1263	0	1241	24	0
1	C	1287	0	1285	33	0
1	E	1282	0	1283	37	0
1	G	1286	0	1294	35	0
2	B	2442	0	2400	74	0
2	D	2197	0	2087	68	0
2	F	2200	0	2111	59	0
2	H	2386	0	2324	85	0
3	A	32	0	12	2	0
3	C	32	0	12	3	0
3	E	32	0	12	2	0
3	G	32	0	12	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	G	1	0	0	0	0
5	A	1	0	0	0	0
5	B	13	0	0	0	0
5	C	1	0	0	0	0
5	D	12	0	0	0	0
5	E	1	0	0	0	0
5	F	14	0	0	0	0
5	G	1	0	0	0	0
5	H	14	0	0	0	0
All	All	14532	0	14073	391	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 391 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:320:GLY:HA3	2:H:348:MET:HE1	1.45	0.98
2:B:320:GLY:HA3	2:B:348:MET:HE1	1.49	0.94
2:H:43:ALA:HB2	2:H:417:MET:HE2	1.55	0.86
2:B:43:ALA:HB2	2:B:417:MET:HE2	1.57	0.85
2:F:320:GLY:HA3	2:F:348:MET:HE1	1.59	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/169 (97%)	155 (94%)	9 (6%)	0	100	100
1	C	164/169 (97%)	158 (96%)	6 (4%)	0	100	100
1	E	164/169 (97%)	155 (94%)	9 (6%)	0	100	100
1	G	164/169 (97%)	154 (94%)	10 (6%)	0	100	100
2	B	326/425 (77%)	290 (89%)	32 (10%)	4 (1%)	10	34
2	D	294/425 (69%)	269 (92%)	24 (8%)	1 (0%)	36	64
2	F	296/425 (70%)	271 (92%)	23 (8%)	2 (1%)	18	47
2	H	321/425 (76%)	288 (90%)	28 (9%)	5 (2%)	7	28
All	All	1893/2376 (80%)	1740 (92%)	141 (7%)	12 (1%)	21	50

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	65	GLY
2	B	113	LEU
2	B	130	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	65	GLY
2	D	65	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/142 (92%)	127 (97%)	4 (3%)	35	60
1	C	137/142 (96%)	129 (94%)	8 (6%)	18	44
1	E	136/142 (96%)	127 (93%)	9 (7%)	15	40
1	G	137/142 (96%)	134 (98%)	3 (2%)	45	67
2	B	250/352 (71%)	225 (90%)	25 (10%)	7	25
2	D	214/352 (61%)	198 (92%)	16 (8%)	12	36
2	F	218/352 (62%)	201 (92%)	17 (8%)	11	34
2	H	242/352 (69%)	210 (87%)	32 (13%)	4	15
All	All	1465/1976 (74%)	1351 (92%)	114 (8%)	11	34

5 of 114 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	58	VAL
2	H	377	LYS
2	F	308	VAL
2	H	360	LYS
2	H	266	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	257	ASN
2	F	69	ASN
2	H	59	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 65 ligands modelled in this entry, 61 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GTP	C	600	4	33,34,34	1.14	4 (12%)	50,54,54	1.73	11 (22%)
3	GTP	A	600	4	33,34,34	0.97	2 (6%)	50,54,54	1.55	9 (18%)
3	GTP	E	600	4	33,34,34	1.12	4 (12%)	50,54,54	1.63	11 (22%)
3	GTP	G	600	4	33,34,34	1.05	3 (9%)	50,54,54	1.54	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	C	600	4	-	7/22/38/38	0/3/3/3
3	GTP	A	600	4	-	3/22/38/38	0/3/3/3
3	GTP	E	600	4	-	4/22/38/38	0/3/3/3
3	GTP	G	600	4	-	1/22/38/38	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	600	GTP	PA-O3A	3.26	1.63	1.59
3	C	600	GTP	PB-O3A	2.89	1.62	1.59
3	G	600	GTP	PA-O3A	2.78	1.62	1.59
3	E	600	GTP	PB-O3A	2.74	1.62	1.59
3	C	600	GTP	PA-O3A	2.63	1.62	1.59

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	600	GTP	C2-N3-C4	4.86	120.67	112.30
3	C	600	GTP	C5-C4-N3	-4.58	121.11	128.39
3	G	600	GTP	C5-C4-N3	-4.44	121.33	128.39
3	A	600	GTP	C5-C4-N3	-4.43	121.34	128.39
3	G	600	GTP	C2-N3-C4	4.34	119.77	112.30

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	600	GTP	C5'-O5'-PA-O1A
3	C	600	GTP	O4'-C4'-C5'-O5'
3	G	600	GTP	PB-O3B-PG-O2G
3	C	600	GTP	PG-O3B-PB-O1B
3	A	600	GTP	PB-O3B-PG-O1G


There are no ring outliers.



3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	600	GTP	3	0
3	A	600	GTP	2	0
3	E	600	GTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	166/169 (98%)	-0.30	2 (1%) 76 57	37, 62, 88, 105	0
1	C	166/169 (98%)	-0.58	1 (0%) 85 70	19, 31, 56, 74	0
1	E	166/169 (98%)	-0.56	0 100 100	20, 35, 68, 83	0
1	G	166/169 (98%)	-0.47	0 100 100	32, 51, 70, 80	0
2	B	330/425 (77%)	-0.41	8 (2%) 59 38	21, 43, 77, 101	0
2	D	304/425 (71%)	-0.21	10 (3%) 49 28	23, 67, 106, 118	0
2	F	304/425 (71%)	-0.03	12 (3%) 43 23	23, 66, 110, 120	0
2	H	325/425 (76%)	-0.34	4 (1%) 76 57	24, 44, 79, 106	0
All	All	1927/2376 (81%)	-0.33	37 (1%) 66 45	19, 48, 101, 120	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	252	THR	5.4
2	F	104	TYR	5.3
2	H	147	GLU	4.5
2	D	226	ILE	4.1
2	F	140	PRO	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	HG	F	1434	1/1	0.42	0.20	113,113,113,113	1
5	HG	H	1434	1/1	0.68	0.26	79,79,79,79	1
5	HG	B	1432	1/1	0.79	0.26	69,69,69,69	1
5	HG	D	1436	1/1	0.82	0.15	100,100,100,100	1
5	HG	G	1181	1/1	0.88	0.11	50,50,50,50	1
5	HG	A	1181	1/1	0.90	0.12	54,54,54,54	1
5	HG	D	1433	1/1	0.93	0.12	82,82,82,82	1
5	HG	F	1435	1/1	0.93	0.20	80,80,80,80	1
5	HG	B	1437	1/1	0.93	0.10	39,39,39,39	1
5	HG	F	1431	1/1	0.93	0.17	108,108,108,108	1
5	HG	C	1181	1/1	0.95	0.17	53,53,53,53	1
3	GTP	G	600	32/32	0.95	0.06	40,44,47,50	0
5	HG	D	1434	1/1	0.95	0.09	80,80,80,80	1
3	GTP	A	600	32/32	0.95	0.06	55,60,67,72	0
5	HG	E	1181	1/1	0.95	0.15	51,51,51,51	1
5	HG	D	1435	1/1	0.96	0.15	109,109,109,109	1
5	HG	B	1426	1/1	0.96	0.09	38,38,38,38	1
4	MG	A	601	1/1	0.97	0.08	58,58,58,58	0
5	HG	F	1433	1/1	0.97	0.13	118,118,118,118	1
4	MG	C	601	1/1	0.97	0.12	27,27,27,27	0
5	HG	B	1435	1/1	0.97	0.13	40,40,40,40	1
4	MG	G	601	1/1	0.97	0.10	45,45,45,45	0
5	HG	H	1433	1/1	0.97	0.13	43,43,43,43	1
3	GTP	E	600	32/32	0.97	0.05	26,32,40,41	0
5	HG	F	1430	1/1	0.98	0.09	82,82,82,82	1
5	HG	B	1434	1/1	0.98	0.09	45,45,45,45	1
3	GTP	C	600	32/32	0.98	0.04	27,31,37,39	0
5	HG	B	1428	1/1	0.98	0.12	61,61,61,61	1
4	MG	E	601	1/1	0.98	0.06	31,31,31,31	0
5	HG	F	1436	1/1	0.98	0.07	73,73,73,73	1
5	HG	D	1437	1/1	0.98	0.12	87,87,87,87	1
5	HG	H	1429	1/1	0.98	0.11	68,68,68,68	1
5	HG	D	1427	1/1	0.98	0.11	85,85,85,85	1
5	HG	F	1428	1/1	0.98	0.10	102,102,102,102	1
5	HG	H	1435	1/1	0.98	0.12	42,42,42,42	1
5	HG	H	1436	1/1	0.98	0.11	46,46,46,46	1
5	HG	H	1439	1/1	0.98	0.10	41,41,41,41	1

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	HG	D	1429	1/1	0.99	0.09	82,82,82,82	1
5	HG	F	1432	1/1	0.99	0.09	73,73,73,73	1
5	HG	D	1430	1/1	0.99	0.10	98,98,98,98	1
5	HG	D	1431	1/1	0.99	0.10	109,109,109,109	1
5	HG	B	1431	1/1	0.99	0.12	43,43,43,43	1
5	HG	B	1436	1/1	0.99	0.08	51,51,51,51	1
5	HG	F	1437	1/1	0.99	0.12	122,122,122,122	1
5	HG	F	1438	1/1	0.99	0.07	76,76,76,76	1
5	HG	F	1439	1/1	0.99	0.12	68,68,68,68	1
5	HG	B	1429	1/1	0.99	0.09	64,64,64,64	1
5	HG	H	1427	1/1	0.99	0.06	41,41,41,41	1
5	HG	H	1428	1/1	0.99	0.08	49,49,49,49	1
5	HG	B	1438	1/1	0.99	0.06	23,23,23,23	1
5	HG	H	1430	1/1	0.99	0.08	62,62,62,62	1
5	HG	H	1431	1/1	0.99	0.09	43,43,43,43	1
5	HG	H	1432	1/1	0.99	0.10	41,41,41,41	1
5	HG	B	1433	1/1	0.99	0.11	48,48,48,48	1
5	HG	D	1426	1/1	0.99	0.10	63,63,63,63	1
5	HG	B	1430	1/1	0.99	0.10	45,45,45,45	1
5	HG	F	1429	1/1	0.99	0.10	78,78,78,78	1
5	HG	H	1437	1/1	0.99	0.09	30,30,30,30	1
5	HG	H	1438	1/1	0.99	0.08	60,60,60,60	1
5	HG	D	1428	1/1	0.99	0.09	96,96,96,96	1
5	HG	B	1427	1/1	1.00	0.09	42,42,42,42	1
5	HG	H	1426	1/1	1.00	0.11	41,41,41,41	1
5	HG	D	1432	1/1	1.00	0.10	62,62,62,62	1
5	HG	F	1426	1/1	1.00	0.10	57,57,57,57	1
5	HG	F	1427	1/1	1.00	0.10	66,66,66,66	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GTP G 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP A 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP E 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP C 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.