



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 08:54 AM EDT

PDB ID : 7V0D / pdb_00007v0d
Title : Crystal structure of halogenase CtcP from Kitasatospora aureofaciens
Authors : Hou, C.; Tsodikov, O.V.
Deposited on : 2022-05-10
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	
1	B	558	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	606	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetracycline 7-halogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4207	2649	758	781	19			
1	B	536	Total	C	N	O	S	0	0	0
			4269	2685	770	795	19			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A0A1E7MYN1
A	-1	PRO	-	expression tag	UNP A0A1E7MYN1
A	0	HIS	-	expression tag	UNP A0A1E7MYN1
B	-2	GLY	-	expression tag	UNP A0A1E7MYN1
B	-1	PRO	-	expression tag	UNP A0A1E7MYN1
B	0	HIS	-	expression tag	UNP A0A1E7MYN1

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	7	4		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is SULFATE ION (CCD ID: SO₄) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	155	Total	O	0	0
			155	155		
8	B	168	Total	O	0	0
			168	168		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tetracycline 7-halogenase



• Molecule 1: Tetracycline 7-halogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	98.59Å 106.03Å 178.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.60 35.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (35.00-2.60) 99.7 (35.00-2.60)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.186 , 0.219 0.190 , 0.222	Depositor DCC
R_{free} test set	2833 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtriage
Anisotropy	0.747	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8905	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, EPE, 1PE, GOL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.98	0/4323	1.39	0/5874
1	B	0.98	0/4385	1.40	0/5958
All	All	0.98	0/8708	1.40	0/11832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4207	0	4011	30	0
1	B	4269	0	4070	25	0
2	A	18	0	24	0	0
2	B	12	0	16	2	0
3	A	11	0	13	2	0
4	A	1	0	0	0	0
5	A	7	0	10	8	0
5	B	7	0	10	1	0
6	A	10	0	0	2	0
6	B	25	0	0	0	0
7	B	15	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	155	0	0	1	0
8	B	168	0	0	1	0
All	All	8905	0	8172	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:ILE:HB	5:A:606:PEG:H32	1.72	0.70
1:B:200:ARG:NH2	1:B:339:ALA:O	2.25	0.65
1:B:522:ILE:HG12	1:B:547:VAL:HG11	1.80	0.62
1:B:420:ALA:HB2	1:B:462:TYR:CE1	2.34	0.61
1:B:488:ARG:NH1	1:B:492:GLN:HE22	1.98	0.61
1:A:186:ALA:HB2	1:A:336:MET:CE	2.36	0.56
1:A:61:GLN:HB2	5:A:606:PEG:H22	1.87	0.56
1:B:336:MET:HE3	1:B:361:LEU:HD23	1.88	0.56
1:B:250:ARG:HG3	1:B:279:ARG:HG3	1.88	0.55
1:A:250:ARG:HG3	1:A:279:ARG:HD3	1.91	0.53
1:A:59:ILE:HB	5:A:606:PEG:C3	2.39	0.52
1:B:420:ALA:HB2	1:B:462:TYR:CZ	2.45	0.52
1:B:42:MET:HE1	1:B:153:ARG:HG3	1.92	0.52
1:B:45:ASP:HA	2:B:604:GOL:H31	1.91	0.51
1:B:537:ARG:O	1:B:541:ALA:HB3	2.10	0.51
1:B:30:LEU:HD22	1:B:336:MET:HE1	1.93	0.50
1:A:36:LYS:HG3	1:A:148:TYR:CE1	2.47	0.50
1:A:250:ARG:HG3	1:A:279:ARG:CD	2.43	0.49
1:A:61:GLN:HB2	5:A:606:PEG:C2	2.42	0.49
1:A:421:SER:OG	5:A:606:PEG:H41	2.13	0.48
1:B:522:ILE:CG1	1:B:547:VAL:HG11	2.44	0.48
1:B:332:ARG:NH2	8:B:701:HOH:O	2.46	0.48
1:B:183:LEU:O	1:B:333:TRP:HA	2.14	0.48
1:A:421:SER:HB2	5:A:606:PEG:H41	1.96	0.47
1:B:84:PHE:CZ	1:B:88:LEU:HD11	2.49	0.47
1:A:491:GLU:OE2	1:A:495:ARG:NH1	2.48	0.47
1:A:421:SER:CB	5:A:606:PEG:H41	2.44	0.47
1:A:332:ARG:NH2	8:A:701:HOH:O	2.47	0.47
1:A:421:SER:HA	5:A:606:PEG:H11	1.96	0.47
1:A:413:VAL:HG22	1:A:469:LEU:HD11	1.96	0.46
1:A:239:TRP:CE3	1:A:244:MET:HE1	2.52	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:506:PHE:CZ	1:A:514:ILE:HG21	2.51	0.45
1:A:289:GLU:OE2	3:A:604:1PE:C25	2.65	0.45
1:A:102:GLY:O	1:A:245:HIS:HD2	2.01	0.44
1:B:252:TRP:CZ3	1:B:254:TRP:HB3	2.52	0.44
1:A:183:LEU:O	1:A:333:TRP:HA	2.18	0.44
1:B:36:LYS:HE2	1:B:74:GLY:O	2.18	0.44
1:A:131:HIS:CG	1:A:256:ILE:HG23	2.53	0.44
1:A:452:TYR:CE1	1:A:463:LYS:HD2	2.53	0.44
1:A:26:SER:HB2	6:A:608:SO4:O4	2.18	0.43
1:A:20:ILE:HB	1:A:43:LEU:HD23	2.00	0.43
1:B:318:THR:HB	5:B:601:PEG:H22	2.01	0.43
1:A:311:SER:H	3:A:604:1PE:C25	2.31	0.43
1:B:336:MET:HE3	1:B:361:LEU:CD2	2.49	0.43
1:A:165:ASP:O	1:A:332:ARG:NH1	2.52	0.42
1:A:500:MET:O	1:A:501:LEU:C	2.62	0.42
1:B:259:ASN:HB3	1:B:266:ASN:O	2.19	0.42
1:B:289:GLU:OE2	2:B:602:GOL:H11	2.19	0.42
1:A:69:ILE:HD11	1:A:355:CYS:O	2.21	0.41
1:B:134:ARG:HA	1:B:137:THR:HB	2.03	0.41
1:B:212:SER:HB2	1:B:321:MET:SD	2.61	0.41
1:B:413:VAL:HG22	1:B:469:LEU:HD11	2.02	0.41
1:A:245:HIS:HE1	6:A:607:SO4:O4	2.04	0.41
1:B:407:TYR:HB3	1:B:408:PRO:HD3	2.03	0.40
1:A:202:GLU:HA	1:A:203:PRO:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	525/558 (94%)	503 (96%)	22 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	534/558 (96%)	514 (96%)	20 (4%)	0	100	100
All	All	1059/1116 (95%)	1017 (96%)	42 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/470 (95%)	439 (98%)	7 (2%)	55	79
1	B	451/470 (96%)	446 (99%)	5 (1%)	65	84
All	All	897/940 (95%)	885 (99%)	12 (1%)	61	82

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ARG
1	A	57	SER
1	A	63	LEU
1	A	92	SER
1	A	287	THR
1	A	336	MET
1	A	349	ARG
1	B	137	THR
1	B	312	VAL
1	B	501	LEU
1	B	504	LEU
1	B	547	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	ASN
1	A	143	HIS
1	A	245	HIS
1	A	450	ASN
1	B	115	ASN
1	B	128	ASN
1	B	160	ASN
1	B	245	HIS
1	B	406	HIS
1	B	443	HIS
1	B	492	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 1 is monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	602	-	5,5,5	0.08	0	5,5,5	0.24	0
2	GOL	A	601	-	5,5,5	0.09	0	5,5,5	0.25	0
5	PEG	B	601	-	6,6,6	0.14	0	5,5,5	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	B	605	-	4,4,4	0.34	0	6,6,6	0.12	0
7	EPE	B	603	-	15,15,15	1.88	1 (6%)	19,20,20	1.11	2 (10%)
6	SO4	A	608	-	4,4,4	0.36	0	6,6,6	0.05	0
6	SO4	A	607	-	4,4,4	0.34	0	6,6,6	0.11	0
2	GOL	A	603	-	5,5,5	0.08	0	5,5,5	0.25	0
3	1PE	A	604	-	10,10,15	0.54	0	9,9,14	0.29	0
6	SO4	B	607	-	4,4,4	0.33	0	6,6,6	0.08	0
5	PEG	A	606	-	6,6,6	0.16	0	5,5,5	0.14	0
6	SO4	B	606	-	4,4,4	0.34	0	6,6,6	0.07	0
6	SO4	B	609	-	4,4,4	0.34	0	6,6,6	0.07	0
6	SO4	B	608	-	4,4,4	0.33	0	6,6,6	0.07	0
2	GOL	B	602	-	5,5,5	0.10	0	5,5,5	0.29	0
2	GOL	B	604	-	5,5,5	0.10	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	602	-	-	2/4/4/4	-
2	GOL	A	601	-	-	2/4/4/4	-
5	PEG	B	601	-	-	3/4/4/4	-
7	EPE	B	603	-	-	4/9/19/19	0/1/1/1
2	GOL	A	603	-	-	2/4/4/4	-
3	1PE	A	604	-	-	4/8/8/13	-
5	PEG	A	606	-	-	2/4/4/4	-
2	GOL	B	602	-	-	2/4/4/4	-
2	GOL	B	604	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	603	EPE	C10-S	-6.95	1.67	1.77

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	603	EPE	O1S-S-C10	3.05	111.33	106.73
7	B	603	EPE	O2S-S-C10	2.10	109.91	106.73

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	602	GOL	O1-C1-C2-C3
2	A	603	GOL	O1-C1-C2-C3
2	B	602	GOL	C1-C2-C3-O3
7	B	603	EPE	S-C10-C9-N1
3	A	604	1PE	OH5-C14-C24-OH4
3	A	604	1PE	OH4-C13-C23-OH3
7	B	603	EPE	N4-C7-C8-O8
5	B	601	PEG	O1-C1-C2-O2
2	A	601	GOL	O1-C1-C2-C3
2	B	604	GOL	C1-C2-C3-O3
5	B	601	PEG	O2-C3-C4-O4
2	A	602	GOL	O1-C1-C2-O2
2	A	603	GOL	O1-C1-C2-O2
2	A	601	GOL	O1-C1-C2-O2
2	B	602	GOL	O2-C2-C3-O3
7	B	603	EPE	C10-C9-N1-C2
7	B	603	EPE	C10-C9-N1-C6
5	A	606	PEG	O2-C3-C4-O4
5	B	601	PEG	C1-C2-O2-C3
5	A	606	PEG	O1-C1-C2-O2
3	A	604	1PE	C12-C22-OH3-C23
3	A	604	1PE	C14-C24-OH4-C13
2	B	604	GOL	O2-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	601	PEG	1	0
6	A	608	SO4	1	0
6	A	607	SO4	1	0
3	A	604	1PE	2	0
5	A	606	PEG	8	0
2	B	602	GOL	1	0
2	B	604	GOL	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/558 (94%)	-0.12	11 (2%) 63 58	28, 42, 67, 107	0
1	B	536/558 (96%)	-0.02	15 (2%) 55 49	28, 44, 67, 130	0
All	All	1063/1116 (95%)	-0.07	26 (2%) 59 54	28, 43, 67, 130	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	550	ALA	5.8
1	A	436	LYS	4.4
1	B	547	VAL	4.2
1	B	549	ALA	4.2
1	A	439	GLY	4.1
1	B	523	ARG	3.7
1	B	542	ALA	3.4
1	B	543	SER	3.3
1	B	319	ASP	3.3
1	A	15	PRO	3.2
1	B	544	ALA	3.1
1	A	437	GLU	2.8
1	B	228	ASP	2.8
1	B	202	GLU	2.8
1	A	541	ALA	2.7
1	B	203	PRO	2.6
1	B	488	ARG	2.5
1	A	279	ARG	2.4
1	A	442	SER	2.4
1	A	488	ARG	2.3
1	A	202	GLU	2.2
1	A	71	ASP	2.2
1	B	15	PRO	2.2
1	B	518	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	441	ASP	2.0
1	A	518	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	602	6/6	0.83	0.18	49,51,52,55	0
5	PEG	A	606	7/7	0.83	0.26	45,52,59,60	0
6	SO4	B	609	5/5	0.84	0.18	88,89,93,93	0
5	PEG	B	601	7/7	0.86	0.20	61,62,64,65	0
3	1PE	A	604	11/16	0.86	0.18	27,45,52,53	0
2	GOL	B	604	6/6	0.87	0.17	53,55,56,57	0
7	EPE	B	603	15/15	0.87	0.21	68,75,91,93	0
2	GOL	A	601	6/6	0.88	0.17	59,62,64,64	0
6	SO4	B	608	5/5	0.89	0.16	85,88,88,90	0
6	SO4	B	607	5/5	0.92	0.16	92,94,95,96	0
4	CL	A	605	1/1	0.94	0.11	53,53,53,53	0
2	GOL	A	602	6/6	0.95	0.13	56,57,57,57	0
2	GOL	A	603	6/6	0.95	0.11	54,55,56,57	0
6	SO4	A	607	5/5	0.95	0.16	54,56,56,59	0
6	SO4	A	608	5/5	0.95	0.12	62,64,65,66	0
6	SO4	B	605	5/5	0.96	0.13	56,58,59,60	0
6	SO4	B	606	5/5	0.96	0.10	61,61,63,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.