



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:14 PM UTC

PDB ID : 4V1U / pdb_00004v1u
Title : Heterocyclase in complex with substrate and Cofactor
Authors : Koehnke, J.; Naismith, J.H.
Deposited on : 2014-10-02
Resolution : 2.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	775	2% 87% 9% ••
1	B	775	2% 88% 7% •
2	C	64	12% 6% 81%
2	D	64	17% • 81%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYND.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	746	Total	C	N	O	S	0	0	0
			5864	3734	1014	1100	16			
1	B	743	Total	C	N	O	S	0	0	0
			5844	3721	1011	1096	16			

- Molecule 2 is a protein called PATE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	0	0	0
			88	53	13	22			
2	D	12	Total	C	N	O	0	0	0
			88	53	13	22			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		
5	B	2	Total	Ca	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	12	Total	O	0	0
			12	12		
6	B	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: LYND



• Molecule 1: LYND



• Molecule 2: PATE



• Molecule 2: PATE



MET	ASP	LYS	ASN	ILE	LEU	PRO	GLN	GLN	GLY	GLN	PRO	VAL	ILE	ARG	LEU	THR	ALA	GLY	GLN	LEU	S23	E31	L34	GLY	ASP	ALA	GLY	LEU	GLU	ALA	SER	LYS	ILE	THR	ALA	CYS	ILE	THR	PHE	CYS	ALA	TYR	ASP	GLY	GLU	LEU	GLU	HIS	HIS	HIS	HIS	HIS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.80Å 152.81Å 182.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.40 – 2.86 76.40 – 2.86	Depositor EDS
% Data completeness (in resolution range)	96.9 (76.40-2.86) 96.9 (76.40-2.86)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.86Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.201 , 0.251 0.201 , 0.249	Depositor DCC
R_{free} test set	2131 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	60.5	Xtriage
Anisotropy	0.566	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11955	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/6005	0.70	1/8191 (0.0%)
1	B	0.40	0/5982	0.68	0/8158
2	C	0.32	0/87	0.55	0/116
2	D	0.32	0/87	0.61	0/116
All	All	0.41	0/12161	0.69	1/16581 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	752	VAL	CB-CA-C	-5.17	108.87	114.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5864	0	5819	45	1
1	B	5844	0	5802	51	1
2	C	88	0	83	3	0
2	D	88	0	83	8	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	23	0	12	0	0
4	B	23	0	12	1	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	12	0	0	1	0
6	B	7	0	0	0	0
All	All	11955	0	11811	93	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:LYS:HE2	2:D:31:GLU:CB	1.48	1.44
1:B:78:LYS:CE	2:D:31:GLU:HB2	1.63	1.29
1:B:78:LYS:HE2	2:D:31:GLU:HB3	1.25	1.15
1:B:78:LYS:HE2	2:D:31:GLU:HB2	1.15	1.05
1:B:78:LYS:CE	2:D:31:GLU:CB	2.31	0.99
1:B:78:LYS:NZ	2:D:31:GLU:HB2	1.79	0.95
1:A:372:ILE:HD13	1:A:384:ARG:HE	1.33	0.93
1:A:372:ILE:CD1	1:A:384:ARG:HE	1.84	0.89
1:A:216:GLU:CD	1:A:390:GLY:O	2.18	0.86
1:A:216:GLU:OE1	1:A:390:GLY:O	1.94	0.85
1:B:335:LYS:NZ	1:B:694:SER:O	2.11	0.83
1:A:472:GLU:OE2	6:A:2009:HOH:O	1.98	0.79
1:B:78:LYS:HD3	1:B:80:TYR:CE2	2.20	0.77
1:B:638:VAL:O	1:B:642:ASN:ND2	2.20	0.74
1:B:434:GLY:O	1:B:438:ARG:NH2	2.23	0.72
1:B:399:ARG:NH2	2:C:32:GLU:OE2	2.23	0.71
1:B:161:TYR:CE2	1:B:182:LEU:HD23	2.24	0.71
1:B:56:VAL:O	1:B:60:ASP:N	2.24	0.70
1:A:738:ARG:NH2	1:A:745:GLY:O	2.27	0.67
1:B:654:GLU:N	1:B:654:GLU:OE1	2.31	0.64
1:A:226:GLN:N	1:A:226:GLN:OE1	2.32	0.63
1:B:78:LYS:CE	1:B:80:TYR:CE2	2.82	0.62
1:A:14:HIS:NE2	1:A:16:GLU:OE1	2.31	0.62
1:A:678:ASP:OD2	1:A:681:GLN:NE2	2.33	0.61
1:B:78:LYS:CE	1:B:80:TYR:HE2	2.14	0.61
1:B:562:ARG:NH2	1:B:767:GLN:O	2.33	0.61
1:A:22:GLN:NE2	1:B:395:LEU:HD22	2.17	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:SER:HG	1:A:126:THR:HG1	1.28	0.60
1:A:619:ARG:NH2	1:A:653:ASP:OD1	2.34	0.59
1:B:78:LYS:CD	1:B:80:TYR:CE2	2.85	0.58
1:A:372:ILE:HD13	1:A:384:ARG:NE	2.13	0.58
1:B:78:LYS:HZ3	2:D:31:GLU:HB2	1.68	0.57
1:B:368:GLU:N	1:B:368:GLU:OE1	2.38	0.56
1:B:161:TYR:CE2	1:B:182:LEU:CD2	2.88	0.56
1:B:161:TYR:CZ	1:B:182:LEU:HG	2.40	0.56
1:A:372:ILE:HD12	1:A:384:ARG:HE	1.70	0.54
1:A:22:GLN:HE22	1:B:395:LEU:HD22	1.73	0.53
1:B:287:ASP:OD1	1:B:287:ASP:N	2.42	0.52
1:A:303:HIS:ND1	1:A:360:SER:OG	2.36	0.52
1:A:216:GLU:OE1	1:A:391:SER:HA	2.12	0.50
1:B:409:LYS:NZ	4:B:777:AMP:P	2.85	0.50
1:B:415:GLN:NE2	1:B:415:GLN:O	2.45	0.50
1:A:656:LEU:HB3	1:A:661:THR:HG22	1.94	0.49
1:A:254:GLN:HB3	1:B:258:GLN:HG2	1.94	0.49
1:B:78:LYS:HD3	1:B:80:TYR:HE2	1.75	0.49
1:A:662:ASP:O	1:A:666:THR:OG1	2.23	0.49
1:A:123:SER:OG	1:A:126:THR:OG1	2.09	0.49
1:A:389:PHE:CZ	1:A:427:ARG:NH2	2.79	0.48
1:A:184:LYS:HB3	1:A:191:TRP:HB2	1.96	0.48
1:B:45:LEU:HB2	1:B:55:ILE:HD11	1.95	0.48
1:B:597:LEU:HD11	1:B:607:ALA:HB2	1.96	0.47
1:A:567:ALA:HB2	1:A:677:PRO:HB3	1.96	0.47
1:A:303:HIS:HB3	1:A:363:THR:HG22	1.96	0.47
1:B:78:LYS:HE2	1:B:80:TYR:HE2	1.79	0.46
1:B:78:LYS:CD	1:B:80:TYR:HE2	2.29	0.46
1:A:298:LEU:HD21	1:B:291:ILE:HG13	1.98	0.46
1:A:724:ASP:OD1	1:A:724:ASP:N	2.49	0.46
1:B:629:ASP:OD1	1:B:631:THR:OG1	2.33	0.45
1:A:702:MET:HA	1:A:705:VAL:HG12	1.98	0.45
1:B:184:LYS:HB3	1:B:191:TRP:HB2	2.00	0.44
1:B:39:TYR:OH	2:D:31:GLU:OE2	2.33	0.44
2:C:31:GLU:OE1	2:C:31:GLU:N	2.51	0.44
1:A:298:LEU:HD11	1:B:259:PHE:CZ	2.53	0.44
1:B:161:TYR:CD2	1:B:182:LEU:CD2	3.00	0.44
1:B:78:LYS:HD3	1:B:80:TYR:CD2	2.52	0.43
1:B:161:TYR:CG	1:B:182:LEU:HD21	2.53	0.43
1:A:6:LEU:HD23	1:A:51:THR:HG23	2.01	0.43
1:B:255:THR:O	1:B:259:PHE:HB2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:562:ARG:HG2	1:B:674:TYR:CD1	2.54	0.42
1:A:500:THR:HG22	1:A:705:VAL:CG2	2.49	0.42
1:B:436:GLU:OE1	1:B:436:GLU:N	2.50	0.42
1:B:590:ARG:NE	1:B:649:ASP:OD2	2.47	0.42
1:A:491:ILE:CD1	1:A:508:PRO:HB3	2.50	0.42
1:A:739:HIS:CD2	1:A:739:HIS:N	2.87	0.42
1:A:141:VAL:HG12	1:A:142:GLN:N	2.34	0.42
1:B:528:ASP:O	1:B:731:LYS:NZ	2.53	0.42
1:B:115:THR:HG23	1:B:151:THR:HG21	2.02	0.42
1:A:59:LEU:HD11	1:A:68:ILE:HD11	2.02	0.42
1:A:373:THR:HG21	1:A:381:HIS:HA	2.03	0.41
1:A:423:GLU:O	1:A:427:ARG:HG2	2.19	0.41
1:B:718:ILE:CG2	1:B:731:LYS:HB3	2.50	0.41
1:B:161:TYR:CD2	1:B:182:LEU:HD21	2.55	0.41
1:A:101:ILE:HD13	1:A:269:VAL:CG1	2.51	0.41
1:A:389:PHE:CE1	1:A:427:ARG:NH2	2.89	0.41
1:A:671:ASP:HB2	1:A:672:HIS:CD2	2.56	0.40
1:B:371:ARG:HG2	1:B:383:TYR:CZ	2.57	0.40
1:A:389:PHE:CE2	1:A:515:HIS:HE1	2.39	0.40
1:A:560:TYR:HA	1:A:769:ASN:HB2	2.03	0.40
1:B:78:LYS:HE3	1:B:80:TYR:CE2	2.56	0.40
2:C:23:SER:OG	2:C:24:SER:N	2.53	0.40
1:A:277:ALA:O	1:A:280:THR:OG1	2.39	0.40
1:A:498:SER:O	1:A:502:GLN:N	2.52	0.40
1:A:500:THR:HG22	1:A:705:VAL:HG22	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:TYR:OH	1:B:316:ASP:OD2[1_455]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	738/775 (95%)	709 (96%)	25 (3%)	4 (0%)	24	42
1	B	731/775 (94%)	704 (96%)	25 (3%)	2 (0%)	36	54
2	C	10/64 (16%)	10 (100%)	0	0	100	100
2	D	10/64 (16%)	9 (90%)	1 (10%)	0	100	100
All	All	1489/1678 (89%)	1432 (96%)	51 (3%)	6 (0%)	30	48

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	276	THR
1	A	224	GLN
1	A	280	THR
1	B	20	PRO
1	B	604	PRO
1	A	20	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	636/661 (96%)	624 (98%)	12 (2%)	50	72
1	B	634/661 (96%)	621 (98%)	13 (2%)	47	70
2	C	10/52 (19%)	10 (100%)	0	100	100
2	D	10/52 (19%)	10 (100%)	0	100	100
All	All	1290/1426 (90%)	1265 (98%)	25 (2%)	50	72

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	ILE
1	A	122	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	123	SER
1	A	217	VAL
1	A	224	GLN
1	A	373	THR
1	A	539	GLU
1	A	551	GLU
1	A	552	ARG
1	A	661	THR
1	A	739	HIS
1	A	748	ARG
1	B	111	ARG
1	B	119	VAL
1	B	188	SER
1	B	289	LYS
1	B	316	ASP
1	B	319	ILE
1	B	346	THR
1	B	373	THR
1	B	438	ARG
1	B	467	ARG
1	B	539	GLU
1	B	739	HIS
1	B	740	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	HIS
1	A	22	GLN
1	A	154	ASN
1	A	172	GLN
1	A	209	HIS
1	A	258	GLN
1	A	295	HIS
1	A	356	GLN
1	A	381	HIS
1	A	478	HIS
1	A	504	HIS
1	A	515	HIS
1	A	588	ASN
1	A	611	ASN
1	B	10	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	32	HIS
1	B	154	ASN
1	B	254	GLN
1	B	258	GLN
1	B	272	HIS
1	B	336	GLN
1	B	343	HIS
1	B	353	GLN
1	B	387	HIS
1	B	504	HIS
1	B	706	ASN
1	B	720	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	AMP	B	777	-	25,25,25	1.40	4 (16%)	37,38,38	1.93	11 (29%)
4	AMP	A	777	-	25,25,25	1.42	4 (16%)	37,38,38	2.02	11 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	B	777	-	-	0/10/26/26	0/3/3/3
4	AMP	A	777	-	-	0/10/26/26	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	777	AMP	C5-C4	4.62	1.47	1.39
4	B	777	AMP	C5-C4	4.58	1.47	1.39
4	A	777	AMP	C5-C6	2.74	1.48	1.41
4	B	777	AMP	C5-C6	2.51	1.48	1.41
4	A	777	AMP	C8-N7	2.43	1.36	1.31
4	B	777	AMP	C5-N7	-2.38	1.34	1.39
4	B	777	AMP	C8-N7	2.31	1.36	1.31
4	A	777	AMP	C5-N7	-2.00	1.35	1.39

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	777	AMP	C5-C4-N3	-5.42	119.25	126.72
4	A	777	AMP	C5-C4-N3	-5.25	119.48	126.72
4	B	777	AMP	N3-C4-N9	4.50	134.82	127.17
4	A	777	AMP	N3-C4-N9	4.50	134.81	127.17
4	A	777	AMP	N3-C2-N1	-3.99	122.54	128.58
4	A	777	AMP	C2-N3-C4	3.80	121.11	111.83
4	A	777	AMP	C4-N9-C8	3.79	109.71	105.74
4	B	777	AMP	N3-C2-N1	-3.69	122.99	128.58
4	B	777	AMP	C2-N3-C4	3.62	120.68	111.83
4	A	777	AMP	C4-C5-N7	-3.40	106.69	110.58
4	B	777	AMP	C4-C5-N7	-3.31	106.80	110.58
4	B	777	AMP	C4-N9-C8	3.18	109.08	105.74
4	B	777	AMP	C5-N7-C8	2.70	107.69	103.45
4	A	777	AMP	C5-N7-C8	2.68	107.66	103.45
4	A	777	AMP	N9-C8-N7	-2.61	110.24	113.94
4	A	777	AMP	O3P-P-O5'	-2.45	100.29	106.67
4	A	777	AMP	C6-C5-N7	2.44	136.79	132.09
4	B	777	AMP	N9-C8-N7	-2.42	110.50	113.94
4	A	777	AMP	C2-N1-C6	2.29	122.49	118.73
4	B	777	AMP	C2-N1-C6	2.17	122.30	118.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	777	AMP	O3P-P-O2P	2.09	115.65	107.80
4	B	777	AMP	C6-C5-N7	2.04	136.02	132.09

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	777	AMP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	746/775 (96%)	-0.09	12 (1%) 70 64	36, 61, 116, 152	0
1	B	743/775 (95%)	0.03	12 (1%) 70 64	41, 76, 125, 155	0
2	C	12/64 (18%)	0.39	0 100 100	67, 85, 104, 105	0
2	D	12/64 (18%)	0.48	0 100 100	90, 113, 137, 145	0
All	All	1513/1678 (90%)	-0.02	24 (1%) 70 64	36, 69, 122, 155	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	387	HIS	4.1
1	B	387	HIS	3.6
1	B	216	GLU	3.2
1	A	476	VAL	3.2
1	B	297	ILE	3.0
1	A	718	ILE	2.8
1	B	478	HIS	2.8
1	B	402	LEU	2.7
1	A	228	GLN	2.7
1	B	143	THR	2.6
1	B	384	ARG	2.4
1	A	276	THR	2.4
1	B	244	THR	2.4
1	A	646	LEU	2.4
1	B	371	ARG	2.3
1	A	388	SER	2.3
1	B	658	SER	2.3
1	A	390	GLY	2.3
1	B	379	LEU	2.2
1	B	390	GLY	2.2
1	A	6	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	478	HIS	2.2
1	A	391	SER	2.2
1	A	19	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	A	1776	1/1	0.79	0.17	107,107,107,107	0
5	CA	A	1777	1/1	0.85	0.17	109,109,109,109	0
5	CA	B	1776	1/1	0.87	0.16	123,123,123,123	0
5	CA	B	1777	1/1	0.94	0.09	104,104,104,104	0
4	AMP	B	777	23/23	0.95	0.07	63,71,93,99	0
4	AMP	A	777	23/23	0.96	0.07	52,65,78,79	0
3	ZN	A	776	1/1	0.98	0.04	53,53,53,53	0
3	ZN	B	776	1/1	0.98	0.09	56,56,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AMP B 777:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AMP A 777:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.