



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:35 PM UTC

PDB ID : 4V2Q / pdb_00004v2q
Title : Ironing out their differences: Dissecting the structural determinants of a phenylalanine aminomutase and ammonia lyase
Authors : Heberling, M.; Masman, M.; Bartsch, S.; Wybenga, G.G.; Dijkstra, B.W.; Marrink, S.; Janssen, D.
Deposited on : 2014-10-14
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	705	
1	B	705	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9878 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLALANINE AMMONIA-LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4770	3031	816	901	22			
1	B	621	Total	C	N	O	S	0	0	0
			4829	3068	826	913	22			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q68G84
A	-18	GLY	-	expression tag	UNP Q68G84
A	-17	SER	-	expression tag	UNP Q68G84
A	-16	SER	-	expression tag	UNP Q68G84
A	-15	HIS	-	expression tag	UNP Q68G84
A	-14	HIS	-	expression tag	UNP Q68G84
A	-13	HIS	-	expression tag	UNP Q68G84
A	-12	HIS	-	expression tag	UNP Q68G84
A	-11	HIS	-	expression tag	UNP Q68G84
A	-10	HIS	-	expression tag	UNP Q68G84
A	-9	SER	-	expression tag	UNP Q68G84
A	-8	SER	-	expression tag	UNP Q68G84
A	-7	GLY	-	expression tag	UNP Q68G84
A	-6	LEU	-	expression tag	UNP Q68G84
A	-5	VAL	-	expression tag	UNP Q68G84
A	-4	PRO	-	expression tag	UNP Q68G84
A	-3	ARG	-	expression tag	UNP Q68G84
A	-2	GLY	-	expression tag	UNP Q68G84
A	-1	SER	-	expression tag	UNP Q68G84
A	0	HIS	-	expression tag	UNP Q68G84
A	89	THR	CYS	engineered mutation	UNP Q68G84
A	97	GLY	LEU	engineered mutation	UNP Q68G84
A	175	MDO	ALA	chromophore	UNP Q68G84
A	175	MDO	SER	chromophore	UNP Q68G84
A	175	MDO	GLY	chromophore	UNP Q68G84

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	expression tag	UNP Q68G84
B	-18	GLY	-	expression tag	UNP Q68G84
B	-17	SER	-	expression tag	UNP Q68G84
B	-16	SER	-	expression tag	UNP Q68G84
B	-15	HIS	-	expression tag	UNP Q68G84
B	-14	HIS	-	expression tag	UNP Q68G84
B	-13	HIS	-	expression tag	UNP Q68G84
B	-12	HIS	-	expression tag	UNP Q68G84
B	-11	HIS	-	expression tag	UNP Q68G84
B	-10	HIS	-	expression tag	UNP Q68G84
B	-9	SER	-	expression tag	UNP Q68G84
B	-8	SER	-	expression tag	UNP Q68G84
B	-7	GLY	-	expression tag	UNP Q68G84
B	-6	LEU	-	expression tag	UNP Q68G84
B	-5	VAL	-	expression tag	UNP Q68G84
B	-4	PRO	-	expression tag	UNP Q68G84
B	-3	ARG	-	expression tag	UNP Q68G84
B	-2	GLY	-	expression tag	UNP Q68G84
B	-1	SER	-	expression tag	UNP Q68G84
B	0	HIS	-	expression tag	UNP Q68G84
B	89	THR	CYS	engineered mutation	UNP Q68G84
B	97	GLY	LEU	engineered mutation	UNP Q68G84
B	175	MDO	ALA	chromophore	UNP Q68G84
B	175	MDO	SER	chromophore	UNP Q68G84
B	175	MDO	GLY	chromophore	UNP Q68G84


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	154	Total O 154 154	0	0
2	B	125	Total O 125 125	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PHENYLALANINE AMMONIA-LYASE



• Molecule 1: PHENYLALANINE AMMONIA-LYASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.53Å 76.17Å 120.45Å 90.00° 120.42° 90.00°	Depositor
Resolution (Å)	103.87 – 1.95 103.86 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (103.87-1.95) 99.7 (103.86-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.190 , 0.216 0.227 , 0.238	Depositor DCC
R_{free} test set	5159 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.084	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9878	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/4837	0.81	0/6563
1	B	0.65	0/4895	0.81	1/6640 (0.0%)
All	All	0.66	0/9732	0.81	1/13203 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	374	SER	N-CA-C	5.22	116.97	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4770	0	4813	20	0
1	B	4829	0	4885	20	0
2	A	154	0	0	0	0
2	B	125	0	0	0	0
All	All	9878	0	9698	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:ALA:HB3	1:B:279:VAL:HG13	1.66	0.75
1:B:501:GLU:OE1	1:B:619:ARG:HD2	1.91	0.70
1:A:446:PRO:HD3	1:B:446:PRO:HD3	1.76	0.67
1:B:227:LEU:O	1:B:231:ASN:HB2	1.97	0.64
1:A:294:TRP:CH2	1:A:624:LYS:HE2	2.35	0.62
1:A:174:SER:C	1:A:175:MDO:CA1	2.72	0.62
1:A:227:LEU:O	1:A:231:ASN:HB2	2.04	0.58
1:A:437:SER:O	1:A:441:GLN:HG2	2.06	0.55
1:B:55:ALA:O	1:B:59:ARG:HB3	2.11	0.51
1:A:27:ILE:CD1	1:A:43:LEU:HB2	2.40	0.51
1:B:31:GLY:HA3	1:B:61:ARG:HH12	1.78	0.48
1:A:352:ALA:HB3	1:B:279:VAL:CG1	2.39	0.48
1:B:178:ASP:HB3	1:B:181:PRO:HG2	1.96	0.47
1:B:437:SER:O	1:B:441:GLN:HG2	2.14	0.47
1:B:545:CYS:SG	1:B:589:SER:HA	2.56	0.45
1:B:552:LEU:O	1:B:556:LYS:HG3	2.16	0.45
1:A:460:ASP:OD2	1:B:400:GLU:OE2	2.34	0.45
1:B:114:THR:HG22	1:B:122:ASP:HB2	1.98	0.45
1:A:51:VAL:HG21	1:A:159:LEU:HD13	1.98	0.45
1:B:224:LYS:HE2	1:B:356:PRO:HD2	2.01	0.43
1:B:59:ARG:HD2	1:B:63:GLU:CD	2.44	0.43
1:A:131:SER:HB2	1:A:238:ALA:HB1	2.00	0.43
1:A:384:ARG:HG2	1:A:444:ALA:HA	2.00	0.42
1:A:175:MDO:O2	1:A:231:ASN:ND2	2.48	0.42
1:A:111:GLY:HA3	1:A:168:PRO:HG2	2.02	0.42
1:B:209:GLU:O	1:B:213:ARG:HG3	2.19	0.42
1:B:500:GLU:HG3	1:B:537:VAL:HG12	2.01	0.42
1:A:257:LEU:HB3	1:A:490:MET:HE1	2.02	0.42
1:A:330:TRP:O	1:A:333:PRO:HD2	2.20	0.42
1:B:51:VAL:HG21	1:B:159:LEU:HD13	2.01	0.41
1:A:43:LEU:HD22	1:A:134:LEU:HD22	2.03	0.41
1:B:537:VAL:HA	1:B:540:TYR:CE2	2.56	0.41
1:B:174:SER:C	1:B:175:MDO:CA1	2.81	0.41
1:A:59:ARG:HD3	1:A:149:TRP:CG	2.56	0.40
1:A:182:LEU:HD21	1:A:234:SER:HB2	2.03	0.40
1:A:559:CYS:O	1:A:562:THR:HG22	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	596/705 (84%)	585 (98%)	11 (2%)	0	100	100
1	B	604/705 (86%)	592 (98%)	12 (2%)	0	100	100
All	All	1200/1410 (85%)	1177 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	521/591 (88%)	510 (98%)	11 (2%)	47	41
1	B	528/591 (89%)	511 (97%)	17 (3%)	34	25
All	All	1049/1182 (89%)	1021 (97%)	28 (3%)	39	32

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	ARG
1	A	71	ARG
1	A	107	CYS
1	A	231	ASN
1	A	334	LEU
1	A	353	ASN
1	A	455	GLU
1	A	481	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	555	LEU
1	A	602	LEU
1	A	619	ARG
1	B	43	LEU
1	B	59	ARG
1	B	114	THR
1	B	123	GLU
1	B	147	ILE
1	B	231	ASN
1	B	279	VAL
1	B	304	LEU
1	B	334	LEU
1	B	353	ASN
1	B	455	GLU
1	B	481	LEU
1	B	555	LEU
1	B	589	SER
1	B	602	LEU
1	B	640	VAL
1	B	673	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	353	ASN
1	B	231	ASN
1	B	302	GLN
1	B	353	ASN
1	B	413	ASN
1	B	462	ASN
1	B	654	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	MDO	A	175	1	11,13,14	2.89	4 (36%)	15,18,20	1.97	7 (46%)
1	MDO	B	175	1	11,13,14	2.91	4 (36%)	15,18,20	2.05	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MDO	A	175	1	-	3/4/23/24	0/1/1/1
1	MDO	B	175	1	-	1/4/23/24	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	175	MDO	O2-C2	7.24	1.37	1.23
1	A	175	MDO	O2-C2	6.92	1.37	1.23
1	A	175	MDO	O3-C3	4.20	1.43	1.20
1	B	175	MDO	O3-C3	4.02	1.42	1.20
1	A	175	MDO	C2-N3	-3.30	1.32	1.40
1	B	175	MDO	C2-N3	-3.15	1.32	1.40
1	A	175	MDO	CA2-N2	-2.65	1.34	1.39
1	B	175	MDO	CA2-N2	-2.60	1.34	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	MDO	CA2-C2-N3	3.65	106.57	103.50
1	B	175	MDO	CA1-C1-N2	3.28	128.15	124.04
1	B	175	MDO	CB-CA1-C1	3.10	115.79	111.43
1	A	175	MDO	CA1-C1-N2	2.99	127.78	124.04
1	B	175	MDO	CA2-C2-N3	2.93	105.97	103.50
1	B	175	MDO	C2-N3-C1	2.91	109.42	108.07
1	B	175	MDO	N3-C1-N2	-2.87	109.22	111.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	MDO	N3-C1-N2	-2.53	109.49	111.48
1	A	175	MDO	CB-CA1-C1	2.25	114.59	111.43
1	A	175	MDO	CA2-N2-C1	2.24	107.40	105.39
1	B	175	MDO	CA2-N2-C1	2.22	107.38	105.39
1	B	175	MDO	O3-C3-CA3	-2.18	115.66	125.47
1	A	175	MDO	C2-N3-C1	2.09	109.03	108.07
1	A	175	MDO	C2-CA2-N2	-2.03	107.50	108.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	175	MDO	C3-CA3-N3-C2
1	A	175	MDO	C3-CA3-N3-C2
1	A	175	MDO	N2-C1-CA1-CB
1	A	175	MDO	N3-C1-CA1-CB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	175	MDO	2	0
1	B	175	MDO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2
1	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	174:SER	C	175:MDO	N1	1.71
1	B	174:SER	C	175:MDO	N1	1.70
1	B	175:MDO	C3	178:ASP	N	1.65
1	A	175:MDO	C3	178:ASP	N	1.63

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	612/705 (86%)	-0.05	13 (2%) 63 70	17, 26, 43, 57	0
1	B	620/705 (87%)	0.13	22 (3%) 47 53	17, 28, 49, 66	0
All	All	1232/1410 (87%)	0.04	35 (2%) 55 62	17, 27, 46, 66	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	618	VAL	4.2
1	B	203	VAL	4.0
1	B	314	LEU	3.7
1	A	114	THR	3.2
1	B	201	ASP	3.1
1	B	71	ARG	2.9
1	A	573	GLU	2.9
1	A	567	HIS	2.8
1	B	70	GLN	2.7
1	A	75	ASP	2.6
1	B	576	THR	2.6
1	B	55	ALA	2.6
1	A	122	ASP	2.6
1	B	67	SER	2.5
1	B	606	LYS	2.5
1	A	203	VAL	2.5
1	B	587	ARG	2.4
1	A	312	ASP	2.4
1	A	424	TYR	2.3
1	B	424	TYR	2.3
1	A	99	GLU	2.3
1	B	99	GLU	2.2
1	B	573	GLU	2.2
1	A	589	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	313	LYS	2.2
1	A	73	ALA	2.2
1	B	69	VAL	2.2
1	B	566	LEU	2.2
1	B	610	THR	2.2
1	A	586	LYS	2.1
1	B	589	SER	2.1
1	B	222	GLN	2.1
1	B	215	GLY	2.1
1	B	613	ASN	2.0
1	B	519	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MDO	B	175	13/14	0.81	0.12	25,28,29,31	0
1	MDO	A	175	13/14	0.82	0.12	23,25,26,28	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.