



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:44 PM UTC

PDB ID : 2V5P / pdb_00002v5p
Title : COMPLEX STRUCTURE OF HUMAN IGF2R DOMAINS 11-13 BOUND TO IGF-II
Authors : Brown, J.; Delaine, C.; Zaccheo, O.J.; Siebold, C.; Gilbert, R.J.; van Boxel, G.; Denley, A.; Wallace, J.C.; Hassan, A.B.; Forbes, B.E.; Jones, E.Y.
Deposited on : 2007-07-06
Resolution : 4.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1243 (4.40-3.80)
Clashscore	190562	1293 (4.40-3.80)
Ramachandran outliers	187476	1206 (4.40-3.80)
Sidechain outliers	187428	1193 (4.40-3.80)
RSRZ outliers	180081	1240 (4.40-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	492	 2% 77% 17% • 5%
1	B	492	 3% 76% 18% • 5%
2	C	67	 4% 66% 6% 28%
2	D	67	 % 57% 13% • 28%
3	E	3	 67% 33%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	466	Total	C	N	O	S	0	0	0
			3571	2243	604	690	34			
1	B	466	Total	C	N	O	S	0	0	0
			3571	2243	604	690	34			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1703	ALA	GLY	conflict	UNP P11717
B	1703	ALA	GLY	conflict	UNP P11717

- Molecule 2 is a protein called INSULIN-LIKE GROWTH FACTOR II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	48	Total	C	N	O	S	0	0	0
			369	232	58	73	6			
2	D	48	Total	C	N	O	S	0	0	0
			369	232	58	73	6			


- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




• Molecule 1: CATION-INDEPENDENT MANNOSE-6-PHOSPHATE RECEPTOR




• Molecule 2: INSULIN-LIKE GROWTH FACTOR II



● Molecule 2: INSULIN-LIKE GROWTH FACTOR II



● Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	165.98Å 117.31Å 116.67Å 90.00° 123.41° 90.00°	Depositor
Resolution (Å)	47.51 – 4.10 47.51 – 4.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.51-4.10) 99.3 (47.51-4.10)	Depositor EDS
R_{merge}	0.29	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 4.14Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.291 , 0.327 0.289 , 0.318	Depositor DCC
R_{free} test set	741 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	66.9	Xtriage
Anisotropy	1.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 72.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.79	EDS
Total number of atoms	7989	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/3658	0.76	4/4970 (0.1%)
1	B	0.45	0/3658	0.77	3/4970 (0.1%)
2	C	0.37	0/374	0.73	0/503
2	D	0.40	0/374	0.71	0/503
All	All	0.45	0/8064	0.76	7/10946 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1598	CYS	CA-C-N	7.20	128.84	119.84
1	B	1598	CYS	C-N-CA	7.20	128.84	119.84
1	A	1988	PRO	N-CA-C	6.51	118.65	110.70
1	A	1598	CYS	CA-C-N	5.63	126.88	119.84
1	A	1598	CYS	C-N-CA	5.63	126.88	119.84
1	B	1988	PRO	N-CA-C	5.60	117.53	110.70
1	A	1823	SER	N-CA-C	5.11	114.99	108.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3571	0	3411	41	0
1	B	3571	0	3411	45	0
2	C	369	0	339	3	0
2	D	369	0	339	5	0
3	E	39	0	34	1	0
4	A	28	0	26	2	0
4	B	42	0	39	1	0
All	All	7989	0	7599	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1567:PHE:HD2	1:B:1572:ILE:HB	1.40	0.86
1:B:1888:VAL:HG23	1:B:1950:THR:HG22	1.64	0.79
1:B:1770:VAL:HA	1:B:1888:VAL:HG21	1.67	0.74
1:B:1829:ARG:HD3	1:B:1968:VAL:HB	1.69	0.73
1:A:1888:VAL:HG23	1:A:1950:THR:HG22	1.73	0.69
2:C:44:GLU:HA	2:C:48:PHE:CD2	2.29	0.66
1:A:1829:ARG:HD3	1:A:1968:VAL:HB	1.79	0.65
1:A:1641:HIS:CE1	1:A:1700:PRO:HD3	2.31	0.64
1:A:1770:VAL:HA	1:A:1888:VAL:HG21	1.81	0.63
1:B:1629:LEU:HD22	1:B:1636:LEU:HA	1.81	0.63
1:B:1567:PHE:CD2	1:B:1572:ILE:HB	2.29	0.62
1:A:1800:ARG:HH12	1:A:1874:LEU:HD12	1.65	0.61
1:B:1610:ILE:HA	1:B:1638:PHE:HB2	1.84	0.60
1:A:1762:ILE:HG12	1:A:1787:PHE:HB2	1.83	0.60
1:B:1885:LEU:HD23	1:B:1953:ILE:HD12	1.82	0.59
1:A:1567:PHE:HD2	1:A:1572:ILE:HB	1.66	0.59
1:B:1734:ASN:HB3	1:B:1739:GLU:H	1.68	0.57
1:B:1805:THR:HG22	1:B:1816:ASN:HA	1.86	0.56
1:B:1542:TYR:HD2	1:B:1548:VAL:HG23	1.71	0.55
1:A:1664:SER:N	1:A:1665:PRO:HD2	2.21	0.55
1:A:1610:ILE:HA	1:A:1638:PHE:HB2	1.88	0.54
3:E:1:NAG:H61	3:E:2:NAG:HN2	1.73	0.54
1:B:1664:SER:N	1:B:1665:PRO:HD2	2.23	0.54
1:B:1591:VAL:HG22	1:B:1609:VAL:HG12	1.89	0.53
1:A:1781:SER:HB3	1:A:1784:ASP:HB3	1.91	0.53
1:B:1920:GLU:H	1:B:1926:TRP:HE1	1.57	0.53
1:A:1862:LYS:HE3	1:A:1915:GLU:HG3	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:21:CYS:SG	2:D:26:PHE:HB3	2.49	0.53
1:A:1909:PHE:HB3	1:A:1914:TYR:CE1	2.45	0.52
1:A:1987:CYS:O	1:A:1989:PRO:HD3	2.09	0.52
1:B:1888:VAL:HG22	1:B:1889:ASN:H	1.76	0.51
1:A:1549:TYR:HB3	1:A:1558:ASN:HD22	1.75	0.51
1:A:1805:THR:HG22	1:A:1816:ASN:HA	1.92	0.51
1:B:1862:LYS:HE3	1:B:1915:GLU:HG3	1.93	0.51
1:B:1627:ILE:HD11	1:B:1639:SER:HB2	1.93	0.50
1:B:1599:PRO:HB3	2:C:51:CYS:O	2.11	0.50
1:B:1615:ARG:HD2	1:B:1641:HIS:HB3	1.94	0.49
1:B:1651:GLU:HB3	1:B:1730:PRO:HB3	1.94	0.49
1:B:1694:ILE:HG21	1:B:1789:TRP:HZ3	1.77	0.49
1:A:1615:ARG:HD2	1:A:1641:HIS:HB3	1.93	0.49
1:A:1542:TYR:HD2	1:A:1548:VAL:HG23	1.79	0.48
1:B:1794:VAL:HG12	1:B:1794:VAL:O	2.14	0.48
1:A:1738:ASN:CG	4:A:2991:NAG:HN2	2.22	0.48
2:D:8:LEU:HD12	2:D:46:CYS:HB3	1.96	0.47
1:B:1543:SER:HB2	2:C:22:GLY:HA3	1.96	0.47
1:B:1888:VAL:HG22	1:B:1889:ASN:N	2.28	0.47
1:A:1772:MET:HG3	1:A:1948:TYR:CD2	2.49	0.47
1:B:1621:THR:HA	1:B:1919:ILE:HD11	1.96	0.47
1:B:1987:CYS:H	1:B:1988:PRO:CD	2.27	0.47
1:A:1909:PHE:HB3	1:A:1914:TYR:HE1	1.79	0.47
1:B:1560:PRO:HD3	1:B:1571:ARG:HB3	1.98	0.46
1:B:1851:LYS:C	1:B:1853:GLY:H	2.23	0.46
1:B:1839:CYS:SG	1:B:1872:MET:HE1	2.55	0.46
1:B:1749:PRO:HB3	4:B:2992:NAG:H81	1.97	0.46
1:A:1866:PHE:HB3	1:A:1951:SER:HB3	1.98	0.45
1:A:1552:ILE:C	1:A:1554:GLY:H	2.25	0.45
1:A:1843:VAL:CG1	1:A:1847:GLN:HB3	2.47	0.45
1:B:1762:ILE:HG12	1:B:1787:PHE:HB2	1.98	0.45
1:A:1651:GLU:HB3	1:A:1730:PRO:HB3	1.98	0.45
1:A:1908:ILE:HA	1:A:1912:LYS:O	2.16	0.45
1:A:1567:PHE:CD2	1:A:1572:ILE:HB	2.50	0.44
2:D:14:VAL:HA	2:D:17:LEU:HD12	1.99	0.44
1:A:1712:VAL:HB	1:A:1725:GLY:HA3	1.99	0.44
1:A:1540:ALA:HB3	1:A:1548:VAL:HB	2.00	0.44
1:B:1575:GLY:HA3	1:B:1592:TYR:HB3	1.98	0.44
1:A:1738:ASN:HB3	4:A:2991:NAG:HN2	1.83	0.44
1:A:1884:VAL:HG22	1:A:1954:ILE:HG22	1.98	0.44
1:A:1872:MET:HE3	1:A:1885:LEU:HD13	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1663:LEU:HD12	1:B:1731:PRO:HG3	2.00	0.43
1:B:1772:MET:HG3	1:B:1948:TYR:CD2	2.53	0.43
1:B:1953:ILE:HG12	1:B:1979:PHE:HB2	2.01	0.43
1:B:1884:VAL:HG22	1:B:1954:ILE:HG22	2.00	0.43
1:A:1955:PHE:HB3	1:A:1986:VAL:HG21	2.00	0.43
1:A:1575:GLY:HA3	1:A:1592:TYR:HB3	2.00	0.43
2:D:15:ASP:HA	2:D:18:GLN:HE21	1.83	0.42
2:D:43:VAL:O	2:D:47:CYS:HB3	2.18	0.42
1:A:1675:ALA:CB	1:A:1690:PHE:HB2	2.48	0.42
1:A:1577:ALA:HA	1:A:1592:TYR:CE1	2.54	0.42
1:B:1852:ASP:HB3	1:B:1868:ARG:HG2	2.01	0.42
1:A:1851:LYS:C	1:A:1853:GLY:H	2.28	0.42
1:B:1691:TYR:C	1:B:1692:ILE:HG13	2.43	0.42
1:A:1987:CYS:H	1:A:1988:PRO:HD3	1.85	0.42
1:A:1830:ASP:C	1:A:1832:ARG:H	2.28	0.41
1:B:1678:GLU:H	1:B:1678:GLU:CD	2.27	0.41
1:B:1830:ASP:C	1:B:1832:ARG:H	2.28	0.41
1:B:1779:ARG:HB2	1:B:1786:VAL:HB	2.02	0.41
1:B:1909:PHE:HB3	1:B:1914:TYR:CE1	2.54	0.41
1:A:1909:PHE:C	1:A:1911:GLY:H	2.29	0.41
1:B:1556:ASN:HD21	1:B:1558:ASN:HD22	1.70	0.40
1:B:1827:VAL:HG12	1:B:1968:VAL:HG23	2.03	0.40
1:A:1897:THR:HA	1:A:1941:PHE:O	2.21	0.40
1:B:1675:ALA:CB	1:B:1690:PHE:HB2	2.52	0.40
1:A:1624:PRO:HD3	1:A:1640:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	460/492 (94%)	406 (88%)	44 (10%)	10 (2%)	5 31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	460/492 (94%)	394 (86%)	55 (12%)	11 (2%)	4	30
2	C	44/67 (66%)	42 (96%)	2 (4%)	0	100	100
2	D	44/67 (66%)	42 (96%)	2 (4%)	0	100	100
All	All	1008/1118 (90%)	884 (88%)	103 (10%)	21 (2%)	5	32

All (21) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1600	SER
1	B	1599	PRO
1	B	1600	SER
1	B	1738	ASN
1	A	1568	GLY
1	B	1568	GLY
1	B	1736	ILE
1	B	1803	GLY
1	A	1580	ARG
1	A	1599	PRO
1	A	1736	ILE
1	A	1987	CYS
1	B	1695	CYS
1	B	1987	CYS
1	A	1695	CYS
1	B	1745	GLU
1	B	1988	PRO
1	A	1803	GLY
1	A	1988	PRO
1	B	1986	VAL
1	A	1794	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/432 (94%)	393 (97%)	11 (3%)	39	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	404/432 (94%)	393 (97%)	11 (3%)	39	60
2	C	41/57 (72%)	41 (100%)	0	100	100
2	D	41/57 (72%)	40 (98%)	1 (2%)	43	63
All	All	890/978 (91%)	867 (97%)	23 (3%)	40	61

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1609	VAL
1	A	1629	LEU
1	A	1652	CYS
1	A	1695	CYS
1	A	1778	LEU
1	A	1802	ASP
1	A	1809	GLU
1	A	1811	LEU
1	A	1885	LEU
1	A	1919	ILE
1	A	1983	THR
1	B	1629	LEU
1	B	1650	THR
1	B	1652	CYS
1	B	1695	CYS
1	B	1761	LEU
1	B	1802	ASP
1	B	1811	LEU
1	B	1901	VAL
1	B	1919	ILE
1	B	1983	THR
1	B	1985	VAL
2	D	8	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1558	ASN
1	A	1693	ASN
1	A	1699	ASN
1	A	1738	ASN
1	A	1765	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1879	GLN
1	A	1910	ASN
1	A	1937	HIS
1	B	1556	ASN
1	B	1558	ASN
1	B	1641	HIS
1	B	1702	HIS
1	B	1738	ASN
1	B	1765	HIS
1	B	1879	GLN
1	B	1910	ASN
2	D	18	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	E	1	3,1	14,14,15	0.53	0	17,19,21	1.17	1 (5%)
3	NAG	E	2	3	14,14,15	0.44	0	17,19,21	0.94	0
3	BMA	E	3	3	11,11,12	0.61	0	15,15,17	0.85	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	6/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	1	NAG	C1-O5-C5	3.18	116.44	112.19
3	E	3	BMA	C1-C2-C3	2.08	112.68	109.64

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
3	E	2	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
3	E	2	NAG	C1-C2-N2-C7
3	E	2	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1	NAG	1	0
3	E	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	2992	1	14,14,15	0.50	0	17,19,21	1.01	1 (5%)
4	NAG	B	2991	1	14,14,15	0.50	0	17,19,21	0.95	1 (5%)
4	NAG	B	2993	1	14,14,15	0.51	0	17,19,21	1.02	1 (5%)
4	NAG	B	2992	1	14,14,15	0.50	0	17,19,21	0.98	1 (5%)
4	NAG	A	2991	1	14,14,15	0.47	0	17,19,21	1.04	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	2992	1	-	2/6/23/26	0/1/1/1
4	NAG	B	2991	1	-	4/6/23/26	0/1/1/1
4	NAG	B	2993	1	-	4/6/23/26	0/1/1/1
4	NAG	B	2992	1	-	3/6/23/26	0/1/1/1
4	NAG	A	2991	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2991	NAG	C1-O5-C5	3.75	117.21	112.19
4	B	2993	NAG	C1-O5-C5	3.21	116.49	112.19
4	B	2991	NAG	C1-O5-C5	3.19	116.46	112.19
4	B	2992	NAG	C1-O5-C5	3.13	116.38	112.19
4	A	2992	NAG	C1-O5-C5	2.76	115.89	112.19

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2991	NAG	C8-C7-N2-C2
4	A	2991	NAG	O7-C7-N2-C2
4	A	2992	NAG	C8-C7-N2-C2
4	A	2992	NAG	O7-C7-N2-C2
4	B	2991	NAG	C8-C7-N2-C2
4	B	2991	NAG	O7-C7-N2-C2
4	B	2992	NAG	C8-C7-N2-C2
4	B	2992	NAG	O7-C7-N2-C2
4	B	2993	NAG	C8-C7-N2-C2
4	B	2993	NAG	O7-C7-N2-C2
4	B	2993	NAG	O5-C5-C6-O6
4	B	2991	NAG	C4-C5-C6-O6
4	A	2991	NAG	O5-C5-C6-O6
4	B	2991	NAG	O5-C5-C6-O6
4	B	2993	NAG	C4-C5-C6-O6
4	B	2992	NAG	O5-C5-C6-O6
4	A	2991	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2992	NAG	1	0
4	A	2991	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/492 (94%)	0.50	11 (2%) 59 45	46, 93, 135, 189	0
1	B	466/492 (94%)	0.55	14 (3%) 52 39	51, 87, 124, 179	0
2	C	48/67 (71%)	0.75	3 (6%) 26 24	74, 84, 94, 130	0
2	D	48/67 (71%)	0.57	1 (2%) 63 48	51, 67, 90, 165	0
All	All	1028/1118 (91%)	0.54	29 (2%) 55 41	46, 89, 129, 189	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1753	ASP	4.8
1	A	1753	ASP	4.6
1	B	1677	ASP	3.1
1	B	1860	GLY	3.0
1	A	1738	ASN	2.9
1	A	1861	THR	2.8
1	A	1924	LYS	2.7
2	C	8	LEU	2.6
1	B	1623	ARG	2.5
1	B	1923	ALA	2.5
1	B	1586	GLN	2.5
1	B	1846	GLU	2.4
1	B	1568	GLY	2.4
1	A	1847	GLN	2.4
1	B	1837	GLY	2.3
2	C	63	PRO	2.3
2	C	16	THR	2.2
1	B	1847	GLN	2.2
1	A	1757	ASN	2.2
1	B	1569	GLN	2.2
1	A	1687	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1849	GLY	2.1
1	B	1703	ALA	2.1
1	B	1693	ASN	2.1
1	A	1812	LEU	2.1
1	A	1573	SER	2.1
1	B	1640	TRP	2.1
1	A	1606	TYR	2.0
2	D	9	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BMA	E	3	11/12	0.78	0.11	59,59,59,59	0
3	NAG	E	1	14/15	0.81	0.13	60,60,61,61	0
3	NAG	E	2	14/15	0.86	0.09	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	2991	14/15	0.73	0.23	59,59,59,59	0
4	NAG	B	2991	14/15	0.73	0.18	59,59,59,60	0
4	NAG	B	2993	14/15	0.73	0.12	63,63,63,63	0
4	NAG	B	2992	14/15	0.76	0.13	61,61,62,62	0
4	NAG	A	2992	14/15	0.79	0.12	60,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.