



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:35 PM UTC

PDB ID : 7V50 / pdb_00007v50
Title : Structure of cyclohexanone monooxygenase mutant from *Acinetobacter calcoaceticus*
Authors : Wu, Y.; Yu, H.
Deposited on : 2021-08-16
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4388 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative flavin-binding monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	0	0
			4074	2597	680	779	18			

There are 35 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A0A8XFY0
A	-18	GLY	-	expression tag	UNP A0A0A8XFY0
A	-17	SER	-	expression tag	UNP A0A0A8XFY0
A	-16	SER	-	expression tag	UNP A0A0A8XFY0
A	-15	HIS	-	expression tag	UNP A0A0A8XFY0
A	-14	HIS	-	expression tag	UNP A0A0A8XFY0
A	-13	HIS	-	expression tag	UNP A0A0A8XFY0
A	-12	HIS	-	expression tag	UNP A0A0A8XFY0
A	-11	HIS	-	expression tag	UNP A0A0A8XFY0
A	-10	HIS	-	expression tag	UNP A0A0A8XFY0
A	-9	SER	-	expression tag	UNP A0A0A8XFY0
A	-8	SER	-	expression tag	UNP A0A0A8XFY0
A	-7	GLY	-	expression tag	UNP A0A0A8XFY0
A	-6	LEU	-	expression tag	UNP A0A0A8XFY0
A	-5	VAL	-	expression tag	UNP A0A0A8XFY0
A	-4	PRO	-	expression tag	UNP A0A0A8XFY0
A	-3	ARG	-	expression tag	UNP A0A0A8XFY0
A	-2	GLY	-	expression tag	UNP A0A0A8XFY0
A	-1	SER	-	expression tag	UNP A0A0A8XFY0
A	0	HIS	-	expression tag	UNP A0A0A8XFY0
A	246	TYR	PHE	engineered mutation	UNP A0A0A8XFY0
A	277	LEU	PHE	engineered mutation	UNP A0A0A8XFY0
A	326	CYS	LYS	engineered mutation	UNP A0A0A8XFY0
A	386	SER	ASN	engineered mutation	UNP A0A0A8XFY0
A	388	LYS	ILE	engineered mutation	UNP A0A0A8XFY0
A	390	ILE	MET	engineered mutation	UNP A0A0A8XFY0
A	426	PHE	LEU	engineered mutation	UNP A0A0A8XFY0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	432	LEU	PHE	engineered mutation	UNP A0A0A8XFY0
A	433	ALA	THR	engineered mutation	UNP A0A0A8XFY0
A	435	SER	LEU	engineered mutation	UNP A0A0A8XFY0
A	438	ILE	SER	engineered mutation	UNP A0A0A8XFY0
A	488	LYS	GLU	engineered mutation	UNP A0A0A8XFY0
A	489	CYS	SER	engineered mutation	UNP A0A0A8XFY0
A	490	ARG	TRP	engineered mutation	UNP A0A0A8XFY0
A	505	LEU	PHE	engineered mutation	UNP A0A0A8XFY0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	213	Total O 213 213	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative flavin-binding monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.30Å 52.90Å 102.99Å 90.00° 95.82° 90.00°	Depositor
Resolution (Å)	38.87 – 2.30 38.87 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.87-2.30) 99.7 (38.87-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 2.08Å)	Xtriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.211 , 0.252 0.211 , 0.251	Depositor DCC
R_{free} test set	1616 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4388	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/4164	0.31	0/5635

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	147	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4074	0	3988	62	0
2	A	53	0	31	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	48	0	25	2	0
4	A	213	0	0	13	0
All	All	4388	0	4044	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:ASP:OD2	4:A:701:HOH:O	1.83	0.96
1:A:448:ASP:OD2	4:A:702:HOH:O	1.85	0.93
1:A:444:ARG:O	4:A:703:HOH:O	2.00	0.79
1:A:516:GLN:NE2	4:A:709:HOH:O	2.19	0.76
1:A:458:ILE:HD11	1:A:461:ILE:HG13	1.69	0.73
1:A:455:ASP:N	4:A:706:HOH:O	2.13	0.71
1:A:198:GLN:OE1	4:A:705:HOH:O	2.11	0.68
1:A:57:ASP:OD2	1:A:327:ARG:NH2	2.27	0.68
1:A:294:ASN:OD1	1:A:298:GLN:NE2	2.27	0.65
1:A:419:PHE:O	1:A:422:MET:N	2.30	0.65
1:A:50:GLN:NE2	4:A:717:HOH:O	2.36	0.59
1:A:458:ILE:HD12	1:A:460:GLN:H	1.68	0.58
1:A:395:LYS:HD3	1:A:462:GLU:OE2	2.02	0.58
1:A:416:VAL:O	1:A:419:PHE:HD1	1.88	0.57
1:A:280:MET:HE2	1:A:327:ARG:HA	1.88	0.56
1:A:166:SER:HB3	1:A:380:PHE:HE2	1.71	0.55
1:A:388:LYS:HG2	1:A:401:LYS:HA	1.90	0.54
1:A:416:VAL:H	1:A:419:PHE:HB3	1.71	0.54
1:A:356:ILE:HG12	1:A:361:VAL:HG22	1.89	0.54
1:A:61:HIS:O	1:A:232:TYR:OH	2.20	0.53
1:A:102:LYS:NZ	1:A:103:ASP:OD2	2.41	0.53
1:A:419:PHE:O	1:A:420:PRO:C	2.52	0.52
1:A:178:ARG:NH1	4:A:713:HOH:O	2.28	0.52
1:A:174:PHE:HB3	1:A:199:VAL:HG12	1.92	0.51
1:A:291:GLU:HA	1:A:483:LEU:HD11	1.92	0.51
1:A:472:THR:HG23	1:A:503:VAL:HG21	1.92	0.51
1:A:57:ASP:N	2:A:601:FAD:O4	2.44	0.51
1:A:88:LEU:HG	1:A:92:LYS:HE3	1.92	0.51
1:A:41:GLU:HG2	1:A:92:LYS:HE2	1.93	0.51
1:A:411:TYR:HB3	1:A:415:MET:HG3	1.92	0.50
1:A:114:TYR:CE1	1:A:393:ARG:HG3	2.47	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:GLU:HG3	1:A:483:LEU:HD21	1.95	0.49
1:A:49:ASN:O	1:A:84:GLN:HG3	2.12	0.48
1:A:143:LEU:HD21	1:A:383:VAL:HG13	1.97	0.47
1:A:82:ILE:HG12	1:A:87:VAL:HG23	1.95	0.47
1:A:119:ASN:HB2	1:A:121:PHE:CD1	2.49	0.47
1:A:119:ASN:HB2	1:A:121:PHE:CE1	2.49	0.47
1:A:184:THR:HG21	1:A:206:GLN:HB2	1.96	0.47
1:A:298:GLN:HE22	1:A:324:TYR:HA	1.80	0.47
1:A:96:ASP:OD1	1:A:101:ARG:NH1	2.38	0.46
1:A:161:GLU:HB2	1:A:373:MET:HG3	1.98	0.46
1:A:512:GLU:O	1:A:516:GLN:HG2	2.16	0.46
1:A:71:LEU:HD21	1:A:94:VAL:HG22	1.99	0.45
1:A:416:VAL:C	1:A:418:ASN:H	2.25	0.45
1:A:327:ARG:HD3	3:A:602:NAP:H72N	1.81	0.44
1:A:51:TYR:CZ	1:A:187:THR:HG23	2.53	0.44
1:A:486:LYS:HD2	1:A:486:LYS:HA	1.68	0.44
1:A:69:LYS:NZ	4:A:704:HOH:O	2.07	0.44
1:A:146:ALA:HB3	1:A:147:PRO:HD3	1.99	0.44
1:A:337:ILE:HD13	4:A:898:HOH:O	2.19	0.43
1:A:416:VAL:C	1:A:418:ASN:N	2.76	0.43
1:A:464:THR:HG22	1:A:529:LEU:HD21	2.00	0.43
1:A:452:TYR:C	4:A:706:HOH:O	2.63	0.41
1:A:333:GLY:O	1:A:337:ILE:HG12	2.20	0.41
1:A:79:ARG:NH2	4:A:707:HOH:O	2.14	0.41
1:A:288:ALA:N	4:A:711:HOH:O	2.27	0.41
1:A:462:GLU:O	1:A:528:CYS:HA	2.20	0.41
1:A:44:GLY:HA2	2:A:601:FAD:O3B	2.21	0.41
1:A:421:ASN:OD1	1:A:460:GLN:HA	2.21	0.41
1:A:435:SER:HB2	1:A:436:PRO:HD3	2.02	0.41
3:A:602:NAP:H2D	3:A:602:NAP:H2N	1.51	0.41
1:A:72:LEU:HB3	1:A:232:TYR:CD2	2.56	0.41
1:A:81:TYR:CE1	1:A:215:ILE:HG12	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/562 (90%)	490 (96%)	16 (3%)	2 (0%)	30	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	485	ALA
1	A	420	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/483 (91%)	429 (98%)	11 (2%)	42	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	SER
1	A	125	THR
1	A	144	LEU
1	A	222	GLU
1	A	223	THR
1	A	259	SER
1	A	326	CYS
1	A	362	VAL
1	A	378	THR
1	A	458	ILE
1	A	464	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	A	198	GLN
1	A	206	GLN
1	A	522	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	601	-	58,58,58	4.19	21 (36%)	85,89,89	3.25	17 (20%)
3	NAP	A	602	-	50,52,52	2.12	14 (28%)	71,80,80	1.88	12 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	8/34/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	602	-	-	4/35/67/67	0/5/5/5

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C5A-C4A	14.23	1.64	1.39
2	A	601	FAD	C8A-N7A	-12.39	1.07	1.31
2	A	601	FAD	C5A-N7A	-11.73	1.17	1.39
2	A	601	FAD	C2A-N3A	-9.70	1.16	1.33
2	A	601	FAD	C4A-N9A	8.78	1.55	1.37
2	A	601	FAD	C8A-N9A	-8.52	1.23	1.37
3	A	602	NAP	O4D-C1D	8.49	1.52	1.40
2	A	601	FAD	C4A-N3A	-7.52	1.20	1.34
2	A	601	FAD	C5A-C6A	6.26	1.58	1.41
3	A	602	NAP	C7N-N7N	5.67	1.43	1.33
2	A	601	FAD	C2-N1	5.20	1.48	1.36
2	A	601	FAD	C2A-N1A	-4.64	1.25	1.33
2	A	601	FAD	C6A-N6A	4.45	1.45	1.34
2	A	601	FAD	C8M-C8	4.07	1.58	1.51
3	A	602	NAP	C3N-C7N	3.87	1.56	1.50
2	A	601	FAD	C5X-N5	3.57	1.46	1.39
3	A	602	NAP	O4B-C1B	3.57	1.50	1.42
2	A	601	FAD	O4B-C1B	3.51	1.50	1.42
3	A	602	NAP	C2N-N1N	3.25	1.38	1.35
3	A	602	NAP	C2D-C3D	-3.06	1.45	1.53
3	A	602	NAP	C3B-C2B	-3.04	1.46	1.53
2	A	601	FAD	C2-N3	2.76	1.45	1.39
2	A	601	FAD	C6A-N1A	2.62	1.43	1.35
3	A	602	NAP	C5A-N7A	-2.56	1.34	1.39
3	A	602	NAP	C2B-C1B	-2.55	1.46	1.53
2	A	601	FAD	C2B-C1B	-2.54	1.45	1.53
3	A	602	NAP	C1B-N9A	-2.39	1.39	1.46
2	A	601	FAD	C3B-C4B	-2.31	1.47	1.53
3	A	602	NAP	C6A-N6A	2.25	1.40	1.34
2	A	601	FAD	O4B-C4B	2.21	1.49	1.45
2	A	601	FAD	C9-C9A	2.19	1.43	1.39
2	A	601	FAD	C2B-C3B	-2.17	1.47	1.53
3	A	602	NAP	C3D-C4D	-2.08	1.47	1.53
3	A	602	NAP	O4D-C4D	2.07	1.49	1.45
3	A	602	NAP	C3B-C4B	-2.01	1.47	1.53

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	N3A-C4A-N9A	13.51	150.15	127.17
2	A	601	FAD	C5A-N7A-C8A	11.91	122.16	103.45
2	A	601	FAD	C5A-C4A-N9A	-11.58	93.19	105.81
2	A	601	FAD	C6A-C5A-N7A	9.60	150.60	132.09
2	A	601	FAD	C4A-C5A-N7A	-8.06	101.37	110.58
2	A	601	FAD	C2A-N3A-C4A	7.47	130.07	111.83
3	A	602	NAP	C4D-O4D-C1D	-7.42	103.13	109.92
2	A	601	FAD	C5A-C4A-N3A	-7.31	116.65	126.72
2	A	601	FAD	C6A-C5A-C4A	-6.69	108.04	117.18
3	A	602	NAP	N3A-C2A-N1A	-5.65	120.03	128.58
3	A	602	NAP	C5A-C4A-N3A	-5.61	119.00	126.72
2	A	601	FAD	N9A-C8A-N7A	4.64	120.52	113.94
3	A	602	NAP	N3A-C4A-N9A	4.61	135.01	127.17
3	A	602	NAP	C5A-N7A-C8A	3.85	109.49	103.45
3	A	602	NAP	C2A-N3A-C4A	3.78	121.06	111.83
2	A	601	FAD	C4-N3-C2	-3.34	119.71	125.64
2	A	601	FAD	C4B-O4B-C1B	-3.25	102.29	109.47
3	A	602	NAP	C4A-C5A-N7A	-2.99	107.17	110.58
3	A	602	NAP	N9A-C8A-N7A	-2.87	109.86	113.94
2	A	601	FAD	C4A-N9A-C8A	-2.84	102.75	105.74
2	A	601	FAD	C4X-C4-N3	2.76	120.28	113.25
3	A	602	NAP	C6N-N1N-C2N	-2.61	119.66	121.88
2	A	601	FAD	O4-C4-C4X	-2.48	119.98	126.53
3	A	602	NAP	C2D-C3D-C4D	2.25	106.95	102.61
2	A	601	FAD	C4X-C10-N10	2.19	119.61	116.48
3	A	602	NAP	P2B-O2B-C2B	-2.08	117.88	123.43
3	A	602	NAP	C4B-O4B-C1B	-2.08	104.88	109.47
2	A	601	FAD	C2B-C3B-C4B	2.01	106.49	102.61
2	A	601	FAD	C10-C4X-N5	-2.01	120.71	124.81

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	FAD	N10-C1'-C2'-O2'
2	A	601	FAD	N10-C1'-C2'-C3'
3	A	602	NAP	C2D-C1D-N1N-C2N
3	A	602	NAP	C2D-C1D-N1N-C6N
2	A	601	FAD	C2'-C3'-C4'-O4'
2	A	601	FAD	O3'-C3'-C4'-C5'
2	A	601	FAD	C2'-C3'-C4'-C5'
2	A	601	FAD	O3'-C3'-C4'-O4'
2	A	601	FAD	O4B-C4B-C5B-O5B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	601	FAD	C3B-C4B-C5B-O5B
3	A	602	NAP	PN-O3-PA-O2A
3	A	602	NAP	C2B-O2B-P2B-O2X


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	2	0
3	A	602	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	512/562 (91%)	0.46	40 (7%) 19 21	6, 30, 63, 83	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	PHE	7.3
1	A	144	LEU	5.9
1	A	147	PRO	4.6
1	A	503	VAL	4.3
1	A	504	TYR	4.0
1	A	383	VAL	3.6
1	A	484	PHE	3.6
1	A	232	TYR	3.6
1	A	417	SER	3.6
1	A	145	ALA	3.5
1	A	416	VAL	3.3
1	A	325	ALA	3.2
1	A	380	PHE	3.1
1	A	397	GLY	3.1
1	A	146	ALA	3.1
1	A	324	TYR	2.9
1	A	418	ASN	2.8
1	A	466	ASP	2.8
1	A	148	ASN	2.8
1	A	462	GLU	2.8
1	A	143	LEU	2.7
1	A	398	ILE	2.6
1	A	420	PRO	2.6
1	A	326	CYS	2.5
1	A	458	ILE	2.4
1	A	384	ASP	2.4
1	A	452	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	463	ALA	2.3
1	A	150	PRO	2.3
1	A	486	LYS	2.3
1	A	57	ASP	2.3
1	A	121	PHE	2.3
1	A	532	GLN	2.3
1	A	530	LEU	2.2
1	A	396	ASP	2.2
1	A	382	ALA	2.2
1	A	485	ALA	2.1
1	A	129	ASP	2.0
1	A	323	LEU	2.0
1	A	528	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FAD	A	601	53/53	0.90	0.13	16,41,63,64	0
3	NAP	A	602	48/48	0.95	0.09	9,25,63,69	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.