



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:35 PM UTC

PDB ID : 1V7V / pdb_00001v7v
Title : Crystal structure of Vibrio proteolyticus chitobiose phosphorylase
Authors : Hidaka, M.; Honda, Y.; Nirasawa, S.; Kitaoka, M.; Hayashi, K.; Wakagi, T.;
Shoun, H.; Fushinobu, S.
Deposited on : 2003-12-24
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	807	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called chitobiose phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	779	6222	3939	1070	1188	25	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	802	HIS	-	expression tag	UNP Q76IQ9
A	803	HIS	-	expression tag	UNP Q76IQ9
A	804	HIS	-	expression tag	UNP Q76IQ9
A	805	HIS	-	expression tag	UNP Q76IQ9
A	806	HIS	-	expression tag	UNP Q76IQ9
A	807	HIS	-	expression tag	UNP Q76IQ9

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	676	Total	O	0	0
			676	676		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: chitinobiose phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.95Å 70.58Å 80.05Å 90.00° 98.40° 90.00°	Depositor
Resolution (Å)	47.78 – 1.80 47.78 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.2 (47.78-1.80) 98.3 (47.78-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.73 (at 1.81Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.162 , 0.186 0.163 , 0.164	Depositor DCC
R_{free} test set	3578 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6901	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/6400	0.88	23/8698 (0.3%)

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	357	SER	N-CA-C	9.76	123.70	111.69
1	A	320	TRP	N-CA-C	8.09	121.12	111.33
1	A	643	SER	N-CA-C	6.87	118.77	111.28
1	A	334	PHE	N-CA-C	6.77	120.84	112.59
1	A	207	ASP	N-CA-C	-6.72	105.21	113.41
1	A	85	SER	N-CA-C	-6.17	100.52	110.32
1	A	25	THR	N-CA-C	5.93	119.24	110.46
1	A	545	HIS	N-CA-C	5.84	120.69	113.50
1	A	187	LEU	N-CA-C	-5.79	97.79	107.99
1	A	281	GLN	N-CA-C	-5.75	105.53	112.54
1	A	264	TYR	N-CA-C	-5.62	101.01	109.95
1	A	87	GLN	N-CA-C	-5.54	102.09	110.39
1	A	532	THR	N-CA-C	-5.45	105.42	111.36
1	A	425	SER	N-CA-C	5.44	122.39	110.80
1	A	636	LYS	CB-CA-C	-5.35	110.39	116.54
1	A	70	TYR	N-CA-C	5.31	118.39	109.95
1	A	57	LYS	N-CA-C	5.26	117.32	110.53
1	A	60	PRO	N-CA-C	5.26	118.96	111.13
1	A	736	ASP	CA-C-N	5.11	125.09	120.03
1	A	736	ASP	C-N-CA	5.11	125.09	120.03
1	A	188	TYR	N-CA-C	5.09	118.26	111.75
1	A	347	GLY	N-CA-C	-5.07	101.42	111.31
1	A	170	SER	N-CA-C	5.03	117.49	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6222	0	5818	68	0
2	A	3	0	0	0	0
3	A	676	0	0	3	0
All	All	6901	0	5818	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:487:ARG:HD2	1:A:492:ASP:HA	1.56	0.85
1:A:644:HIS:HB3	1:A:645:PRO:HD3	1.65	0.78
1:A:1:MET:CE	1:A:2:LYS:H	1.99	0.74
1:A:385:HIS:H	1:A:418:HIS:HE1	1.34	0.74
1:A:295:ALA:O	1:A:299:GLU:HG3	1.89	0.73
1:A:432:ILE:HB	1:A:433:PRO:HD3	1.75	0.68
1:A:227:LEU:O	1:A:231:GLN:HG2	1.94	0.67
1:A:478:GLY:HA3	1:A:500:GLU:OE2	1.95	0.66
1:A:1:MET:HE2	1:A:2:LYS:H	1.60	0.66
1:A:219:LEU:HG	1:A:220:TYR:CD2	2.31	0.65
1:A:277:ARG:O	1:A:281:GLN:HG3	1.98	0.64
1:A:182:LEU:HD22	1:A:274:GLU:HG3	1.81	0.62
1:A:425:SER:HB3	1:A:496:LEU:HD21	1.81	0.62
1:A:1:MET:HE3	1:A:1:MET:HA	1.81	0.62
1:A:1:MET:HE2	1:A:2:LYS:O	2.04	0.58
1:A:437:LYS:HE3	1:A:437:LYS:HA	1.87	0.57
1:A:384:LEU:HD11	1:A:417:ILE:HD13	1.87	0.56
1:A:689:VAL:HG21	1:A:703:ALA:HB1	1.88	0.56
1:A:644:HIS:O	1:A:647:PRO:HD2	2.06	0.55
1:A:1:MET:HE3	1:A:2:LYS:H	1.70	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ASN:O	1:A:12:GLU:HG3	2.06	0.54
1:A:47:TYR:HB2	1:A:268:ILE:CD1	2.38	0.54
1:A:674:GLN:HG2	1:A:677:ILE:HD11	1.90	0.53
1:A:567:THR:OG1	1:A:570:GLN:HG3	2.11	0.51
1:A:91:LYS:HE3	1:A:113:TYR:OH	2.11	0.49
1:A:421:LYS:HG3	1:A:422:ASP:N	2.27	0.49
1:A:24:TRP:CG	1:A:684:GLU:HG3	2.48	0.49
1:A:420:ILE:HD12	1:A:421:LYS:N	2.28	0.49
1:A:635:VAL:HG12	1:A:636:LYS:HG2	1.93	0.49
1:A:684:GLU:HG2	1:A:686:TYR:CZ	2.48	0.48
1:A:374:LEU:HD13	1:A:452:ILE:HD12	1.95	0.48
1:A:585:LEU:C	1:A:585:LEU:HD23	2.39	0.48
1:A:487:ARG:HB3	3:A:1582:HOH:O	2.14	0.47
1:A:20:VAL:HG11	1:A:24:TRP:CE2	2.50	0.47
1:A:270:LYS:HB3	1:A:270:LYS:NZ	2.29	0.47
1:A:622:ASN:OD1	1:A:624:ASP:HB2	2.16	0.46
1:A:240:THR:O	1:A:241:CYS:HB2	2.16	0.46
1:A:570:GLN:O	1:A:574:ARG:HD3	2.16	0.46
1:A:421:LYS:HE2	1:A:422:ASP:OD1	2.16	0.45
1:A:209:TYR:O	1:A:236:ASN:HA	2.16	0.45
1:A:642:PHE:O	1:A:645:PRO:HD2	2.16	0.45
1:A:417:ILE:HD11	1:A:455:ALA:HB1	1.98	0.45
1:A:588:LEU:C	1:A:588:LEU:HD23	2.41	0.45
1:A:126:LYS:NZ	1:A:296:HIS:HD2	2.14	0.44
1:A:179:ARG:HH22	1:A:184:GLU:CD	2.25	0.44
1:A:204:PHE:C	1:A:204:PHE:CD1	2.96	0.44
1:A:161:GLN:HG2	3:A:1086:HOH:O	2.18	0.43
1:A:512:LEU:O	1:A:516:ILE:HG13	2.18	0.43
1:A:349:ARG:O	1:A:353:GLN:HG3	2.19	0.43
1:A:487:ARG:HH22	1:A:627:PHE:HD2	1.65	0.43
1:A:606:PHE:HA	1:A:611:LEU:HD23	1.99	0.42
1:A:425:SER:HB3	1:A:496:LEU:CD2	2.47	0.42
1:A:151:SER:HB3	1:A:264:TYR:CZ	2.55	0.42
1:A:559:THR:HG22	1:A:565:ILE:HD11	2.01	0.42
1:A:491:ASN:O	1:A:493:CYS:N	2.53	0.41
1:A:66:ARG:HB3	1:A:67:PRO:HA	2.01	0.41
1:A:487:ARG:HH12	1:A:490:TRP:HA	1.84	0.41
1:A:132:ILE:HD11	1:A:280:TYR:CZ	2.56	0.41
1:A:204:PHE:CE1	1:A:250:LYS:HB2	2.56	0.41
1:A:485:GLY:HA2	1:A:495:ASN:O	2.21	0.41
1:A:197:PHE:HB3	3:A:1546:HOH:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:SER:HB3	1:A:53:ASN:HD22	1.86	0.40
1:A:647:PRO:HA	1:A:650:TRP:CD2	2.56	0.40
1:A:100:ILE:HD12	1:A:100:ILE:N	2.36	0.40
1:A:699:ASP:O	1:A:700:HIS:C	2.64	0.40
1:A:133:TRP:HB2	1:A:264:TYR:HB2	2.03	0.40
1:A:163:ASP:O	1:A:167:HIS:HE1	2.05	0.40
1:A:393:ASP:HA	1:A:394:PRO:HD2	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	775/807 (96%)	740 (96%)	34 (4%)	1 (0%)	48 34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	707	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	653/680 (96%)	650 (100%)	3 (0%)	81 80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	PRO
1	A	420	ILE
1	A	437	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	161	GLN
1	A	164	ASN
1	A	167	HIS
1	A	255	GLN
1	A	296	HIS
1	A	305	GLN
1	A	311	GLN
1	A	418	HIS
1	A	428	HIS
1	A	571	GLN
1	A	704	ASN
1	A	721	ASN
1	A	751	GLN
1	A	782	GLN
1	A	795	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	779/807 (96%)	-0.20	21 (2%) 56 56	9, 15, 31, 59	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	490	TRP	5.8
1	A	417	ILE	4.5
1	A	420	ILE	3.8
1	A	197	PHE	3.8
1	A	492	ASP	3.6
1	A	496	LEU	3.6
1	A	497	GLY	3.5
1	A	486	LEU	3.2
1	A	488	ALA	3.1
1	A	418	HIS	2.9
1	A	394	PRO	2.8
1	A	493	CYS	2.8
1	A	498	GLY	2.6
1	A	422	ASP	2.6
1	A	391	TRP	2.5
1	A	489	ASP	2.5
1	A	494	LEU	2.3
1	A	421	LYS	2.3
1	A	480	THR	2.2
1	A	491	ASN	2.1
1	A	782	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	A	1003	1/1	0.97	0.14	40,40,40,40	0
2	CA	A	1002	1/1	0.98	0.14	25,25,25,25	0
2	CA	A	1001	1/1	0.99	0.11	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.