



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:22 AM UTC

PDB ID : 3V8O / pdb_00003v8o
Title : Human Filamin C Ig - like Domains 4 and 5
Authors : Sethi, R.; Ylanne, J.
Deposited on : 2011-12-23
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	7% 72% 20% 8% .
1	B	194	6% 64% 23% 9% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Filamin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	0	0	0
			1444	916	241	280	7			
1	B	188	Total	C	N	O	S	0	0	0
			1410	896	234	273	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	568	GLN	-	expression tag	UNP Q14315
B	568	GLN	-	expression tag	UNP Q14315

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	K	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Filamin-C



• Molecule 1: Filamin-C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.62Å 91.35Å 103.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.63 – 2.80 46.63 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.63-2.80) 97.8 (46.63-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.200 , 0.255 0.209 , 0.253	Depositor DCC
R_{free} test set	1512 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2902	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	4/1482 (0.3%)	1.18	6/2019 (0.3%)
1	B	1.20	1/1448 (0.1%)	1.22	8/1973 (0.4%)
All	All	1.20	5/2930 (0.2%)	1.20	14/3992 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	692	ILE	CA-CB	-5.66	1.47	1.54
1	A	757	ASN	CA-C	-5.52	1.45	1.52
1	A	744	TRP	C-O	-5.25	1.18	1.23
1	A	590	ALA	CA-CB	-5.09	1.46	1.53
1	B	631	TRP	C-O	-5.04	1.18	1.24

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	759	GLY	N-CA-C	-7.07	101.80	112.41
1	A	759	GLY	N-CA-C	-7.05	101.84	112.41
1	A	654	PHE	O-C-N	-6.14	116.22	123.22
1	A	617	GLU	N-CA-C	5.88	118.03	108.26
1	B	758	VAL	CA-C-N	-5.81	116.52	122.27
1	B	758	VAL	C-N-CA	-5.81	116.52	122.27
1	B	746	GLY	N-CA-C	-5.73	106.07	114.61
1	B	749	VAL	N-CA-CB	-5.68	103.25	111.21
1	A	719	ILE	N-CA-C	5.43	113.92	107.73
1	B	719	ILE	N-CA-C	5.34	113.32	107.60
1	A	758	VAL	CA-C-N	-5.33	117.00	122.27
1	A	758	VAL	C-N-CA	-5.33	117.00	122.27
1	B	626	CYS	N-CA-C	5.29	117.23	109.24
1	B	662	PRO	O-C-N	5.08	123.54	121.15

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1444	0	1404	47	0
1	B	1410	0	1371	75	0
2	A	1	0	0	0	0
3	A	21	0	0	1	0
3	B	26	0	0	8	0
All	All	2902	0	2775	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:721:ASN:ND2	1:A:725:THR:HG22	1.63	1.11
1:A:721:ASN:HD22	1:A:725:THR:CG2	1.68	1.06
1:B:721:ASN:ND2	1:B:725:THR:HG22	1.70	1.04
1:B:615:LYS:HB2	3:B:823:HOH:O	1.59	1.00
1:B:721:ASN:HD22	1:B:725:THR:HG22	0.87	1.00
1:B:721:ASN:HD22	1:B:725:THR:CG2	1.76	0.98
1:A:721:ASN:HD22	1:A:725:THR:HG22	0.79	0.95
1:B:723:ASP:OD1	1:B:725:THR:HB	1.65	0.95
1:B:650:ARG:O	1:B:651:ASP:HB2	1.69	0.92
1:B:750:PRO:O	1:B:751:LYS:HB2	1.70	0.92
1:A:723:ASP:OD1	1:A:725:THR:HB	1.71	0.90
1:A:573:LYS:HG2	1:A:598:GLY:HA3	1.56	0.87
1:A:750:PRO:O	1:A:751:LYS:HB2	1.74	0.86
1:B:641:HIS:NE2	1:B:653:PRO:HG3	1.92	0.83
1:B:683:ILE:HG21	1:B:686:LYS:HG3	1.62	0.82
1:A:684:VAL:HG23	1:A:760:GLU:HB2	1.62	0.80
1:B:759:GLY:O	1:B:760:GLU:HB2	1.79	0.80
1:A:759:GLY:O	1:A:760:GLU:HB2	1.81	0.79

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:650:ARG:HH11	1:B:650:ARG:HB2	1.48	0.79
1:A:749:VAL:HG13	1:A:750:PRO:HD2	1.64	0.77
1:B:575:ARG:NH2	1:B:597:ILE:HD11	1.99	0.77
1:A:749:VAL:CG1	1:A:750:PRO:HD2	2.15	0.76
1:B:742:ILE:O	1:B:749:VAL:HB	1.86	0.75
1:B:650:ARG:HH11	1:B:650:ARG:CG	2.02	0.73
1:A:684:VAL:HG23	1:A:760:GLU:CB	2.18	0.73
1:B:683:ILE:CG2	1:B:686:LYS:HG3	2.18	0.73
1:B:684:VAL:O	1:B:685:ASP:HB2	1.87	0.72
1:B:650:ARG:HH11	1:B:650:ARG:CB	2.04	0.69
1:B:650:ARG:O	1:B:651:ASP:CB	2.42	0.68
1:A:760:GLU:CD	1:A:760:GLU:O	2.38	0.67
1:B:651:ASP:O	3:B:821:HOH:O	2.11	0.67
1:B:750:PRO:O	1:B:751:LYS:CB	2.45	0.64
1:B:760:GLU:CD	1:B:760:GLU:C	2.66	0.64
1:B:683:ILE:HG21	1:B:686:LYS:CG	2.29	0.63
1:A:572:GLN:HB3	1:A:650:ARG:NH2	2.14	0.62
1:B:760:GLU:CD	1:B:760:GLU:O	2.43	0.61
1:B:760:GLU:O	1:B:760:GLU:OE2	2.18	0.61
1:B:641:HIS:CD2	1:B:653:PRO:HB3	2.36	0.61
1:B:641:HIS:CD2	1:B:653:PRO:HG3	2.37	0.60
1:B:650:ARG:NH1	1:B:650:ARG:HG3	2.16	0.59
1:A:760:GLU:CD	1:A:760:GLU:C	2.71	0.59
1:B:683:ILE:HG22	1:B:686:LYS:HB2	1.84	0.58
1:B:599:THR:HG21	1:B:603:THR:HG21	1.85	0.58
1:A:568:GLN:HE21	1:A:573:LYS:HE2	1.68	0.58
1:A:712:CYS:SG	1:B:712:CYS:SG	3.02	0.58
1:B:650:ARG:CG	1:B:650:ARG:NH1	2.64	0.58
1:B:743:SER:OG	3:B:820:HOH:O	2.17	0.57
1:B:733:THR:HG22	1:B:734:LYS:HG3	1.87	0.57
1:A:712:CYS:HB3	1:B:712:CYS:SG	2.45	0.57
1:B:668:ASP:HA	1:B:750:PRO:CG	2.35	0.57
1:A:601:VAL:HG21	3:A:920:HOH:O	2.05	0.56
1:A:749:VAL:CG1	1:A:750:PRO:CD	2.84	0.56
1:B:603:THR:CG2	1:B:603:THR:O	2.53	0.55
1:A:634:GLU:HG3	1:A:635:PRO:HD2	1.86	0.55
1:A:657:HIS:HE1	3:B:826:HOH:O	1.89	0.55
1:B:684:VAL:HB	1:B:760:GLU:HG3	1.88	0.54
1:B:575:ARG:HH22	1:B:597:ILE:HD11	1.73	0.53
1:A:571:VAL:HG23	1:B:715:ASP:OD1	2.08	0.53
1:B:652:SER:OG	1:B:653:PRO:HA	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:683:ILE:HG23	1:B:684:VAL:N	2.24	0.53
1:B:745:GLY:O	1:B:747:VAL:HG23	2.09	0.53
1:B:717:LYS:O	1:B:728:CYS:HA	2.09	0.52
1:A:650:ARG:NH1	1:B:705:TYR:CD1	2.78	0.52
1:B:683:ILE:CG2	1:B:684:VAL:N	2.70	0.52
1:B:650:ARG:HH11	1:B:650:ARG:HG3	1.72	0.51
1:B:751:LYS:O	1:B:754:PHE:CZ	2.64	0.50
1:B:759:GLY:O	1:B:760:GLU:CB	2.49	0.50
1:A:749:VAL:HG12	1:A:750:PRO:HD2	1.94	0.50
1:A:712:CYS:SG	1:B:712:CYS:HB3	2.51	0.50
1:A:575:ARG:NH2	1:A:597:ILE:HD11	2.27	0.50
1:B:751:LYS:O	1:B:754:PHE:CE1	2.65	0.49
1:A:572:GLN:HB3	1:A:650:ARG:HH21	1.78	0.49
1:B:684:VAL:HG23	1:B:760:GLU:HB2	1.96	0.48
1:A:721:ASN:ND2	1:A:725:THR:CG2	2.48	0.48
1:B:575:ARG:HH22	1:B:597:ILE:CD1	2.27	0.47
1:B:731:VAL:HG23	1:B:731:VAL:O	2.14	0.47
1:A:648:ASP:HB3	1:A:652:SER:HB3	1.96	0.47
1:A:573:LYS:CG	1:A:598:GLY:HA3	2.36	0.46
1:A:717:LYS:O	1:A:728:CYS:HA	2.16	0.46
1:A:760:GLU:O	1:A:760:GLU:OE1	2.34	0.46
1:B:630:TYR:CD2	1:B:632:PRO:HD3	2.50	0.46
1:B:680:THR:HG22	3:B:809:HOH:O	2.14	0.46
1:A:581:LEU:HD12	1:A:581:LEU:HA	1.72	0.46
1:B:603:THR:O	1:B:603:THR:HG23	2.15	0.45
1:B:683:ILE:N	3:B:817:HOH:O	2.46	0.45
1:B:641:HIS:CD2	1:B:653:PRO:CG	3.00	0.45
1:A:649:ILE:O	1:A:650:ARG:C	2.60	0.44
1:A:655:ILE:N	1:A:655:ILE:HD13	2.33	0.44
1:B:683:ILE:CG2	1:B:686:LYS:CG	2.89	0.44
1:B:721:ASN:ND2	1:B:725:THR:CG2	2.55	0.44
1:B:577:TRP:O	1:B:592:PHE:HB2	2.17	0.44
1:A:668:ASP:OD1	1:A:668:ASP:N	2.47	0.44
1:B:731:VAL:O	1:B:731:VAL:CG2	2.66	0.43
1:B:645:ASP:HA	3:B:824:HOH:O	2.18	0.43
1:B:615:LYS:CB	3:B:823:HOH:O	2.41	0.43
1:B:608:ILE:HG21	1:B:614:ALA:HB3	2.00	0.43
1:B:683:ILE:CG2	1:B:686:LYS:HB2	2.47	0.43
1:A:733:THR:HG22	1:A:734:LYS:HG3	2.02	0.42
1:B:652:SER:HG	1:B:653:PRO:HA	1.83	0.42
1:A:715:ASP:OD1	1:B:573:LYS:HE3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:731:VAL:HA	1:A:732:PRO:HD2	1.91	0.42
1:B:581:LEU:HG	1:B:654:PHE:CG	2.55	0.42
1:B:688:ALA:O	1:B:729:SER:HA	2.19	0.42
1:A:615:LYS:H	1:A:615:LYS:HE2	1.85	0.42
1:B:641:HIS:CG	1:B:653:PRO:HB3	2.55	0.42
1:A:612:SER:HB3	1:A:634:GLU:HB2	2.03	0.41
1:A:653:PRO:HD2	1:B:655:ILE:HB	2.03	0.41
1:A:600:GLU:O	1:A:603:THR:HB	2.21	0.41
1:B:599:THR:HG21	1:B:603:THR:CG2	2.49	0.41
1:A:603:THR:O	1:A:603:THR:CG2	2.68	0.41
1:A:608:ILE:HD13	1:A:608:ILE:HA	1.81	0.41
1:B:618:CYS:SG	1:B:626:CYS:SG	3.17	0.40
1:A:712:CYS:CB	1:B:712:CYS:SG	3.09	0.40
1:A:749:VAL:HG12	1:A:750:PRO:CD	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/194 (98%)	184 (96%)	7 (4%)	0	100	100
1	B	186/194 (96%)	178 (96%)	6 (3%)	2 (1%)	11	36
All	All	377/388 (97%)	362 (96%)	13 (3%)	2 (0%)	24	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	651	ASP
1	B	751	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	155/155 (100%)	135 (87%)	20 (13%)	4	14
1	B	152/155 (98%)	129 (85%)	23 (15%)	3	10
All	All	307/310 (99%)	264 (86%)	43 (14%)	3	12

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	573	LYS
1	A	581	LEU
1	A	597	ILE
1	A	601	VAL
1	A	603	THR
1	A	608	ILE
1	A	609	GLU
1	A	615	LYS
1	A	617	GLU
1	A	643	ILE
1	A	650	ARG
1	A	652	SER
1	A	680	THR
1	A	683	ILE
1	A	716	ILE
1	A	717	LYS
1	A	725	THR
1	A	731	VAL
1	A	733	THR
1	A	751	LYS
1	B	581	LEU
1	B	597	ILE
1	B	601	VAL
1	B	603	THR
1	B	608	ILE
1	B	609	GLU
1	B	613	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	616	ILE
1	B	617	GLU
1	B	643	ILE
1	B	650	ARG
1	B	655	ILE
1	B	680	THR
1	B	683	ILE
1	B	686	LYS
1	B	716	ILE
1	B	717	LYS
1	B	725	THR
1	B	731	VAL
1	B	733	THR
1	B	743	SER
1	B	749	VAL
1	B	751	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	568	GLN
1	A	572	GLN
1	A	641	HIS
1	A	657	HIS
1	A	721	ASN
1	B	641	HIS
1	B	657	HIS
1	B	721	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	193/194 (99%)	0.07	13 (6%)	24 17	12, 25, 56, 70	0
1	B	188/194 (96%)	0.08	11 (5%)	28 21	12, 25, 56, 70	0
All	All	381/388 (98%)	0.08	24 (6%)	26 19	12, 25, 56, 70	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	683	ILE	4.7
1	A	568	GLN	4.2
1	B	745	GLY	4.1
1	B	602	GLY	4.0
1	B	597	ILE	3.7
1	A	599	THR	3.7
1	B	617	GLU	3.4
1	B	599	THR	3.3
1	B	653	PRO	3.3
1	B	751	LYS	3.1
1	A	601	VAL	3.0
1	B	601	VAL	3.0
1	B	651	ASP	3.0
1	A	602	GLY	2.4
1	A	616	ILE	2.4
1	A	650	ARG	2.4
1	A	760	GLU	2.4
1	A	647	GLU	2.3
1	A	615	LYS	2.3
1	A	597	ILE	2.2
1	B	760	GLU	2.2
1	A	569	ALA	2.2
1	A	651	ASP	2.1
1	A	603	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	K	A	801	1/1	0.97	0.04	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.