



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 05:35 AM UTC

PDB ID : 3V8W / pdb_00003v8w
Title : Crystal Structure of Interleukin-2 Inducible T-cell Kinase Itk Catalytic Domain with Thienopyrazolylindole Inhibitor 469
Authors : McLean, L.R.; Zhang, Y.
Deposited on : 2011-12-23
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	27% 87% 5% 9%
1	B	266	23% 87% • • 9%

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase ITK/TSK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total 1841	C 1179	N 311	O 336	S 15	0	0	0
1	B	242	Total 1839	C 1181	N 307	O 336	S 15	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	355	GLY	-	expression tag	UNP Q08881
A	356	SER	-	expression tag	UNP Q08881
B	355	GLY	-	expression tag	UNP Q08881
B	356	SER	-	expression tag	UNP Q08881

- Molecule 2 is 3-[2-(5-phenyl-2H-thieno[3,2-c]pyrazol-3-yl)-1H-indol-6-yl]pentan-3-ol (CCD ID: 0G2) (formula: C₂₄H₂₃N₃OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			29	24	3	1	1		
2	B	1	Total	C	N	O	S	0	0
			29	24	3	1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	22	Total	O	0	0
			22	22		
4	B	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase ITK/TSK



- Molecule 1: Tyrosine-protein kinase ITK/TSK



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.08Å 69.29Å 68.58Å 90.00° 108.04° 90.00°	Depositor
Resolution (Å)	38.50 – 2.27 38.50 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.50-2.27) 99.7 (38.50-2.27)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 2.27Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.9.1, BUSTER 2.9.1	Depositor
R, R_{free}	0.251 , 0.268 0.271 , 0.283	Depositor DCC
R_{free} test set	1521 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.297	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3792	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0G2, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1879	1.28	0/2542
1	B	0.71	0/1879	1.27	2/2546 (0.1%)
All	All	0.71	0/3758	1.28	2/5088 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	438	MET	CA-C-N	5.20	127.50	120.38
1	B	438	MET	C-N-CA	5.20	127.50	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1841	0	1739	5	0
1	B	1839	0	1754	5	0
2	A	29	0	23	1	0
2	B	29	0	23	1	0
3	B	5	0	0	0	0
4	A	22	0	0	0	0
4	B	27	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3792	0	3539	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:438:MET:H	2:B:702:OG2:H17	1.43	0.67
1:B:393:ILE:HG12	1:B:431:ILE:HG13	1.78	0.65
1:A:438:MET:H	2:A:701:OG2:H17	1.48	0.59
1:A:468:GLU:HG3	1:A:609:LEU:HD11	1.91	0.53
1:B:468:GLU:HG3	1:B:609:LEU:HD11	1.96	0.47
1:B:419:VAL:HG21	1:B:489:LEU:HD12	1.96	0.46
1:A:528:GLU:HB3	1:A:534:ARG:HB2	2.01	0.43
1:A:421:LEU:HD11	1:A:433:LEU:HB3	2.01	0.42
1:B:608:LEU:HA	1:B:611:GLN:HE21	1.84	0.41
1:A:608:LEU:HA	1:A:611:GLN:HE21	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/266 (88%)	226 (97%)	6 (3%)	1 (0%)	30	36
1	B	234/266 (88%)	228 (97%)	6 (3%)	0	100	100
All	All	467/532 (88%)	454 (97%)	12 (3%)	1 (0%)	43	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	398	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/233 (79%)	183 (99%)	2 (1%)	65	79
1	B	190/233 (82%)	187 (98%)	3 (2%)	55	71
All	All	375/466 (80%)	370 (99%)	5 (1%)	61	76

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	500	ASP
1	A	590	ILE
1	B	393	ILE
1	B	421	LEU
1	B	426	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	420	GLN
1	A	494	GLN
1	A	592	ASN
1	A	611	GLN
1	B	420	GLN
1	B	494	GLN
1	B	592	ASN
1	B	611	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	701	-	4,4,4	0.29	0	6,6,6	0.07	0
2	0G2	B	702	-	31,33,33	2.15	11 (35%)	39,49,49	1.68	6 (15%)
2	0G2	A	701	-	31,33,33	2.15	10 (32%)	39,49,49	1.63	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0G2	B	702	-	-	4/20/20/20	0/5/5/5
2	0G2	A	701	-	-	1/20/20/20	0/5/5/5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	0G2	NAK-NAL	-5.91	1.24	1.36
2	B	702	0G2	NAK-NAL	-5.85	1.24	1.36
2	A	701	0G2	CAJ-CAA	-3.99	1.40	1.45
2	B	702	0G2	CAJ-CAA	-3.93	1.40	1.45
2	B	702	0G2	CAN-CAM	-3.89	1.39	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	0G2	CAN-CAM	-3.77	1.39	1.44
2	A	701	0G2	CAR-CAP	-3.71	1.40	1.47
2	B	702	0G2	CAR-CAP	-3.67	1.40	1.47
2	B	702	0G2	CAE-CAB	-3.48	1.33	1.44
2	A	701	0G2	CAE-CAB	-3.46	1.33	1.44
2	A	701	0G2	CAA-NAC	-3.29	1.33	1.38
2	B	702	0G2	CAA-NAC	-3.19	1.33	1.38
2	A	701	0G2	CAD-NAC	-3.01	1.33	1.38
2	B	702	0G2	CAM-SAQ	-2.92	1.68	1.74
2	B	702	0G2	CAD-NAC	-2.90	1.33	1.38
2	A	701	0G2	CAM-SAQ	-2.86	1.68	1.74
2	B	702	0G2	CAB-CAA	-2.46	1.33	1.37
2	A	701	0G2	CAB-CAA	-2.45	1.34	1.37
2	A	701	0G2	CAO-CAN	-2.26	1.35	1.41
2	B	702	0G2	CAO-CAN	-2.23	1.35	1.41
2	B	702	0G2	CAS-CAH	2.03	1.55	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	702	0G2	CAN-NAL-NAK	5.65	112.27	104.58
2	A	701	0G2	CAN-NAL-NAK	5.65	112.27	104.58
2	B	702	0G2	CAP-SAQ-CAM	5.43	96.09	91.75
2	A	701	0G2	CAP-SAQ-CAM	5.06	95.79	91.75
2	B	702	0G2	CAM-CAN-NAL	-3.55	104.96	111.52
2	A	701	0G2	CAM-CAN-NAL	-3.51	105.03	111.52
2	B	702	0G2	CAN-CAO-CAP	2.97	116.07	108.23
2	A	701	0G2	CAN-CAO-CAP	2.92	115.95	108.23
2	B	702	0G2	CAI-CAD-CAE	-2.47	120.15	122.59
2	A	701	0G2	CAI-CAD-CAE	-2.44	120.18	122.59
2	B	702	0G2	CAO-CAP-SAQ	-2.00	107.40	110.35

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	702	0G2	CAY-CAS-CAZ-CBC
2	B	702	0G2	OBA-CAS-CAZ-CBC
2	B	702	0G2	CAH-CAS-CAZ-CBC
2	B	702	0G2	NAC-CAA-CAJ-CAM
2	A	701	0G2	NAC-CAA-CAJ-CAM


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	702	0G2	1	0
2	A	701	0G2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/266 (91%)	1.68	73 (30%)  	36, 52, 77, 88	0
1	B	242/266 (90%)	1.53	62 (25%)  	35, 51, 74, 92	0
All	All	485/532 (91%)	1.61	135 (27%)  	35, 52, 76, 92	0

All (135) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	619	GLY	7.7
1	A	575	PHE	6.6
1	A	619	GLY	6.3
1	A	560	GLU	6.1
1	B	372	GLY	5.4
1	A	372	GLY	5.2
1	B	429	ALA	5.1
1	A	578	TYR	5.1
1	A	567	VAL	4.9
1	B	394	ARG	4.9
1	A	564	ASN	4.6
1	B	502	GLY	4.5
1	A	576	ARG	4.4
1	A	577	LEU	4.4
1	B	403	PHE	4.2
1	B	564	ASN	4.2
1	A	562	ARG	4.2
1	A	397	ALA	4.1
1	A	393	ILE	4.0
1	B	404	ILE	4.0
1	B	400	GLU	4.0
1	B	575	PHE	4.0
1	A	565	SER	3.9
1	A	426	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	570	ASP	3.9
1	A	521	PRO	3.8
1	A	617	GLU	3.7
1	A	502	GLY	3.7
1	A	573	THR	3.7
1	B	393	ILE	3.6
1	A	574	GLY	3.6
1	A	569	GLU	3.6
1	A	559	TYR	3.5
1	A	532	PHE	3.5
1	B	399	SER	3.5
1	B	402	ASP	3.5
1	B	532	PHE	3.5
1	A	403	PHE	3.4
1	A	571	ILE	3.3
1	B	401	GLU	3.3
1	A	563	SER	3.3
1	B	477	CYS	3.3
1	A	572	SER	3.3
1	A	568	VAL	3.3
1	B	425	CYS	3.2
1	B	568	VAL	3.2
1	A	579	LYS	3.2
1	B	572	SER	3.2
1	A	615	ILE	3.2
1	B	398	MET	3.1
1	A	536	SER	3.1
1	A	371	SER	3.1
1	B	371	SER	3.0
1	B	426	LEU	3.0
1	B	427	GLU	3.0
1	A	616	ALA	2.9
1	A	535	TYR	2.9
1	A	399	SER	2.9
1	A	527	PRO	2.9
1	A	409	VAL	2.9
1	B	616	ALA	2.9
1	A	522	VAL	2.9
1	A	530	PHE	2.8
1	B	523	LYS	2.8
1	B	521	PRO	2.8
1	B	617	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	404	ILE	2.8
1	A	355	GLY	2.8
1	A	421	LEU	2.8
1	B	531	SER	2.7
1	B	406	GLU	2.7
1	B	442	CYS	2.7
1	B	355	GLY	2.6
1	A	428	GLN	2.6
1	B	601	ASP	2.6
1	A	370	GLY	2.6
1	B	499	SER	2.6
1	A	385	LYS	2.6
1	B	501	PHE	2.6
1	A	398	MET	2.5
1	A	524	TRP	2.5
1	A	557	ILE	2.5
1	B	529	VAL	2.5
1	B	574	GLY	2.5
1	B	561	ASN	2.5
1	B	405	GLU	2.4
1	A	586	HIS	2.4
1	A	377	VAL	2.4
1	A	501	PHE	2.4
1	B	567	VAL	2.4
1	B	570	ASP	2.4
1	B	618	SER	2.4
1	B	530	PHE	2.4
1	A	410	MET	2.3
1	A	590	ILE	2.3
1	A	442	CYS	2.3
1	B	433	LEU	2.3
1	B	578	TYR	2.3
1	A	431	ILE	2.3
1	A	525	ALA	2.3
1	B	409	VAL	2.3
1	A	429	ALA	2.3
1	A	391	LYS	2.3
1	B	576	ARG	2.2
1	A	585	THR	2.2
1	B	496	ILE	2.2
1	A	529	VAL	2.2
1	B	562	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	455	ALA	2.2
1	A	456	ALA	2.2
1	B	458	THR	2.2
1	B	615	ILE	2.2
1	B	392	THR	2.2
1	A	551	VAL	2.2
1	B	428	GLN	2.2
1	B	377	VAL	2.1
1	B	522	VAL	2.1
1	B	408	GLU	2.1
1	B	421	LEU	2.1
1	A	394	ARG	2.1
1	A	538	LYS	2.1
1	A	406	GLU	2.1
1	B	528	GLU	2.1
1	A	357	VAL	2.1
1	A	587	VAL	2.1
1	B	370	GLY	2.1
1	A	356	SER	2.1
1	B	407	ALA	2.1
1	A	427	GLU	2.1
1	A	582	LEU	2.1
1	A	365	PHE	2.1
1	B	391	LYS	2.0
1	A	364	THR	2.0
1	B	585	THR	2.0
1	B	431	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	701	5/5	0.93	0.11	52,56,57,57	0
2	OG2	A	701	29/29	0.94	0.12	25,39,75,121	0
2	OG2	B	702	29/29	0.95	0.10	24,41,71,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OG2 A 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.