



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 02:31 PM UTC

PDB ID : 6V8C / pdb_00006v8c
Title : Design, Synthesis, and Mechanism of Fluorine-substituted Cyclohexene Analogues of GAMA-Aminobutyric Acid (GABA) as Selective Ornithine Amino-transferase Inactivators
Authors : Zhu, W.; Doubleday, P.T.; Catlin, D.S.; Weerawarna, P.; Butrin, A.; Shen, S.; Kelleher, N.L.; Liu, D.; Silverman, R.B.
Deposited on : 2019-12-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	20% 91% 9%
1	B	404	6% 90% 10%
1	C	404	20% 90% 10%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 10360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ornithine aminotransferase, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	0	0
			3157	2028	532	585	12			
1	B	404	Total	C	N	O	S	0	0	0
			3161	2030	533	586	12			
1	C	404	Total	C	N	O	S	0	0	0
			3157	2028	532	585	12			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is 3-aminocyclohexa-1,3-diene-1-carboxylic acid (CCD ID: QRM) (formula: $C_7H_9NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	7	1	2		
3	B	1	Total	C	N	O	0	0
			10	7	1	2		
3	C	1	Total	C	N	O	0	0
			10	7	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	269	Total	O	0	0
			269	269		
4	B	286	Total	O	0	0
			286	286		
4	C	255	Total	O	0	0
			255	255		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ornithine aminotransferase, mitochondrial




- Molecule 1: Ornithine aminotransferase, mitochondrial



- Molecule 1: Ornithine aminotransferase, mitochondrial





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	193.78Å 193.78Å 57.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.97 – 1.90 27.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.2 (27.97-1.90) 96.2 (27.97-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.01 (at 1.91Å)	Xtriage
Refinement program	PHENIX 1.17.1	Depositor
R, R_{free}	0.239 , 0.268 0.239 , 0.268	Depositor DCC
R_{free} test set	4746 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.045 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10360	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 81.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6362e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QRM, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3231	0.44	0/4388
1	B	0.24	0/3235	0.44	0/4393
1	C	0.22	0/3231	0.43	0/4388
All	All	0.23	0/9697	0.44	0/13169

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3157	0	3156	30	1
1	B	3161	0	3162	30	1
1	C	3157	0	3156	33	0
2	A	15	0	7	1	0
2	B	15	0	7	1	0
2	C	15	0	7	1	0
3	A	10	0	0	1	0
3	B	10	0	0	1	0
3	C	10	0	0	1	0
4	A	269	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	286	0	0	4	0
4	C	255	0	0	6	0
All	All	10360	0	9495	96	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:LEU:H	1:B:405:LYS:NZ	1.78	0.81
1:C:367:VAL:HG23	1:C:388:TRP:CZ2	2.19	0.78
1:A:58:LEU:H	1:A:405:LYS:HZ1	1.32	0.76
1:A:58:LEU:H	1:A:405:LYS:NZ	1.82	0.76
1:C:314:ILE:HG22	4:C:601:HOH:O	1.85	0.75
1:A:58:LEU:HB2	1:A:405:LYS:HE2	1.66	0.75
1:C:314:ILE:O	4:C:601:HOH:O	2.09	0.70
1:C:274:ARG:NH1	4:C:604:HOH:O	2.25	0.68
1:C:391:TRP:HZ3	1:C:405:LYS:HE3	1.58	0.68
1:B:242:ASP:OD2	4:B:601:HOH:O	2.10	0.67
1:B:254:THR:HG22	4:B:847:HOH:O	1.95	0.67
1:B:367:VAL:HG13	1:B:388:TRP:CH2	2.30	0.66
1:B:58:LEU:H	1:B:405:LYS:HZ1	1.41	0.66
1:B:58:LEU:HB2	1:B:405:LYS:HE2	1.77	0.66
1:A:242:ASP:OD2	4:A:601:HOH:O	2.13	0.66
1:A:367:VAL:HG11	1:A:436:ILE:HG23	1.78	0.65
1:B:367:VAL:HG11	1:B:436:ILE:HG23	1.79	0.64
1:C:275:TRP:HZ2	1:C:341:GLU:HG3	1.63	0.64
1:C:384:GLU:HG3	1:C:389:ASP:HB3	1.81	0.62
1:C:364:PRO:HG2	1:C:367:VAL:CG1	2.30	0.62
1:A:244:GLY:HA2	1:A:247:MET:HE3	1.83	0.60
1:A:167:LYS:HE3	4:A:813:HOH:O	2.01	0.60
1:A:367:VAL:HG13	1:A:388:TRP:CH2	2.37	0.59
1:C:169:LYS:HD2	1:C:203:GLY:HA2	1.85	0.59
1:C:314:ILE:C	4:C:601:HOH:O	2.45	0.59
1:C:340:LEU:HD23	1:C:345:LEU:HD12	1.85	0.58
1:C:132:ASN:HB3	4:C:766:HOH:O	2.03	0.58
1:B:344:ASN:HB3	1:B:347:GLU:OE2	2.05	0.57
1:A:340:LEU:HD23	1:A:345:LEU:HD12	1.88	0.56
1:B:382:ILE:HG23	1:B:388:TRP:CH2	2.40	0.56
1:C:391:TRP:CZ3	1:C:405:LYS:HE3	2.39	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:244:GLY:HA2	1:C:247:MET:HE3	1.88	0.55
1:C:367:VAL:HG23	1:C:388:TRP:HZ2	1.70	0.54
1:C:367:VAL:HG21	1:C:436:ILE:HG23	1.90	0.53
1:B:58:LEU:H	1:B:405:LYS:HZ3	1.54	0.53
1:C:364:PRO:HG2	1:C:367:VAL:HG12	1.91	0.52
1:A:430:GLU:HG2	4:B:777:HOH:O	2.09	0.52
1:A:125:GLU:O	1:A:129:LYS:HG2	2.08	0.52
1:A:178:TRP:CZ2	1:A:184:ALA:HA	2.44	0.52
1:C:367:VAL:HG23	1:C:388:TRP:CH2	2.44	0.52
1:B:351:LYS:HD2	1:B:352:LEU:HD23	1.92	0.51
1:B:244:GLY:HA2	1:B:247:MET:HE3	1.93	0.50
1:A:372:ARG:HD2	4:A:701:HOH:O	2.11	0.50
1:B:367:VAL:HG13	1:B:388:TRP:CZ2	2.47	0.50
1:A:382:ILE:HG23	1:A:388:TRP:CH2	2.46	0.49
1:C:56:HIS:O	1:C:405:LYS:NZ	2.40	0.49
1:A:367:VAL:HG12	1:A:368:VAL:HG23	1.94	0.49
1:A:358:ASN:HA	1:A:361:MET:HE3	1.94	0.49
1:C:178:TRP:CZ2	1:C:184:ALA:HA	2.47	0.49
1:B:58:LEU:N	1:B:405:LYS:HZ3	2.11	0.48
1:B:366:ASP:OD2	1:B:367:VAL:HG23	2.14	0.48
1:A:164:GLN:HB2	1:A:167:LYS:HE2	1.95	0.48
1:C:247:MET:HG3	1:C:281:GLU:HB3	1.95	0.48
1:B:178:TRP:CZ2	1:B:184:ALA:HA	2.50	0.47
1:B:252:LEU:HD23	1:B:255:ARG:HH21	1.80	0.46
1:A:149:ALA:HB2	1:A:289:LEU:HD21	1.98	0.45
1:A:165:LYS:HA	1:A:166:TYR:HA	1.65	0.45
1:A:369:THR:HG21	4:A:769:HOH:O	2.16	0.45
1:B:385:THR:OG1	1:B:388:TRP:N	2.49	0.45
1:C:242:ASP:OD2	4:C:603:HOH:O	2.21	0.45
1:C:364:PRO:HG2	1:C:367:VAL:HG11	1.98	0.45
1:A:247:MET:HG3	1:A:281:GLU:HB3	1.98	0.45
1:B:149:ALA:HB2	1:B:289:LEU:HD21	1.99	0.44
1:B:387:ASP:OD1	1:B:387:ASP:N	2.51	0.44
2:A:501:PLP:O3	3:A:502:QRM:N10	2.48	0.44
1:B:165:LYS:HA	1:B:166:TYR:HA	1.67	0.44
1:B:247:MET:HG3	1:B:281:GLU:HB3	1.99	0.44
1:C:215:LEU:O	1:C:219:LEU:HG	2.17	0.44
1:B:58:LEU:N	1:B:405:LYS:NZ	2.57	0.44
1:C:358:ASN:HA	1:C:361:MET:HE3	2.00	0.44
1:B:119:VAL:HG11	1:B:331:ARG:HG2	2.00	0.44
1:B:62:LEU:HA	1:B:72:ASP:HA	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:501:PLP:O3	3:B:502:QRM:N10	2.50	0.43
1:C:165:LYS:HA	1:C:166:TYR:HA	1.68	0.43
1:C:394:CYS:SG	1:C:406:PRO:HD3	2.58	0.43
1:A:232:ILE:HD11	1:A:374:LYS:HD2	2.01	0.43
1:A:367:VAL:HG13	1:A:388:TRP:CZ2	2.53	0.43
1:C:119:VAL:HG11	1:C:331:ARG:HG2	2.00	0.43
1:A:167:LYS:O	1:A:169:LYS:NZ	2.48	0.42
1:A:341:GLU:HA	1:A:341:GLU:OE2	2.20	0.42
1:C:169:LYS:CD	1:C:203:GLY:HA2	2.50	0.42
1:C:275:TRP:CZ2	1:C:341:GLU:HG3	2.50	0.42
1:C:385:THR:HG1	1:C:388:TRP:H	1.68	0.42
1:A:261:ILE:HG12	1:A:287:ILE:HD12	2.01	0.42
1:A:68:ILE:HA	1:A:417:PRO:HG2	2.02	0.41
1:A:169:LYS:HA	1:A:169:LYS:HD3	1.89	0.41
1:B:128:THR:HB	1:B:133:TYR:O	2.21	0.41
1:B:164:GLN:HB2	1:B:167:LYS:HE2	2.02	0.41
1:B:384:GLU:HG3	1:B:389:ASP:HB3	2.02	0.41
1:B:358:ASN:HA	1:B:361:MET:HE3	2.03	0.41
1:C:149:ALA:HB2	1:C:289:LEU:HD21	2.01	0.41
1:C:423:ASP:OD1	1:C:424:GLU:N	2.54	0.40
2:C:501:PLP:O3	3:C:502:QRM:N10	2.52	0.40
1:A:58:LEU:N	1:A:405:LYS:NZ	2.60	0.40
1:A:367:VAL:HG13	1:A:388:TRP:HH2	1.84	0.40
1:B:64:ARG:NH2	4:B:603:HOH:O	2.30	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ALA:O	1:A:405:LYS:NZ[5_555]	1.85	0.35
1:B:114:ALA:O	1:B:405:LYS:NZ[6_565]	1.92	0.28

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/404 (100%)	383 (95%)	18 (4%)	1 (0%)	43	36
1	B	402/404 (100%)	385 (96%)	16 (4%)	1 (0%)	43	36
1	C	402/404 (100%)	383 (95%)	17 (4%)	2 (0%)	24	16
All	All	1206/1212 (100%)	1151 (95%)	51 (4%)	4 (0%)	36	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	257	GLN
1	A	292	LYS
1	B	292	LYS
1	C	292	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/337 (100%)	336 (100%)	0	100	100
1	B	337/337 (100%)	337 (100%)	0	100	100
1	C	336/337 (100%)	336 (100%)	0	100	100
All	All	1009/1011 (100%)	1009 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	266	GLN
1	B	257	GLN
1	B	266	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	344	ASN
1	C	266	GLN
1	C	400	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	501	3	15,15,16	1.07	2 (13%)	21,22,23	0.94	2 (9%)
3	QRM	A	502	2	9,10,10	3.24	4 (44%)	11,13,13	1.87	4 (36%)
3	QRM	C	502	2	9,10,10	3.24	4 (44%)	11,13,13	1.92	4 (36%)
2	PLP	C	501	3	15,15,16	1.07	2 (13%)	21,22,23	0.98	2 (9%)
3	QRM	B	502	2	9,10,10	3.29	3 (33%)	11,13,13	1.83	4 (36%)
2	PLP	B	501	3	15,15,16	0.97	1 (6%)	21,22,23	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	501	3	-	0/6/6/8	0/1/1/1
3	QRM	A	502	2	-	0/4/14/14	0/1/1/1
3	QRM	C	502	2	-	0/4/14/14	0/1/1/1
2	PLP	C	501	3	-	0/6/6/8	0/1/1/1
3	QRM	B	502	2	-	0/4/14/14	0/1/1/1
2	PLP	B	501	3	-	0/6/6/8	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	QRM	C3-C2	-6.26	1.39	1.50
3	C	502	QRM	C3-C2	-6.11	1.39	1.50
3	A	502	QRM	C3-C2	-6.09	1.39	1.50
3	B	502	QRM	C3-C4	-5.45	1.39	1.52
3	C	502	QRM	C3-C4	-5.39	1.39	1.52
3	A	502	QRM	C3-C4	-5.38	1.39	1.52
3	B	502	QRM	C4-C5	-4.02	1.39	1.50
3	C	502	QRM	C4-C5	-3.92	1.39	1.50
3	A	502	QRM	C4-C5	-3.88	1.40	1.50
2	C	501	PLP	C2-N1	2.49	1.38	1.33
2	A	501	PLP	C2-N1	2.39	1.38	1.33
2	B	501	PLP	C6-N1	2.20	1.38	1.34
2	A	501	PLP	C6-N1	2.12	1.38	1.34
2	C	501	PLP	C6-N1	2.08	1.38	1.34
3	A	502	QRM	C5-C6	2.02	1.40	1.34
3	C	502	QRM	C5-C6	2.02	1.40	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	QRM	C3-C4-C5	3.43	120.51	111.14
3	C	502	QRM	C3-C4-C5	3.38	120.37	111.14
3	A	502	QRM	C3-C4-C5	3.29	120.11	111.14
3	C	502	QRM	C4-C3-C2	2.54	120.79	113.72
3	C	502	QRM	C4-C5-C6	-2.50	118.62	122.75
3	A	502	QRM	C4-C3-C2	2.47	120.58	113.72
3	B	502	QRM	C4-C3-C2	2.40	120.41	113.72
3	C	502	QRM	C5-C6-N10	-2.40	120.01	123.99
3	B	502	QRM	C4-C5-C6	-2.38	118.83	122.75
2	C	501	PLP	C6-C5-C4	2.33	120.01	118.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	PLP	C5-C6-N1	-2.28	120.11	123.83
3	A	502	QRM	C4-C5-C6	-2.28	118.99	122.75
2	A	501	PLP	C6-C5-C4	2.23	119.92	118.10
2	A	501	PLP	C5-C6-N1	-2.22	120.21	123.83
3	A	502	QRM	C5-C6-N10	-2.22	120.31	123.99
3	B	502	QRM	C5-C6-N10	-2.15	120.42	123.99

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

6 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PLP	1	0
3	A	502	QRM	1	0
3	C	502	QRM	1	0
2	C	501	PLP	1	0
3	B	502	QRM	1	0
2	B	501	PLP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand QRM A 502



Ligand QRM C 502





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/404 (100%)	1.19	79 (19%) 3 3	14, 25, 43, 87	0
1	B	404/404 (100%)	-0.12	24 (5%) 28 30	5, 15, 32, 61	0
1	C	404/404 (100%)	1.21	79 (19%) 3 3	13, 25, 45, 86	0
All	All	1212/1212 (100%)	0.76	182 (15%) 5 5	5, 22, 43, 87	0

All (182) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	388	TRP	8.3
1	C	388	TRP	6.7
1	A	41	ASP	6.1
1	A	37	PRO	5.5
1	A	388	TRP	5.3
1	B	257	GLN	5.3
1	C	255	ARG	5.2
1	C	254	THR	4.9
1	C	45	GLU	4.9
1	B	367	VAL	4.9
1	A	275	TRP	4.8
1	B	365	SER	4.7
1	B	347	GLU	4.6
1	A	386	LYS	4.6
1	C	367	VAL	4.5
1	A	45	GLU	4.5
1	B	405	LYS	4.4
1	B	37	PRO	4.4
1	C	386	LYS	4.4
1	C	275	TRP	4.3
1	A	38	PRO	4.3
1	C	385	THR	4.2
1	C	41	ASP	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	347	GLU	4.2
1	B	41	ASP	4.2
1	A	341	GLU	4.1
1	B	254	THR	4.1
1	B	385	THR	4.1
1	A	126	TYR	4.0
1	A	130	LEU	4.0
1	C	48	TYR	3.9
1	C	37	PRO	3.9
1	A	385	THR	3.8
1	C	36	GLY	3.8
1	C	341	GLU	3.8
1	B	387	ASP	3.8
1	B	430	GLU	3.8
1	C	87	ALA	3.7
1	A	334	ILE	3.6
1	A	365	SER	3.6
1	B	255	ARG	3.6
1	B	36	GLY	3.6
1	B	45	GLU	3.6
1	A	347	GLU	3.5
1	A	36	GLY	3.5
1	C	126	TYR	3.5
1	A	86	SER	3.5
1	B	341	GLU	3.5
1	C	351	LYS	3.5
1	B	82	LEU	3.4
1	A	166	TYR	3.4
1	B	351	LYS	3.3
1	C	86	SER	3.3
1	A	43	ILE	3.3
1	A	85	TYR	3.3
1	C	334	ILE	3.3
1	A	405	LYS	3.3
1	C	133	TYR	3.2
1	C	436	ILE	3.2
1	A	48	TYR	3.2
1	A	254	THR	3.1
1	C	130	LEU	3.1
1	C	39	THR	3.1
1	A	258	VAL	3.1
1	C	38	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	85	TYR	3.1
1	A	133	TYR	3.0
1	A	351	LYS	3.0
1	C	320	GLY	3.0
1	C	166	TYR	3.0
1	A	87	ALA	3.0
1	A	156	TRP	3.0
1	C	405	LYS	3.0
1	C	131	PHE	2.9
1	A	387	ASP	2.9
1	A	255	ARG	2.8
1	A	312	LEU	2.8
1	A	345	LEU	2.8
1	C	43	ILE	2.8
1	C	314	ILE	2.8
1	A	128	THR	2.8
1	C	60	VAL	2.8
1	C	55	TYR	2.8
1	A	60	VAL	2.8
1	A	367	VAL	2.7
1	C	304	VAL	2.7
1	A	129	LYS	2.7
1	A	58	LEU	2.7
1	A	257	GLN	2.7
1	C	430	GLU	2.7
1	C	346	ALA	2.7
1	A	352	LEU	2.7
1	C	345	LEU	2.7
1	B	383	LYS	2.7
1	A	134	HIS	2.7
1	A	366	ASP	2.6
1	B	366	ASP	2.6
1	A	131	PHE	2.6
1	C	74	GLU	2.6
1	C	305	LEU	2.6
1	C	312	LEU	2.6
1	A	409	GLY	2.6
1	C	258	VAL	2.6
1	A	305	LEU	2.6
1	C	51	GLY	2.6
1	A	61	ALA	2.6
1	C	44	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	49	LYS	2.5
1	A	382	ILE	2.5
1	C	61	ALA	2.5
1	C	319	HIS	2.5
1	B	386	LYS	2.5
1	A	82	LEU	2.5
1	B	382	ILE	2.5
1	C	274	ARG	2.5
1	C	168	ALA	2.5
1	A	384	GLU	2.5
1	C	387	ASP	2.5
1	A	383	LYS	2.5
1	A	168	ALA	2.5
1	A	340	LEU	2.4
1	C	134	HIS	2.4
1	A	212	LEU	2.4
1	A	251	GLU	2.4
1	A	418	LEU	2.4
1	C	49	LYS	2.4
1	C	259	LEU	2.4
1	A	304	VAL	2.4
1	C	129	LYS	2.4
1	A	62	LEU	2.3
1	C	62	LEU	2.3
1	A	50	TYR	2.3
1	C	192	THR	2.3
1	A	333	ALA	2.3
1	A	358	ASN	2.3
1	A	320	GLY	2.3
1	C	156	TRP	2.3
1	A	430	GLU	2.3
1	A	55	TYR	2.3
1	C	132	ASN	2.3
1	C	365	SER	2.3
1	A	119	VAL	2.3
1	C	246	LEU	2.3
1	C	340	LEU	2.3
1	A	59	PRO	2.2
1	C	333	ALA	2.2
1	C	409	GLY	2.2
1	C	40	SER	2.2
1	A	314	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	42	ASP	2.2
1	A	339	VAL	2.2
1	C	160	VAL	2.2
1	C	344	ASN	2.2
1	C	50	TYR	2.2
1	C	382	ILE	2.2
1	A	313	THR	2.2
1	A	259	LEU	2.2
1	A	39	THR	2.2
1	B	409	GLY	2.2
1	A	274	ARG	2.2
1	A	420	ILE	2.1
1	A	407	THR	2.1
1	A	419	VAL	2.1
1	C	136	VAL	2.1
1	C	59	PRO	2.1
1	A	256	HIS	2.1
1	C	419	VAL	2.1
1	C	165	LYS	2.1
1	A	149	ALA	2.1
1	A	164	GLN	2.1
1	A	132	ASN	2.1
1	C	352	LEU	2.1
1	C	225	ALA	2.1
1	C	366	ASP	2.1
1	B	392	LYS	2.0
1	A	157	GLY	2.0
1	C	283	VAL	2.0
1	C	212	LEU	2.0
1	C	256	HIS	2.0
1	C	260	PHE	2.0
1	A	51	GLY	2.0
1	C	88	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	QRM	C	502	10/10	0.81	0.13	18,24,26,27	0
3	QRM	A	502	10/10	0.84	0.11	19,23,25,27	0
2	PLP	A	501	15/16	0.94	0.09	15,17,21,21	0
2	PLP	C	501	15/16	0.94	0.09	12,16,19,20	0
3	QRM	B	502	10/10	0.95	0.06	15,18,19,21	0
2	PLP	B	501	15/16	0.98	0.06	6,9,12,13	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around QRM A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.