



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:10 AM UTC

PDB ID : 2V9K / pdb_00002v9k
Title : Crystal structure of human PUS10, a novel pseudouridine synthase.
Authors : McCleverty, C.J.; Hornsby, M.; Spraggon, G.; Kreusch, A.
Deposited on : 2007-08-23
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	530	5% 71% 14% • 12%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4093 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UNCHARACTERIZED PROTEIN FLJ32312.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	467	Total	C	N	O	S	5	0	0
			3702	2368	633	681	20			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q3MIT2


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cl	0	0
			4	4		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	350	Total 350	O 350	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UNCHARACTERIZED PROTEIN FLJ32312



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.61Å 80.61Å 153.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	69.84 – 2.00 69.81 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (69.84-2.00) 95.6 (69.81-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.230 0.185 , 0.227	Depositor DCC
R_{free} test set	1909 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4093	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, ZN, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/3784	0.89	14/5124 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	523	TRP	CA-C-N	10.37	131.06	120.38
1	A	523	TRP	C-N-CA	10.37	131.06	120.38
1	A	2	PHE	CA-C-N	8.08	127.80	119.56
1	A	2	PHE	C-N-CA	8.08	127.80	119.56
1	A	524	PRO	N-CA-C	-8.05	100.87	110.70
1	A	267	CYS	N-CA-C	7.26	121.10	108.75
1	A	498	LYS	CA-C-N	7.20	128.83	119.84
1	A	498	LYS	C-N-CA	7.20	128.83	119.84
1	A	268	PRO	N-CA-C	-5.88	103.53	110.70
1	A	310	GLY	N-CA-C	-5.84	107.62	115.21
1	A	523	TRP	N-CA-C	5.67	117.66	109.04
1	A	143	VAL	CB-CA-C	-5.47	102.49	110.81
1	A	230	CYS	CA-C-N	5.15	124.81	119.56
1	A	230	CYS	C-N-CA	5.15	124.81	119.56

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	0	GLY	Peptide
1	A	244	PHE	Peptide
1	A	267	CYS	Peptide
1	A	498	LYS	Peptide
1	A	523	TRP	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3702	0	3639	51	0
2	A	4	0	0	0	0
3	A	1	0	0	0	0
4	A	30	0	34	2	0
5	A	6	0	8	2	0
6	A	350	0	0	6	0
All	All	4093	0	3681	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:ASN:OD1	5:A:1536:GOL:H2	1.82	0.78
1:A:413:THR:HG22	1:A:517:GLU:HB2	1.67	0.77
1:A:113:LEU:HD22	1:A:272:PRO:HG3	1.65	0.77
1:A:442:GLN:HE22	1:A:485:TYR:H	1.34	0.75
1:A:184:GLU:OE1	6:A:2082:HOH:O	2.07	0.72
1:A:473:HIS:NE2	4:A:1535:EPE:O1S	2.22	0.71
1:A:290:VAL:CG1	1:A:392:LEU:HD22	2.23	0.68
1:A:348:ARG:HD3	1:A:515:ASP:OD1	1.93	0.68
1:A:290:VAL:HG12	1:A:392:LEU:HD22	1.76	0.67
1:A:413:THR:HG21	6:A:2314:HOH:O	1.95	0.66
1:A:246:ARG:HG2	1:A:246:ARG:HH21	1.60	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:HIS:HD2	1:A:286:GLY:O	1.82	0.63
1:A:491:HIS:HD2	1:A:493:ASP:H	1.45	0.63
1:A:293:ARG:NH2	1:A:524:PRO:HD2	2.15	0.62
1:A:343:GLU:OE2	1:A:349:THR:HG21	1.99	0.61
1:A:268:PRO:HG2	5:A:1536:GOL:H11	1.82	0.60
1:A:491:HIS:CD2	1:A:493:ASP:H	2.19	0.59
1:A:245:THR:HG23	1:A:248:ALA:H	1.69	0.58
1:A:413:THR:CG2	1:A:517:GLU:HB2	2.32	0.57
1:A:290:VAL:HG12	1:A:392:LEU:CD2	2.36	0.56
1:A:246:ARG:HH21	1:A:246:ARG:CG	2.19	0.55
1:A:143:VAL:HG22	1:A:206:SER:HB3	1.87	0.55
1:A:442:GLN:NE2	1:A:485:TYR:H	2.06	0.52
1:A:3:PRO:HG2	1:A:36:PRO:HB3	1.92	0.52
1:A:413:THR:HB	1:A:518:SER:OG	2.10	0.51
1:A:299:ARG:CZ	1:A:348:ARG:HD2	2.41	0.51
1:A:130:GLN:OE1	6:A:2039:HOH:O	2.19	0.50
1:A:167:MET:HE2	1:A:174:LEU:HG	1.92	0.50
1:A:296:LYS:HB3	1:A:349:THR:HG23	1.93	0.50
1:A:153:SER:HB3	1:A:250:MET:HE3	1.95	0.49
1:A:419:TRP:CD2	4:A:1535:EPE:H61	2.48	0.48
1:A:45:LEU:O	1:A:49:GLN:HG2	2.13	0.48
1:A:6:GLU:HG2	1:A:9:LYS:HE2	1.96	0.48
1:A:2:PHE:HA	1:A:3:PRO:HD3	1.75	0.47
1:A:145:SER:O	1:A:211:SER:HA	2.16	0.46
1:A:290:VAL:CG1	1:A:392:LEU:CD2	2.91	0.45
1:A:285:HIS:HE1	6:A:2149:HOH:O	1.99	0.45
1:A:430:GLU:HG3	6:A:2276:HOH:O	2.16	0.45
1:A:335:PHE:HA	1:A:359:GLU:O	2.17	0.45
1:A:457:ARG:NH2	1:A:483:GLY:O	2.49	0.44
1:A:406:GLY:HA2	1:A:410:LYS:HG2	1.99	0.44
1:A:489:PHE:O	1:A:500:ASN:HB2	2.18	0.44
1:A:429:ILE:HD12	1:A:466:THR:HB	2.01	0.43
1:A:148:PHE:HB3	1:A:152:LEU:HD12	2.00	0.43
1:A:293:ARG:CZ	1:A:524:PRO:HD2	2.48	0.42
1:A:149:PRO:HA	1:A:150:PRO:HD3	1.96	0.42
1:A:464:MET:HB3	1:A:476:LEU:HD11	2.01	0.42
1:A:265:PHE:HA	1:A:266:PRO:HD3	1.93	0.42
1:A:459:ARG:HG2	1:A:484:THR:OG1	2.20	0.42
1:A:324:ASP:C	6:A:2179:HOH:O	2.62	0.41
1:A:246:ARG:CG	1:A:246:ARG:NH2	2.80	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/530 (87%)	449 (97%)	10 (2%)	2 (0%)	30	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	499	PRO
1	A	524	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/477 (85%)	379 (94%)	25 (6%)	16	13

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LEU
1	A	48	LEU
1	A	107	ASN
1	A	131	LYS
1	A	142	LEU
1	A	143	VAL
1	A	162	LEU
1	A	174	LEU
1	A	196	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	220	VAL
1	A	226	LEU
1	A	245	THR
1	A	246	ARG
1	A	250	MET
1	A	271	SER
1	A	279	LEU
1	A	345	VAL
1	A	349	THR
1	A	376	LEU
1	A	392	LEU
1	A	413	THR
1	A	451	ARG
1	A	452	ARG
1	A	459	ARG
1	A	508	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	170	GLN
1	A	285	HIS
1	A	300	ASN
1	A	382	ASN
1	A	393	GLN
1	A	442	GLN
1	A	462	HIS
1	A	472	HIS
1	A	491	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	A	1535	-	15,15,15	0.65	0	19,20,20	1.81	5 (26%)
4	EPE	A	1534	-	15,15,15	0.96	1 (6%)	19,20,20	1.80	5 (26%)
5	GOL	A	1536	-	5,5,5	0.44	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	A	1535	-	-	5/9/19/19	0/1/1/1
4	EPE	A	1534	-	-	5/9/19/19	0/1/1/1
5	GOL	A	1536	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1534	EPE	C10-S	3.28	1.82	1.77

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1535	EPE	C5-N4-C3	4.26	118.02	108.84
4	A	1534	EPE	C5-N4-C3	3.77	116.95	108.84
4	A	1534	EPE	O2S-S-C10	3.59	112.15	106.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1535	EPE	C7-N4-C3	3.02	119.29	111.24
4	A	1534	EPE	C7-N4-C5	2.98	119.19	111.24
4	A	1535	EPE	C6-N1-C2	2.82	114.92	108.84
4	A	1534	EPE	C6-N1-C2	2.79	114.84	108.84
4	A	1535	EPE	O1S-S-C10	2.57	110.61	106.73
4	A	1534	EPE	C7-N4-C3	2.24	117.22	111.24
4	A	1535	EPE	O2S-S-C10	2.06	109.84	106.73

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1534	EPE	C9-C10-S-O1S
4	A	1534	EPE	C9-C10-S-O3S
4	A	1535	EPE	N4-C7-C8-O8
4	A	1534	EPE	C10-C9-N1-C6
4	A	1535	EPE	C10-C9-N1-C6
4	A	1534	EPE	C9-C10-S-O2S
4	A	1535	EPE	C8-C7-N4-C5
4	A	1535	EPE	S-C10-C9-N1
4	A	1534	EPE	C10-C9-N1-C2
4	A	1535	EPE	C8-C7-N4-C3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1535	EPE	2	0
5	A	1536	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	467/530 (88%)	0.46	27 (5%)	29 27	21, 36, 62, 93	11 (2%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	ALA	4.5
1	A	310	GLY	4.3
1	A	244	PHE	3.3
1	A	168	GLY	3.3
1	A	229	ILE	3.1
1	A	1	MET	3.1
1	A	528	ASP	3.1
1	A	308	ILE	3.0
1	A	2	PHE	2.9
1	A	455	ALA	2.7
1	A	172	LEU	2.7
1	A	309	ASP	2.7
1	A	449	LEU	2.6
1	A	0	GLY	2.6
1	A	224	HIS	2.6
1	A	174	LEU	2.6
1	A	230	CYS	2.5
1	A	225	PHE	2.5
1	A	453	PRO	2.5
1	A	223	CYS	2.4
1	A	245	THR	2.3
1	A	6	GLU	2.2
1	A	246	ARG	2.2
1	A	227	ALA	2.2
1	A	41	TYR	2.1
1	A	45	LEU	2.1
1	A	454	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EPE	A	1535	15/15	0.67	0.36	57,59,67,68	15
5	GOL	A	1536	6/6	0.80	0.13	47,52,53,54	6
4	EPE	A	1534	15/15	0.89	0.12	28,47,61,61	0
2	CL	A	1531	1/1	0.96	0.13	43,43,43,43	0
2	CL	A	1530	1/1	0.97	0.12	40,40,40,40	0
2	CL	A	1532	1/1	0.98	0.14	39,39,39,39	0
2	CL	A	1529	1/1	0.98	0.13	36,36,36,36	0
3	ZN	A	1533	1/1	0.99	0.10	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.