



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 10:52 AM EDT

PDB ID : 2V9Y / pdb_00002v9y
Title : Human aminoimidazole ribonucleotide synthetase
Authors : Welin, M.; Lehtio, L.; Arrowsmith, C.H.; Berglund, H.; Busam, R.; Collins, R.; Dahlgren, L.G.; Herman, M.D.; Edwards, A.M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hallberg, B.M.; Holmberg-Schiavone, L.; Johansson, I.; Kallas, A.; Karlberg, T.; Kotenyova, T.; Moche, M.; Nyman, T.; Persson, C.; Sagemark, J.; Stenmark, P.; Sundstrom, M.; Thorsell, A.G.; Tresaugues, L.; van den Berg, S.; Weigelt, J.; Nordlund, P.; Structural Genomics Consortium (SGC)
Deposited on : 2007-08-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	 85% 8% • 6%
1	B	334	 77% 13% • 8%

Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4880 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHORIBOSYLFORMYLGLYCINAMIDINE CYCLO-LIGASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	3	0
			2374	1506	407	448	13			
1	B	307	Total	C	N	O	S	0	2	0
			2321	1473	399	436	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	752	GLY	ASP	engineered mutation	UNP P22102
B	752	GLY	ASP	engineered mutation	UNP P22102

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total	O	0	0
			88	88		
3	B	82	Total	O	0	0
			82	82		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PHOSPHORIBOSYLFORMYLGLYCINAMIDINE CYCLO-LIGASE

Chain A:  85% 8% • 6%



• Molecule 1: PHOSPHORIBOSYLFORMYLGLYCINAMIDINE CYCLO-LIGASE

Chain B:  77% 13% • 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.67Å 80.99Å 98.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.99 – 2.10 14.99 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (14.99-2.10) 99.9 (14.99-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.94 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.3.0032	Depositor
R, R_{free}	0.191 , 0.240 0.189 , 0.237	Depositor DCC
R_{free} test set	1909 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.075 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4880	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	2/2425 (0.1%)	0.96	0/3280
1	B	0.87	2/2369 (0.1%)	0.92	2/3204 (0.1%)
All	All	0.87	4/4794 (0.1%)	0.94	2/6484 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	627	VAL	CA-CB	7.09	1.61	1.54
1	A	735	VAL	CA-CB	5.14	1.59	1.53
1	B	681	ILE	CA-CB	5.05	1.59	1.53
1	B	690	ILE	CA-CB	5.03	1.57	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	628	ARG	NE-CZ-NH1	-5.17	116.33	121.50
1	B	689	ASN	N-CA-C	5.15	118.03	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2374	0	2405	15	0
1	B	2321	0	2349	23	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	88	0	0	1	0
3	B	82	0	0	0	0
All	All	4880	0	4754	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:724:GLU:HG2	1:A:782:ILE:HG12	1.41	1.01
1:B:727:MET:HG3	1:B:785:MET:HE1	1.62	0.82
1:A:500:LEU:HD23	1:A:512:ILE:HG12	1.62	0.80
1:B:717:GLN:HG3	1:B:785:MET:HE3	1.67	0.76
1:A:494:ASP:OD2	1:A:523:ASP:OD2	2.08	0.71
1:A:500:LEU:CD2	1:A:512:ILE:HG12	2.21	0.69
1:B:500:LEU:HD23	1:B:512:ILE:HG12	1.75	0.68
1:B:714:SER:O	1:B:718:GLN:HG3	1.97	0.64
1:A:724:GLU:HG2	1:A:782:ILE:CG1	2.25	0.61
1:B:724:GLU:HG2	1:B:781:LEU:HB3	1.82	0.60
1:B:625:SER:O	1:B:629:LYS:HG3	2.01	0.59
1:A:692:ARG:HH11	1:A:692:ARG:HB2	1.67	0.59
1:B:500:LEU:HD23	1:B:512:ILE:CG1	2.32	0.59
1:A:496:VAL:HG12	1:A:499:LYS:HB3	1.86	0.58
1:A:617:SER:HB3	1:A:660:ARG:HG3	1.86	0.58
1:B:608:GLY:O	1:B:743:LYS:HE3	2.04	0.58
1:A:561:LYS:HD3	3:A:2024:HOH:O	2.04	0.56
1:A:486:ASP:OD2	1:A:595:ARG:NH1	2.38	0.56
1:B:603:GLU:H	1:B:603:GLU:CD	2.16	0.55
1:B:617:SER:HB3	1:B:660:ARG:HG3	1.90	0.53
1:A:500:LEU:HD23	1:A:512:ILE:CG1	2.35	0.52
1:B:715:TRP:CE2	1:B:719:GLU:HG3	2.46	0.50
1:A:611:VAL:HG13	1:A:738:VAL:HG13	1.95	0.48
1:B:715:TRP:CZ2	1:B:719:GLU:HG3	2.49	0.47
1:B:708:ARG:NH1	1:B:759:GLU:OE2	2.35	0.45
1:A:625:SER:O	1:A:629:LYS:HG3	2.17	0.45
1:A:593:MET:HG2	1:A:594:GLU:O	2.17	0.44
1:B:714:SER:O	1:B:718:GLN:CG	2.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:656:LEU:O	1:B:657:THR:C	2.60	0.44
1:B:623:GLY:O	1:B:627:VAL:HG23	2.18	0.43
1:B:753:ILE:HG23	1:B:758:GLU:HB2	2.00	0.43
1:B:637:GLN:H	1:B:637:GLN:HG2	1.65	0.43
1:B:533:PHE:HB2	1:B:567:LEU:HG	2.00	0.42
1:B:780:ASN:O	1:B:784:SER:HB3	2.19	0.42
1:A:656:LEU:O	1:A:657:THR:C	2.61	0.42
1:B:689:ASN:OD1	1:B:692:ARG:NH1	2.53	0.41
1:B:724:GLU:HG3	1:B:782:ILE:HG13	2.03	0.40
1:B:620:HIS:CG	1:B:659:THR:HG23	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/334 (94%)	309 (99%)	4 (1%)	0	100	100
1	B	305/334 (91%)	300 (98%)	5 (2%)	0	100	100
All	All	618/668 (92%)	609 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/268 (96%)	243 (95%)	13 (5%)	21	21
1	B	249/268 (93%)	233 (94%)	16 (6%)	16	14
All	All	505/536 (94%)	476 (94%)	29 (6%)	19	17

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	494	ASP
1	A	501	LYS
1	A	593	MET
1	A	594	GLU
1	A	603	GLU
1	A	688[A]	GLU
1	A	688[B]	GLU
1	A	692	ARG
1	A	700	VAL
1	A	725	GLU
1	A	738	VAL
1	A	748	GLN
1	A	792	LEU
1	B	501	LYS
1	B	512	ILE
1	B	567	LEU
1	B	593	MET
1	B	599	LEU
1	B	603	GLU
1	B	606	THR
1	B	637	GLN
1	B	670	LEU
1	B	696	GLU
1	B	705	GLN
1	B	718	GLN
1	B	725	GLU
1	B	738	VAL
1	B	767	VAL
1	B	782	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	622	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	689	ASN
1	B	507	ASN
1	B	622	ASN
1	B	721	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	1786	-	4,4,4	0.29	0	6,6,6	0.58	0
2	SO4	B	1787	-	4,4,4	0.33	0	6,6,6	0.22	0
2	SO4	A	1793	-	4,4,4	0.30	0	6,6,6	0.41	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/334 (94%)	-1.51	0 100 100	16, 31, 42, 49	3 (0%)
1	B	307/334 (91%)	-1.53	0 100 100	16, 32, 41, 49	2 (0%)
All	All	621/668 (92%)	-1.52	0 100 100	16, 32, 42, 49	5 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	1793	5/5	0.99	0.04	17,21,22,27	5
2	SO4	B	1786	5/5	0.99	0.05	22,24,28,29	5
2	SO4	B	1787	5/5	0.99	0.03	48,49,51,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.