



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 07:41 PM UTC

PDB ID : 5VAF / pdb_00005vaf
Title : Crystal structure of accessory secretion protein 1
Authors : Chen, Y.; Rapoport, T.A.; Jeffrey, P.D.
Deposited on : 2017-03-25
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	533	% 68% 27% .
1	B	533	2% 69% 27% ..
1	C	533	2% 65% 30% ..
1	D	533	% 64% 30% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 17260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Accessory Sec system protein Asp1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	510	Total	C	N	O	S	0	0	0
			4292	2776	705	802	9			
1	B	515	Total	C	N	O	S	0	0	0
			4329	2802	710	808	9			
1	C	516	Total	C	N	O	S	0	0	0
			4338	2807	712	810	9			
1	D	511	Total	C	N	O	S	0	0	0
			4298	2779	706	804	9			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	527	VAL	-	expression tag	UNP Q9AET9
A	528	ASP	-	expression tag	UNP Q9AET9
A	529	LYS	-	expression tag	UNP Q9AET9
A	530	LEU	-	expression tag	UNP Q9AET9
A	531	VAL	-	expression tag	UNP Q9AET9
A	532	PRO	-	expression tag	UNP Q9AET9
A	533	ARG	-	expression tag	UNP Q9AET9
B	527	VAL	-	expression tag	UNP Q9AET9
B	528	ASP	-	expression tag	UNP Q9AET9
B	529	LYS	-	expression tag	UNP Q9AET9
B	530	LEU	-	expression tag	UNP Q9AET9
B	531	VAL	-	expression tag	UNP Q9AET9
B	532	PRO	-	expression tag	UNP Q9AET9
B	533	ARG	-	expression tag	UNP Q9AET9
C	527	VAL	-	expression tag	UNP Q9AET9
C	528	ASP	-	expression tag	UNP Q9AET9
C	529	LYS	-	expression tag	UNP Q9AET9
C	530	LEU	-	expression tag	UNP Q9AET9
C	531	VAL	-	expression tag	UNP Q9AET9
C	532	PRO	-	expression tag	UNP Q9AET9
C	533	ARG	-	expression tag	UNP Q9AET9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	527	VAL	-	expression tag	UNP Q9AET9
D	528	ASP	-	expression tag	UNP Q9AET9
D	529	LYS	-	expression tag	UNP Q9AET9
D	530	LEU	-	expression tag	UNP Q9AET9
D	531	VAL	-	expression tag	UNP Q9AET9
D	532	PRO	-	expression tag	UNP Q9AET9
D	533	ARG	-	expression tag	UNP Q9AET9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total O 2 2	0	0
2	B	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Accessory Sec system protein Asp1




• Molecule 1: Accessory Sec system protein Asp1






• Molecule 1: Accessory Sec system protein Asp1



• Molecule 1: Accessory Sec system protein Asp1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.50Å 99.90Å 179.07Å 100.66° 90.07° 95.78°	Depositor
Resolution (Å)	97.65 – 2.77 97.65 – 2.77	Depositor EDS
% Data completeness (in resolution range)	92.7 (97.65-2.77) 92.7 (97.65-2.77)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.208 , 0.282 0.212 , 0.281	Depositor DCC
R_{free} test set	3346 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	46.4	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17260	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7622e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	1/4400 (0.0%)	0.61	0/5962
1	B	0.43	0/4437	0.62	0/6012
1	C	0.44	1/4447 (0.0%)	0.65	1/6027 (0.0%)
1	D	0.44	0/4406	0.68	4/5970 (0.1%)
All	All	0.43	2/17690 (0.0%)	0.64	5/23971 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	467	GLN	CA-C	-5.84	1.49	1.52
1	C	294	TYR	CA-CB	-5.31	1.45	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	97	LEU	N-CA-C	12.94	128.41	109.14
1	D	96	ASP	N-CA-C	-8.46	101.09	112.26
1	C	294	TYR	N-CA-C	-6.24	99.07	109.24
1	D	101	ASP	N-CA-C	6.21	120.72	111.96
1	D	97	LEU	N-CA-CB	-5.76	102.28	111.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4292	0	4164	104	0
1	B	4329	0	4208	113	1
1	C	4338	0	4217	128	1
1	D	4298	0	4169	119	0
2	A	2	0	0	1	0
2	B	1	0	0	0	0
All	All	17260	0	16758	459	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 459 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:236:GLN:HG3	1:C:237:PRO:HD2	1.49	0.94
1:D:285:MET:HE1	1:D:460:SER:HA	1.46	0.94
1:D:414:ARG:NH1	1:D:425:GLU:OE1	2.09	0.85
1:A:108:THR:HG22	1:A:110:PHE:H	1.43	0.83
1:C:257:PHE:HD1	1:C:262:ASN:HD22	1.27	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:473:ASP:OD2	1:C:390:LYS:NZ[1_445]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	506/533 (95%)	462 (91%)	40 (8%)	4 (1%)	16 40
1	B	509/533 (96%)	461 (91%)	44 (9%)	4 (1%)	16 40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	512/533 (96%)	459 (90%)	47 (9%)	6 (1%)	10	30
1	D	507/533 (95%)	465 (92%)	39 (8%)	3 (1%)	21	47
All	All	2034/2132 (95%)	1847 (91%)	170 (8%)	17 (1%)	16	40

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	GLU
1	D	100	GLU
1	A	266	ASN
1	B	352	ASP
1	C	319	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	471/493 (96%)	470 (100%)	1 (0%)	87	96
1	B	476/493 (97%)	473 (99%)	3 (1%)	78	91
1	C	477/493 (97%)	474 (99%)	3 (1%)	78	91
1	D	472/493 (96%)	469 (99%)	3 (1%)	78	91
All	All	1896/1972 (96%)	1886 (100%)	10 (0%)	81	92

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	100	GLU
1	D	101	ASP
1	D	102	ASP
1	B	528	ASP
1	C	230	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	477	GLN
1	D	303	HIS
1	C	169	GLN
1	D	494	GLN
1	D	147	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/533 (95%)	0.14	5 (0%) 79 74	25, 44, 71, 93	0
1	B	515/533 (96%)	0.31	12 (2%) 61 53	29, 49, 88, 118	0
1	C	516/533 (96%)	0.26	10 (1%) 66 59	32, 49, 89, 116	0
1	D	511/533 (95%)	0.18	7 (1%) 73 67	25, 45, 76, 105	0
All	All	2052/2132 (96%)	0.22	34 (1%) 69 62	25, 46, 84, 118	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	143	LEU	3.4
1	B	118	GLY	3.0
1	C	93	GLN	2.8
1	A	363	TRP	2.8
1	D	92	LEU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.