



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2026 – 01:02 AM EDT

PDB ID : 8VAW / pdb_00008vaw
Title : Magnesium catalyzed primer extension product with 14mer primer and mixed phosphorothioate activated G monomer
Authors : Fang, Z.; Szostak, J.W.
Deposited on : 2023-12-11
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)
RNA backbone	3983	1020 (2.10-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	15	
1	B	15	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Magnesium catalyzed nonenzymatic RNA primer extension product.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	15	Total	C	N	O	P	S	0	0	0
			322	149	56	102	14	1			
1	B	15	Total	C	N	O	P	S	0	0	0
			322	149	56	102	14	1			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 2'-DEOXYGUANOSINE-5'-MONOPHOSPHATE (CCD ID: DGP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			22	10	5	6	1		
3	B	1	Total	C	N	O	P	0	0
			22	10	5	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	75	Total	O	0	0
			75	75		
4	B	90	Total	O	0	0
			90	90		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Magnesium catalyzed nonenzymatic RNA primer extension product

Chain A: 



- Molecule 1: Magnesium catalyzed nonenzymatic RNA primer extension product

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	43.57Å 43.57Å 84.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.73 – 1.25 37.73 – 1.25	Depositor EDS
% Data completeness (in resolution range)	94.0 (37.73-1.25) 94.0 (37.73-1.25)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 1.25Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.188 , 0.191 0.192 , 0.197	Depositor DCC
R_{free} test set	1156 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	13.5	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	857	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.2129e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LCG, LCC, MG, G46, DGP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/236	1.13	0/365
1	B	0.81	0/236	1.24	2/365 (0.5%)
All	All	0.81	0/472	1.18	2/730 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	10	A	O3'-P-O5'	-6.51	94.23	104.00
1	B	10	A	C4'-C3'-O3'	-5.75	104.37	113.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	7	U	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	322	0	169	2	0
1	B	322	0	170	0	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	22	0	12	0	0
3	B	22	0	12	5	0
4	A	75	0	0	0	0
4	B	90	0	0	3	0
All	All	857	0	363	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:203:DGP:C5'	4:B:304:HOH:O	2.05	1.05
3:B:203:DGP:H5''	4:B:304:HOH:O	1.56	1.04
1:A:1:LCC:N4	3:B:203:DGP:H2'	2.26	0.51
3:B:203:DGP:P	4:B:301:HOH:O	2.74	0.45
1:A:1:LCC:N4	3:B:203:DGP:C2'	2.81	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	14/15 (93%)	1 (7%)	0
1	B	14/15 (93%)	0	0
All	All	28/30 (93%)	1 (3%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	2	LCC

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	LCG	A	4	1	24,27,28	1.84	6 (25%)	35,42,45	1.27	4 (11%)
1	LCC	B	1	1	21,21,25	2.68	8 (38%)	31,33,40	2.54	13 (41%)
1	G46	B	15	2,1	21,25,26	1.24	3 (14%)	32,37,40	1.44	4 (12%)
1	LCG	B	4	1	24,27,28	1.40	4 (16%)	35,42,45	1.14	2 (5%)
1	LCC	B	3	1	21,24,25	2.14	7 (33%)	29,37,40	1.75	8 (27%)
1	LCC	A	2	1	21,24,25	2.11	5 (23%)	29,37,40	1.55	6 (20%)
1	LCC	A	1	1	21,21,25	3.12	6 (28%)	31,33,40	1.84	9 (29%)
1	LCC	A	3	1	21,24,25	1.66	4 (19%)	29,37,40	0.95	1 (3%)
1	LCC	B	2	1	21,24,25	2.45	9 (42%)	29,37,40	1.62	5 (17%)
1	G46	A	15	2,1	21,25,26	1.77	5 (23%)	32,37,40	1.89	10 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LCG	A	4	1	-	0/8/35/36	0/5/4/4
1	LCC	B	1	1	-	0/7/32/36	0/4/3/3
1	G46	B	15	2,1	-	0/7/25/26	0/3/3/3
1	LCG	B	4	1	-	0/8/35/36	0/5/4/4
1	LCC	B	3	1	-	0/8/35/36	0/4/3/3
1	LCC	A	2	1	-	0/8/35/36	0/4/3/3
1	LCC	A	1	1	-	0/7/32/36	0/4/3/3
1	LCC	A	3	1	-	0/8/35/36	0/4/3/3
1	LCC	B	2	1	-	0/8/35/36	0/4/3/3
1	G46	A	15	2,1	-	0/7/25/26	0/3/3/3

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	LCC	C5-C4	-10.45	1.36	1.44
1	B	2	LCC	C5-C4	-6.64	1.39	1.44
1	A	2	LCC	C5-C4	-5.92	1.39	1.44
1	A	1	LCC	O4'-C1'	5.88	1.51	1.42
1	B	1	LCC	O2'-C6'	5.62	1.53	1.43
1	B	1	LCC	C5-C4	-5.45	1.40	1.44
1	B	3	LCC	C4'-C3'	-5.31	1.47	1.53
1	B	1	LCC	O4'-C1'	4.65	1.49	1.42
1	A	15	G46	C5-N7	-4.54	1.30	1.39
1	A	4	LCG	O2'-C6'	4.42	1.51	1.43
1	B	3	LCC	C5-C4	-4.32	1.40	1.44
1	A	1	LCC	C5M-C5	-4.31	1.40	1.50
1	B	1	LCC	C4'-C3'	-4.22	1.48	1.53
1	A	4	LCG	C4'-C3'	-4.14	1.48	1.53
1	A	2	LCC	O2'-C6'	3.98	1.50	1.43
1	B	1	LCC	C5M-C5	-3.93	1.41	1.50
1	B	2	LCC	C2-N1	-3.92	1.31	1.40
1	B	2	LCC	O4'-C4'	-3.73	1.40	1.45
1	A	4	LCG	O4'-C1'	3.50	1.48	1.42
1	B	4	LCG	C6'-C4'	-3.47	1.46	1.53
1	A	2	LCC	C2'-C3'	-3.44	1.45	1.53
1	A	3	LCC	C5'-C4'	3.42	1.57	1.51
1	B	3	LCC	C5M-C5	-3.35	1.42	1.50
1	A	1	LCC	C4'-C3'	-3.32	1.49	1.53
1	B	2	LCC	C5M-C5	-3.21	1.42	1.50
1	A	3	LCC	C5M-C5	-3.18	1.43	1.50
1	A	3	LCC	O2'-C6'	3.16	1.49	1.43
1	A	15	G46	C5-C6	-3.07	1.33	1.44
1	A	2	LCC	C5M-C5	-3.05	1.43	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	4	LCG	C2'-C3'	-2.87	1.47	1.53
1	B	1	LCC	C6-C5	2.84	1.39	1.34
1	A	15	G46	C4-N9	-2.82	1.30	1.38
1	B	3	LCC	C6-C5	2.77	1.39	1.34
1	A	3	LCC	C4'-C3'	-2.74	1.50	1.53
1	A	1	LCC	C2-N3	-2.70	1.31	1.36
1	B	2	LCC	O2'-C6'	2.67	1.48	1.43
1	B	1	LCC	C2-N1	-2.63	1.34	1.40
1	A	15	G46	O4'-C4'	2.56	1.50	1.45
1	B	4	LCG	C4'-C3'	-2.54	1.50	1.53
1	A	1	LCC	O2'-C6'	2.54	1.48	1.43
1	B	4	LCG	O3'-C3'	2.53	1.47	1.42
1	A	4	LCG	C6'-C4'	-2.53	1.48	1.53
1	B	1	LCC	C6-N1	2.49	1.42	1.38
1	B	2	LCC	C6-N1	2.49	1.42	1.38
1	A	2	LCC	C2-N1	-2.41	1.35	1.40
1	B	3	LCC	O2'-C6'	2.39	1.47	1.43
1	B	4	LCG	C2'-C3'	-2.39	1.48	1.53
1	B	15	G46	O4'-C1'	2.32	1.47	1.42
1	A	15	G46	C2-N1	2.29	1.43	1.37
1	B	3	LCC	C2'-C3'	-2.23	1.48	1.53
1	B	3	LCC	C5'-C4'	-2.20	1.48	1.51
1	B	15	G46	O6-C6	2.16	1.27	1.23
1	A	4	LCG	O3'-C3'	2.11	1.46	1.42
1	B	15	G46	O4'-C4'	2.08	1.49	1.45
1	B	2	LCC	C6-C5	2.02	1.37	1.34
1	B	2	LCC	C4'-C3'	2.01	1.56	1.53
1	B	2	LCC	O4'-C1'	2.01	1.45	1.42

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1	LCC	O5'-C5'-C4'	-5.91	100.12	111.83
1	B	1	LCC	O4'-C1'-N1	5.13	115.57	108.82
1	A	15	G46	N2-C2-N3	4.72	128.89	119.67
1	B	2	LCC	C5-C6-N1	-4.43	118.50	123.31
1	B	1	LCC	O4'-C4'-C5'	4.41	116.67	108.86
1	B	3	LCC	C5M-C5-C6	-4.25	117.10	122.85
1	A	15	G46	O6-C6-N1	-4.16	112.29	120.11
1	B	1	LCC	C5-C6-N1	-3.91	119.06	123.31
1	A	2	LCC	C5-C6-N1	-3.90	119.07	123.31
1	B	3	LCC	C5M-C5-C4	3.87	126.88	120.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	LCC	O5'-C5'-C4'	-3.83	104.24	111.83
1	B	1	LCC	O2'-C6'-C4'	-3.80	96.52	103.81
1	A	2	LCC	O2'-C6'-C4'	-3.68	96.75	103.81
1	B	2	LCC	O2'-C6'-C4'	-3.66	96.79	103.81
1	B	1	LCC	C6-N1-C2	-3.59	116.16	120.95
1	B	1	LCC	N1-C2-N3	3.53	124.92	118.80
1	A	1	LCC	C5-C6-N1	-3.44	119.58	123.31
1	B	1	LCC	O4'-C4'-C3'	-3.34	97.67	102.32
1	B	2	LCC	O2-C2-N3	-3.33	117.09	122.33
1	A	1	LCC	O4'-C1'-C2'	-3.22	97.73	104.64
1	A	4	LCG	O4'-C1'-N9	-3.12	104.72	108.82
1	B	15	G46	O6-C6-N1	-2.97	114.52	120.11
1	B	1	LCC	O2-C2-N3	-2.94	117.70	122.33
1	B	1	LCC	O4'-C1'-C2'	-2.92	98.39	104.64
1	A	15	G46	C2-N3-C4	2.87	117.24	112.30
1	B	1	LCC	O4'-C4'-C6'	2.85	112.01	105.97
1	A	1	LCC	O4'-C4'-C5'	2.81	113.83	108.86
1	B	3	LCC	O2-C2-N3	-2.78	117.94	122.33
1	A	15	G46	N9-C8-N7	-2.72	108.36	113.40
1	B	3	LCC	C4-N3-C2	-2.71	117.05	120.81
1	A	2	LCC	O4'-C4'-C6'	2.68	111.65	105.97
1	A	4	LCG	C4'-O4'-C1'	-2.67	102.88	107.77
1	B	3	LCC	C5-C4-N4	-2.65	117.66	121.39
1	A	1	LCC	C5M-C5-C6	-2.64	119.28	122.85
1	B	15	G46	C5-C4-N3	-2.63	124.21	128.39
1	B	15	G46	C4-C5-N7	-2.60	106.55	110.67
1	A	4	LCG	O4'-C4'-C6'	2.58	111.43	105.97
1	A	15	G46	C8-N7-C5	2.57	108.84	104.26
1	A	15	G46	N1-C2-N3	-2.56	118.64	123.32
1	A	1	LCC	O4'-C4'-C6'	2.50	111.25	105.97
1	B	15	G46	O4'-C1'-N9	-2.48	102.74	108.36
1	B	3	LCC	C5-C6-N1	-2.48	120.62	123.31
1	B	2	LCC	O4'-C1'-N1	2.48	112.08	108.82
1	A	1	LCC	O4'-C4'-C3'	-2.47	98.87	102.32
1	B	3	LCC	C5-C4-N3	2.46	124.28	121.75
1	B	1	LCC	C6'-C4'-C5'	-2.46	108.87	115.97
1	B	1	LCC	C5-C4-N3	-2.42	119.28	121.75
1	B	4	LCG	C2'-C1'-N9	2.27	118.55	114.24
1	A	2	LCC	C5-C4-N3	2.26	124.07	121.75
1	A	1	LCC	C5-C4-N4	-2.24	118.24	121.39
1	A	4	LCG	O2'-C6'-C4'	-2.23	99.53	103.81
1	A	3	LCC	O2'-C6'-C4'	-2.21	99.57	103.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	LCC	N1-C2-N3	2.20	122.61	118.80
1	A	15	G46	C2-N1-C6	-2.16	121.19	125.11
1	B	2	LCC	N1-C2-N3	2.14	122.52	118.80
1	A	2	LCC	C5-C4-N4	-2.13	118.39	121.39
1	A	15	G46	C4-C5-N7	-2.13	107.29	110.67
1	A	2	LCC	C4-N3-C2	-2.09	117.91	120.81
1	B	4	LCG	O4'-C4'-C3'	-2.07	99.43	102.32
1	A	1	LCC	O2'-C6'-C4'	-2.07	99.84	103.81
1	A	15	G46	N2-C2-N1	-2.06	112.42	116.76
1	A	15	G46	O3'-C3'-C4'	-2.04	105.22	111.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1	LCC	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DGP	B	203	-	21,24,25	0.63	0	30,35,38	1.21	2 (6%)
3	DGP	A	203	-	21,24,25	0.44	0	30,35,38	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DGP	B	203	-	-	5/7/21/22	0/3/3/3
3	DGP	A	203	-	-	0/7/21/22	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	203	DGP	C1'-N9-C8	-3.99	118.95	127.91
3	B	203	DGP	C1'-N9-C4	3.66	135.38	125.50

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	203	DGP	C2'-C1'-N9-C8
3	B	203	DGP	C3'-C4'-C5'-O5'
3	B	203	DGP	C2'-C1'-N9-C4
3	B	203	DGP	O4'-C4'-C5'-O5'
3	B	203	DGP	C4'-C5'-O5'-P


There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	203	DGP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	10/15 (66%)	-0.03	0 100 100	11, 14, 17, 17	0
1	B	10/15 (66%)	-0.04	0 100 100	12, 14, 15, 16	0
All	All	20/30 (66%)	-0.03	0 100 100	11, 14, 17, 17	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LCC	A	1	19/23	0.87	0.13	19,24,30,33	0
1	LCC	B	1	19/23	0.87	0.12	20,25,29,30	0
1	LCC	B	2	22/23	0.94	0.09	16,18,21,25	0
1	G46	A	15	23/24	0.95	0.09	15,21,25,27	0
1	LCC	B	3	22/23	0.96	0.06	12,13,15,17	0
1	LCC	A	2	22/23	0.96	0.06	13,14,20,23	0
1	G46	B	15	23/24	0.96	0.07	12,14,19,23	0
1	LCG	B	4	24/25	0.98	0.04	11,12,13,13	0
1	LCC	A	3	22/23	0.98	0.04	10,11,14,15	0
1	LCG	A	4	24/25	0.98	0.05	10,11,11,12	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DGP	A	203	22/23	0.82	0.18	21,26,96,108	0
3	DGP	B	203	22/23	0.84	0.18	21,26,58,64	0
2	MG	B	202	1/1	0.95	0.08	19,19,19,19	1
2	MG	A	202	1/1	0.97	0.19	27,27,27,27	1
2	MG	B	201	1/1	1.00	0.04	11,11,11,11	0
2	MG	A	201	1/1	1.00	0.08	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.