



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 03:08 AM EDT

PDB ID : 8VBK / pdb\_00008vbk  
Title : Structure of bovine anti-HIV Fab ElsE5  
Authors : Stanfield, R.L.; Wilson, I.A.  
Deposited on : 2023-12-12  
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	 2% 92% 7%
2	H	265	 15% 94%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bovine Fab ElsE5 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	214	1566	966	264	331	5	0	0	0

- Molecule 2 is a protein called Bovine Fab ElsE5 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	259	1930	1206	320	395	9	0	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	103	Total	O	0	0
			103	103		
3	H	108	Total	O	0	0
			108	108		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bovine Fab ElsE5 light chain

Chain L: 



- Molecule 2: Bovine Fab ElsE5 heavy chain

Chain H: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.26Å 70.02Å 109.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.86 – 1.89 43.86 – 1.89	Depositor EDS
% Data completeness (in resolution range)	98.7 (43.86-1.89) 98.6 (43.86-1.89)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.204 , 0.245 0.204 , 0.246	Depositor DCC
$R_{free}$ test set	1773 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtrriage
Anisotropy	0.524	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 34.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3707	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.14	0/1599	0.37	0/2184
2	H	0.15	0/1975	0.36	0/2696
All	All	0.14	0/3574	0.36	0/4880

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1566	0	1502	7	0
2	H	1930	0	1869	4	0
3	H	108	0	0	1	0
3	L	103	0	0	0	0
All	All	3707	0	3371	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:211:CYS:SG	3:H:303:HOH:O	2.55	0.64

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:110:LYS:HD2	1:L:141:PRO:HD3	1.82	0.60
2:H:95:VAL:HG12	2:H:97:GLN:HG3	1.96	0.48
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.95	0.47
1:L:30:ASN:CG	2:H:142:TYR:H	2.22	0.47
2:H:169:PRO:HD3	2:H:255:LYS:HE2	1.97	0.46
1:L:92:GLU:C	1:L:94:SER:H	2.26	0.44
1:L:39:ILE:HG12	1:L:84:ALA:HB2	2.02	0.42
2:H:173:SER:O	2:H:177:THR:HG23	2.20	0.42
1:L:167:GLN:OE1	1:L:173:ALA:HB2	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	212/216 (98%)	205 (97%)	7 (3%)	0	100	100
2	H	255/265 (96%)	247 (97%)	8 (3%)	0	100	100
All	All	467/481 (97%)	452 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	178/180 (99%)	175 (98%)	3 (2%)	53	52
2	H	226/230 (98%)	223 (99%)	3 (1%)	61	61
All	All	404/410 (98%)	398 (98%)	6 (2%)	57	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	27	SER
1	L	27(A)	SER
1	L	27(B)	ASN
2	H	80	LEU
2	H	101	LYS
2	H	115	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	126	GLN
2	H	98	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	214/216 (99%)	0.30	5 (2%) 61 65	16, 31, 47, 68	0
2	H	259/265 (97%)	0.64	40 (15%) 5 5	17, 30, 89, 106	0
All	All	473/481 (98%)	0.49	45 (9%) 14 14	16, 30, 77, 106	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	136	LEU	5.5
2	H	137	SER	5.4
2	H	115	PHE	5.4
2	H	107	TRP	5.3
2	H	130	ILE	4.9
1	L	211	CYS	4.6
2	H	111	TRP	4.6
2	H	124	TYR	4.0
2	H	128	THR	3.8
2	H	123	CYS	3.8
2	H	131	LEU	3.7
2	H	127	CYS	3.7
2	H	106	GLY	3.6
2	H	104	PRO	3.5
2	H	110	GLY	3.5
2	H	114	GLY	3.4
2	H	262	CYS	3.3
1	L	30	ASN	3.2
2	H	139	THR	3.1
2	H	109	PHE	3.1
2	H	125	ASP	3.1
2	H	103	CYS	3.1
2	H	261	SER	3.1
2	H	112	ASP	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	H	113	CYS	2.8
2	H	138	PRO	2.8
2	H	105	ASP	2.8
2	H	174	SER	2.8
2	H	140	ASP	2.6
2	H	2	VAL	2.6
2	H	132	SER	2.6
2	H	121	GLU	2.6
2	H	135	THR	2.5
1	L	2	ALA	2.4
2	H	74	PRO	2.4
2	H	122	ASP	2.4
2	H	134	GLN	2.3
2	H	133	SER	2.3
2	H	236	GLY	2.3
2	H	108	SER	2.2
2	H	129	ASP	2.2
1	L	93	ASP	2.2
2	H	126	ASP	2.1
1	L	153	SER	2.0
2	H	141	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.