



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 03:34 PM UTC

PDB ID : 8VBY / pdb_00008vby
EMDB ID : EMD-43128
Title : Structure of the human dopamine transporter in complex with beta-CFT, MRS7292 and divalent zinc
Authors : Srivastava, D.K.; Gouaux, E.
Deposited on : 2023-12-12
Resolution : 3.19 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14455 (2.69 - 3.69)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 4862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sodium-dependent dopamine transporter.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	553	Total	C	N	O	S	0	0
			4371	2909	694	745	23		

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	51	GLY	-	expression tag	UNP Q01959
A	52	PRO	-	expression tag	UNP Q01959
A	53	GLY	-	expression tag	UNP Q01959
A	54	GLY	-	expression tag	UNP Q01959
A	55	ARG	-	expression tag	UNP Q01959
A	56	MET	-	expression tag	UNP Q01959
A	248	TYR	ILE	engineered mutation	UNP Q01959
A	621	ILE	-	expression tag	UNP Q01959

- Molecule 2 is methyl (1R,2S,3S,5S)-3-(4-fluorophenyl)-8-methyl-8-azabicyclo[3.2.1]octane-2-carboxylate (CCD ID: 42L) (formula: C₁₆H₂₀FNO₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
2	A	1	Total	C	F	N	O	0
			20	16	1	1	2	

- Molecule 3 is methyl (1S,2R,3S,4R,5R)-4-{2-[(5-chlorothiophen-2-yl)ethynyl]-6-(methylamino)-9H-purin-9-yl}-2,3-dihydroxybicyclo[3.1.0]hexane-1-carboxylate (CCD ID: A1AAF) (formula: C₂₀H₁₈ClN₅O₄S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
3	A	1	Total	C	Cl	N	O	S	0
			31	20	1	5	4	1	

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				AltConf
4	A	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 5 is CHOLESTEROL HEMISUCCINATE (CCD ID: Y01) (formula: C₃₁H₅₀O₄).




Mol	Chain	Residues	Atoms				AltConf
5	A	1	Total	C	O		0
			35	31	4		
5	A	1	Total	C	O		0
			35	31	4		
5	A	1	Total	C	O		0
			35	31	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			AltConf
5	A	1	Total	C	O	0
			35	31	4	

- Molecule 6 is DECANE (CCD ID: D10) (formula: $C_{10}H_{22}$).




Mol	Chain	Residues	Atoms		AltConf
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	
6	A	1	Total	C	0
			10	10	

- Molecule 7 is PENTANE (CCD ID: LNK) (formula: C_5H_{12}).



Mol	Chain	Residues	Atoms	AltConf
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0
7	A	1	Total C 5 5	0

- Molecule 8 is HEPTANE (CCD ID: HP6) (formula: C₇H₁₆).



Mol	Chain	Residues	Atoms		AltConf
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	
8	A	1	Total	C	0
			7	7	


- Molecule 9 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
9	A	1	Total	Zn	0
			1	1	

- Molecule 10 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		AltConf
10	A	1	Total	Na	0
			1	1	

- Molecule 11 is DODECANE (CCD ID: D12) (formula: $C_{12}H_{26}$).




Mol	Chain	Residues	Atoms		AltConf
11	A	1	Total	C	0
			12	12	
11	A	1	Total	C	0
			12	12	
11	A	1	Total	C	0
			12	12	
11	A	1	Total	C	0
			12	12	
11	A	1	Total	C	0
			12	12	
11	A	1	Total	C	0
			12	12	

- Molecule 12 is PENTADECANE (CCD ID: MYS) (formula: $C_{15}H_{32}$).



Mol	Chain	Residues	Atoms		AltConf
12	A	1	Total	C	0
			15	15	

- Molecule 13 is N-OCTANE (CCD ID: OCT) (formula: C₈H₁₈).



Mol	Chain	Residues	Atoms		AltConf
13	A	1	Total	C	0
			8	8	
13	A	1	Total	C	0
			8	8	


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		AltConf
14	A	10	Total	O	0
			10	10	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sodium-dependent dopamine transporter



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	177494	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	165000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.252	Depositor
Minimum map value	-0.793	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.024	Depositor
Recommended contour level	0.24	Depositor
Map size (Å)	285.31198, 285.31198, 285.31198	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.743, 0.743, 0.743	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D10, OCT, LNK, NAG, D12, MYS, HP6, Y01, A1AAF, ZN, NA, 42L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/4508	0.25	0/6152

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4371	0	4356	39	0
2	A	20	0	20	0	0
3	A	31	0	0	0	0
4	A	14	0	13	0	0
5	A	140	0	196	3	0
6	A	80	0	176	0	0
7	A	35	0	84	0	0
8	A	56	0	128	1	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
11	A	72	0	156	1	0
12	A	15	0	32	0	0
13	A	16	0	36	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	A	10	0	0	0	0
All	All	4862	0	5197	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:VAL:HG13	1:A:360:SER:HB2	1.68	0.76
1:A:79:ASP:OD1	1:A:156:TYR:OH	2.13	0.65
1:A:60:ARG:NH1	1:A:334:SER:OG	2.35	0.60
1:A:421:ASP:OD1	1:A:422:SER:N	2.36	0.57
1:A:221:VAL:O	1:A:237:ARG:NH1	2.39	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	549/571 (96%)	535 (97%)	14 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/475 (98%)	464 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	ASN
1	A	341	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 41 ligands modelled in this entry, 2 are monoatomic - leaving 39 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	D12	A	726	-	11,11,11	0.23	0	10,10,10	0.21	0
11	D12	A	728	-	11,11,11	0.23	0	10,10,10	0.20	0
3	A1AAF	A	702	-	35,35,35	1.06	2 (5%)	39,54,54	1.20	3 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	703	1	14,14,15	0.23	0	17,19,21	0.44	0
11	D12	A	739	-	11,11,11	0.23	0	10,10,10	0.21	0
5	Y01	A	704	-	38,38,38	0.57	0	57,57,57	0.76	0
6	D10	A	705	-	9,9,9	0.23	0	8,8,8	0.21	0
11	D12	A	716	-	11,11,11	0.23	0	10,10,10	0.24	0
5	Y01	A	710	-	38,38,38	0.54	0	57,57,57	0.78	0
5	Y01	A	738	-	38,38,38	0.56	0	57,57,57	0.57	0
12	MYS	A	725	-	14,14,14	0.21	0	13,13,13	0.22	0
8	HP6	A	712	-	6,6,6	0.25	0	5,5,5	0.19	0
6	D10	A	734	-	9,9,9	0.23	0	8,8,8	0.20	0
11	D12	A	736	-	11,11,11	0.23	0	10,10,10	0.20	0
8	HP6	A	715	-	6,6,6	0.26	0	5,5,5	0.17	0
6	D10	A	735	-	9,9,9	0.23	0	8,8,8	0.21	0
6	D10	A	718	-	9,9,9	0.23	0	8,8,8	0.20	0
13	OCT	A	729	-	7,7,7	0.24	0	6,6,6	0.19	0
2	42L	A	701	-	22,22,22	0.60	0	29,32,32	0.72	0
6	D10	A	740	-	9,9,9	0.24	0	8,8,8	0.19	0
8	HP6	A	727	-	6,6,6	0.25	0	5,5,5	0.19	0
5	Y01	A	711	-	38,38,38	0.56	0	57,57,57	0.75	0
7	LNK	A	721	-	4,4,4	0.27	0	3,3,3	0.24	0
7	LNK	A	706	-	4,4,4	0.27	0	3,3,3	0.24	0
6	D10	A	707	-	9,9,9	0.23	0	8,8,8	0.21	0
8	HP6	A	733	-	6,6,6	0.26	0	5,5,5	0.19	0
7	LNK	A	708	-	4,4,4	0.27	0	3,3,3	0.24	0
7	LNK	A	723	-	4,4,4	0.27	0	3,3,3	0.23	0
7	LNK	A	722	-	4,4,4	0.27	0	3,3,3	0.24	0
7	LNK	A	732	-	4,4,4	0.27	0	3,3,3	0.24	0
8	HP6	A	737	-	6,6,6	0.25	0	5,5,5	0.20	0
6	D10	A	741	-	9,9,9	0.23	0	8,8,8	0.19	0
7	LNK	A	709	-	4,4,4	0.27	0	3,3,3	0.23	0
6	D10	A	724	-	9,9,9	0.24	0	8,8,8	0.18	0
8	HP6	A	719	-	6,6,6	0.25	0	5,5,5	0.21	0
8	HP6	A	720	-	6,6,6	0.25	0	5,5,5	0.20	0
8	HP6	A	731	-	6,6,6	0.26	0	5,5,5	0.18	0
13	OCT	A	730	-	7,7,7	0.24	0	6,6,6	0.18	0
11	D12	A	717	-	11,11,11	0.24	0	10,10,10	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	D12	A	726	-	-	0/9/9/9	-
11	D12	A	728	-	-	0/9/9/9	-
3	A1AAF	A	702	-	-	2/16/45/45	0/5/5/5
4	NAG	A	703	1	-	2/6/23/26	0/1/1/1
11	D12	A	739	-	-	0/9/9/9	-
5	Y01	A	704	-	-	8/19/77/77	0/4/4/4
6	D10	A	705	-	-	2/7/7/7	-
11	D12	A	716	-	-	1/9/9/9	-
5	Y01	A	710	-	-	11/19/77/77	0/4/4/4
5	Y01	A	738	-	-	8/19/77/77	0/4/4/4
12	MYS	A	725	-	-	2/12/12/12	-
8	HP6	A	712	-	-	0/4/4/4	-
6	D10	A	734	-	-	2/7/7/7	-
11	D12	A	736	-	-	1/9/9/9	-
8	HP6	A	715	-	-	0/4/4/4	-
6	D10	A	735	-	-	0/7/7/7	-
6	D10	A	718	-	-	0/7/7/7	-
13	OCT	A	729	-	-	0/5/5/5	-
2	42L	A	701	-	-	0/10/35/35	0/4/3/3
6	D10	A	740	-	-	0/7/7/7	-
8	HP6	A	727	-	-	0/4/4/4	-
5	Y01	A	711	-	-	8/19/77/77	0/4/4/4
7	LNK	A	721	-	-	0/2/2/2	-
7	LNK	A	706	-	-	0/2/2/2	-
6	D10	A	707	-	-	1/7/7/7	-
8	HP6	A	733	-	-	0/4/4/4	-
7	LNK	A	708	-	-	0/2/2/2	-
7	LNK	A	723	-	-	0/2/2/2	-
7	LNK	A	722	-	-	0/2/2/2	-
7	LNK	A	732	-	-	0/2/2/2	-
8	HP6	A	737	-	-	0/4/4/4	-
6	D10	A	741	-	-	0/7/7/7	-
7	LNK	A	709	-	-	0/2/2/2	-
6	D10	A	724	-	-	2/7/7/7	-
8	HP6	A	719	-	-	0/4/4/4	-
8	HP6	A	720	-	-	1/4/4/4	-
8	HP6	A	731	-	-	0/4/4/4	-
13	OCT	A	730	-	-	0/5/5/5	-
11	D12	A	717	-	-	2/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	A1AAF	C2-N2	3.46	1.38	1.34
3	A	702	A1AAF	C16-C19	-2.48	1.48	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	A1AAF	C5-N4-C6	-4.04	110.20	116.16
3	A	702	A1AAF	O3-C19-C16	-2.56	121.10	124.50
3	A	702	A1AAF	C17-C18-C13	2.06	105.72	102.95

There are no chirality outliers.

5 of 53 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	A1AAF	C3-C2-N2-C1
3	A	702	A1AAF	N5-C2-N2-C1
5	A	704	Y01	OAG-CAY-OAW-CBC
5	A	738	Y01	CAM-CAY-OAW-CBC
5	A	704	Y01	CAR-CBC-OAW-CAY


There are no ring outliers.



5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	710	Y01	1	0
5	A	738	Y01	1	0
11	A	736	D12	1	0
5	A	711	Y01	1	0
8	A	720	HP6	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1AAF A 702







Ligand Y01 A 738



Ligand 42L A 701





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-43128. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map




6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 209




Y Index: 196




Z Index: 207


6.3.2 Raw map



X Index: 209



Y Index: 196




Z Index: 207


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.24. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_43128_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 24 nm³; this corresponds to an approximate mass of 22 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.313 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.313 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.19	-	-
Author-provided FSC curve	3.19	3.64	3.22
Unmasked-calculated*	3.78	4.23	3.85

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.78 differs from the reported value 3.19 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-43128 and PDB model 8VBY. Per-residue inclusion information can be found in section [3](#) on page [12](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.24 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.24).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.24) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8140	0.5650
A	0.8140	0.5650

