



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:35 PM UTC

PDB ID : 2VC1 / pdb_00002vc1
Title : Feast or famine regulatory protein (Rv3291c) from *M. tuberculosis* complexed with L-Methionine
Authors : Shrivastava, T.; Ramachandran, R.
Deposited on : 2007-09-18
Resolution : 2.75 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	53% 31% 11% . .
1	B	150	5% 48% 33% 12% 5% .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	0	0
			1127	697	209	220	1			
1	B	147	Total	C	N	O	S	0	0	0
			1111	689	203	218	1			

- Molecule 2 is METHIONINE (CCD ID: MET) (formula: C₅H₁₁NO₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			9	5	1	2	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	18	Total	O	0	0
			18	18		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN



• Molecule 1: TRANSCRIPTIONAL REGULATORY PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	100.92Å 100.92Å 99.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.43 – 2.75 71.36 – 2.75	Depositor EDS
% Data completeness (in resolution range)	92.5 (71.43-2.75) 92.5 (71.36-2.75)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 2.76Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.195 , 0.240 0.175 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	44.9	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,-l,-k 0.008 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2286	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.75	13/1141 (1.1%)	1.89	33/1552 (2.1%)
1	B	1.84	17/1125 (1.5%)	1.77	28/1533 (1.8%)
All	All	1.80	30/2266 (1.3%)	1.83	61/3085 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	91	ILE	CA-C	6.76	1.60	1.52
1	A	40	SER	N-CA	-6.67	1.38	1.46
1	B	94	VAL	C-O	-6.27	1.17	1.23
1	B	33	LEU	CA-C	6.23	1.61	1.53
1	B	91	ILE	CA-CB	6.21	1.61	1.53
1	A	146	ARG	N-CA	-6.21	1.38	1.46
1	A	132	ARG	CB-CG	-6.18	1.33	1.52
1	B	144	SER	N-CA	-5.98	1.38	1.46
1	A	141	THR	C-O	-5.92	1.16	1.24
1	A	138	ILE	CA-CB	-5.79	1.47	1.54
1	B	30	ARG	CA-C	5.75	1.60	1.52
1	B	141	THR	C-O	-5.70	1.17	1.24
1	A	14	ARG	CG-CD	5.70	1.69	1.52
1	A	75	PRO	N-CA	5.68	1.53	1.47
1	A	123	GLN	C-O	-5.67	1.17	1.24
1	B	48	ARG	CA-CB	5.61	1.62	1.53
1	B	110	VAL	C-O	5.59	1.29	1.24
1	B	111	ARG	N-CA	5.43	1.52	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	25	SER	CA-C	5.36	1.59	1.52
1	A	115	ALA	CA-CB	-5.34	1.45	1.53
1	A	37	ALA	CA-CB	-5.32	1.44	1.53
1	B	69	ALA	N-CA	5.23	1.52	1.45
1	B	83	ASP	CA-C	5.15	1.59	1.52
1	B	130	ASN	N-CA	5.12	1.53	1.46
1	B	11	ILE	CA-CB	5.10	1.61	1.54
1	A	29	THR	CB-CG2	5.07	1.69	1.52
1	B	32	GLY	N-CA	5.07	1.52	1.45
1	B	29	THR	CA-CB	5.06	1.61	1.53
1	A	145	ASP	C-O	-5.06	1.17	1.23
1	A	62	ALA	C-O	-5.03	1.17	1.24

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	GLN	CA-C-N	-13.50	107.08	120.31
1	A	80	GLN	C-N-CA	-13.50	107.08	120.31
1	A	83	ASP	N-CA-CB	-10.05	97.63	111.00
1	A	71	VAL	CB-CA-C	9.08	121.53	110.73
1	B	129	ALA	CA-C-N	-9.06	108.50	123.37
1	B	129	ALA	C-N-CA	-9.06	108.50	123.37
1	B	130	ASN	CB-CA-C	8.65	124.48	111.76
1	A	74	THR	CA-C-N	8.36	128.88	119.93
1	A	74	THR	C-N-CA	8.36	128.88	119.93
1	B	91	ILE	CB-CA-C	8.12	122.22	111.33
1	B	87	ARG	N-CA-C	7.84	120.72	111.71
1	A	145	ASP	CA-C-N	-7.61	111.51	122.77
1	A	145	ASP	C-N-CA	-7.61	111.51	122.77
1	A	77	ASP	CA-C-N	7.50	129.21	119.84
1	A	77	ASP	C-N-CA	7.50	129.21	119.84
1	B	128	THR	N-CA-C	7.21	118.83	110.97
1	A	126	ARG	N-CA-C	-7.17	103.47	111.28
1	A	37	ALA	N-CA-C	-7.10	103.58	111.82
1	B	31	ALA	N-CA-C	7.04	121.97	112.88
1	A	48	ARG	CA-C-N	-6.96	107.78	121.41
1	A	48	ARG	C-N-CA	-6.96	107.78	121.41
1	A	146	ARG	CB-CA-C	6.67	122.82	111.68
1	A	82	ASP	N-CA-C	6.54	119.35	108.96
1	B	91	ILE	CA-CB-CG2	6.53	121.61	110.50
1	A	119	GLU	N-CA-C	6.42	119.09	111.71
1	A	21	ARG	N-CA-C	-6.33	105.19	113.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	GLN	N-CA-C	-6.15	101.34	110.20
1	B	56	ALA	N-CA-C	-6.07	99.76	109.96
1	B	63	VAL	N-CA-CB	-6.07	104.21	112.28
1	B	134	ARG	CA-C-N	-6.06	113.30	122.81
1	B	134	ARG	C-N-CA	-6.06	113.30	122.81
1	A	81	PRO	N-CA-C	-5.83	102.70	111.57
1	B	145	ASP	CB-CA-C	-5.74	103.42	111.80
1	B	91	ILE	N-CA-CB	5.72	117.17	110.82
1	A	64	GLY	N-CA-C	5.66	123.41	115.43
1	B	145	ASP	N-CA-C	5.59	119.15	111.54
1	B	61	GLU	N-CA-CB	5.53	118.25	110.12
1	A	93	GLU	CA-C-N	-5.52	115.48	122.43
1	A	93	GLU	C-N-CA	-5.52	115.48	122.43
1	B	94	VAL	CB-CA-C	5.51	117.18	111.23
1	A	25	SER	N-CA-C	5.50	116.96	111.07
1	A	118	LEU	CA-C-N	-5.50	112.36	120.28
1	A	118	LEU	C-N-CA	-5.50	112.36	120.28
1	B	146	ARG	CA-C-N	5.49	128.56	120.82
1	B	146	ARG	C-N-CA	5.49	128.56	120.82
1	B	64	GLY	N-CA-C	5.37	122.73	115.32
1	B	129	ALA	N-CA-C	-5.36	101.44	109.79
1	A	53	GLY	N-CA-C	-5.32	103.02	110.63
1	B	74	THR	CB-CA-C	-5.30	103.11	110.16
1	B	129	ALA	CA-C-O	-5.30	114.99	121.67
1	A	30	ARG	N-CA-C	5.29	117.13	111.36
1	A	41	ARG	CB-CA-C	-5.25	102.41	110.81
1	B	112	VAL	CB-CA-C	5.22	118.77	111.40
1	A	105	SER	N-CA-C	5.21	119.63	113.12
1	B	135	SER	N-CA-C	5.21	118.05	109.72
1	A	131	VAL	N-CA-CB	-5.20	101.74	111.62
1	B	94	VAL	N-CA-CB	5.19	117.41	111.39
1	A	81	PRO	CA-C-N	-5.15	115.35	122.30
1	A	81	PRO	C-N-CA	-5.15	115.35	122.30
1	B	87	ARG	CA-CB-CG	-5.11	103.87	114.10
1	B	29	THR	N-CA-CB	5.04	117.30	109.98

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	7	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1127	0	1130	38	0
1	B	1111	0	1104	47	0
2	A	9	0	8	3	0
3	A	21	0	0	4	0
3	B	18	0	0	4	0
All	All	2286	0	2242	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ARG:HH11	1:A:14:ARG:HB3	1.21	1.05
1:A:14:ARG:HH11	1:A:14:ARG:CB	1.72	1.03
1:B:42:VAL:O	1:B:46:GLU:HG3	1.67	0.94
1:B:23:THR:HG22	3:B:2001:HOH:O	1.66	0.93
1:A:14:ARG:HB3	1:A:14:ARG:NH1	1.83	0.92
1:A:26:GLU:HG3	3:A:2002:HOH:O	1.70	0.89
1:B:24:LEU:HD22	1:B:39:GLN:HG3	1.55	0.88
1:A:124:ARG:HD3	3:A:2017:HOH:O	1.72	0.86
1:B:24:LEU:HD12	3:B:2001:HOH:O	1.74	0.85
1:B:91:ILE:C	1:B:91:ILE:HD12	2.02	0.85
1:A:99:SER:HB2	2:A:1151:MET:CE	2.15	0.77
1:A:14:ARG:HH11	1:A:14:ARG:CG	1.98	0.76
1:A:99:SER:HB2	2:A:1151:MET:HE1	1.68	0.75
1:B:63:VAL:O	1:B:63:VAL:HG12	1.88	0.72
1:B:44:ARG:HH11	1:B:44:ARG:HB3	1.57	0.69
1:B:59:ASN:HD22	1:B:62:ALA:H	1.42	0.68
1:B:44:ARG:HB3	1:B:44:ARG:NH1	2.09	0.67
1:B:59:ASN:ND2	1:B:62:ALA:H	1.94	0.66
1:A:116:ARG:NH1	1:A:119:GLU:OE2	2.29	0.66
1:B:72:ALA:HB1	1:B:104:GLU:HG2	1.77	0.66
1:B:91:ILE:HD12	1:B:92:GLU:N	2.12	0.64
1:A:14:ARG:NH1	1:A:14:ARG:CG	2.55	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ARG:NH2	1:B:89:GLU:O	2.30	0.62
1:B:68:SER:OG	1:B:111:ARG:HD3	1.99	0.62
1:B:124:ARG:O	1:B:128:THR:HG23	2.00	0.61
1:A:90:HIS:HB3	3:A:2010:HOH:O	2.00	0.61
1:B:91:ILE:HD11	1:B:93:GLU:HG2	1.80	0.61
1:A:124:ARG:O	1:A:128:THR:HG23	2.00	0.61
1:B:63:VAL:O	1:B:63:VAL:CG1	2.50	0.59
1:A:124:ARG:O	1:A:128:THR:CG2	2.49	0.59
1:A:104:GLU:CD	1:A:134:ARG:HH21	2.14	0.55
1:B:42:VAL:O	1:B:46:GLU:CG	2.49	0.55
1:B:41:ARG:HA	1:B:44:ARG:NH1	2.22	0.55
1:B:72:ALA:CB	1:B:104:GLU:HG2	2.36	0.55
1:A:87:ARG:HB3	1:A:87:ARG:HH11	1.73	0.53
1:B:8:ILE:HG21	1:B:41:ARG:HH21	1.75	0.52
1:A:14:ARG:CB	1:A:14:ARG:NH1	2.49	0.52
1:B:59:ASN:HD22	1:B:62:ALA:CB	2.23	0.51
1:B:11:ILE:O	1:B:15:GLU:HB2	2.11	0.50
1:A:25:SER:O	1:A:29:THR:HG23	2.12	0.50
1:B:59:ASN:HD21	1:B:61:GLU:HG2	1.77	0.49
1:B:122:LEU:O	1:B:126:ARG:HG3	2.12	0.49
1:A:24:LEU:HB3	1:A:35:VAL:HG13	1.95	0.49
1:A:21:ARG:NH1	1:B:20:GLY:O	2.46	0.48
1:A:73:ILE:HD12	1:A:131:VAL:HG21	1.95	0.48
1:B:24:LEU:CD2	1:B:39:GLN:HG3	2.37	0.47
1:B:90:HIS:HB3	3:B:2005:HOH:O	2.14	0.47
1:B:41:ARG:O	1:B:45:LEU:N	2.28	0.47
1:B:28:ALA:HA	1:B:38:VAL:HG21	1.97	0.47
1:A:100:VAL:HG22	1:B:136:THR:HG22	1.96	0.46
1:A:146:ARG:NH1	1:B:94:VAL:O	2.48	0.46
1:B:21:ARG:O	1:B:22:ALA:C	2.58	0.46
1:A:122:LEU:O	1:A:126:ARG:HG3	2.16	0.46
1:B:146:ARG:HH11	1:B:146:ARG:HB2	1.81	0.46
1:A:45:LEU:HD12	1:A:45:LEU:HA	1.68	0.46
1:B:14:ARG:HA	1:B:149:ILE:HD13	1.97	0.46
1:B:75:PRO:HA	1:B:131:VAL:HG12	1.98	0.46
1:B:124:ARG:O	1:B:128:THR:CG2	2.63	0.45
1:B:6:ASP:O	1:B:9:ASP:N	2.50	0.45
1:B:28:ALA:CA	1:B:38:VAL:HG21	2.47	0.44
1:A:121:LEU:O	1:A:125:ILE:HG13	2.18	0.43
1:B:41:ARG:H	1:B:41:ARG:HG2	1.64	0.43
1:A:73:ILE:HG22	1:A:133:THR:HG22	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:LEU:HD23	1:B:16:LEU:HA	1.72	0.42
1:B:27:LEU:HD23	1:B:27:LEU:HA	1.97	0.42
1:A:8:ILE:HG13	1:A:41:ARG:NH2	2.34	0.42
1:A:127:THR:HG22	3:A:2015:HOH:O	2.19	0.42
1:A:149:ILE:HA	1:A:150:PRO:HD3	1.88	0.42
1:A:8:ILE:HG21	1:A:33:LEU:HD11	2.01	0.42
1:A:21:ARG:NH1	1:A:21:ARG:HG2	2.35	0.42
1:A:124:ARG:O	1:A:128:THR:HG22	2.18	0.41
1:A:67:LEU:HD23	1:A:67:LEU:HA	1.89	0.41
1:A:83:ASP:HB3	1:A:87:ARG:HD2	2.02	0.41
1:B:8:ILE:CG2	1:B:41:ARG:HH21	2.33	0.41
1:A:76:LEU:HA	1:A:76:LEU:HD23	1.54	0.41
1:B:24:LEU:CD1	3:B:2001:HOH:O	2.46	0.41
1:B:51:VAL:O	1:B:51:VAL:CG2	2.68	0.41
1:A:98:TYR:O	1:A:106:TYR:HA	2.21	0.41
1:A:99:SER:HB2	2:A:1151:MET:HE3	1.97	0.41
1:B:59:ASN:HD22	1:B:62:ALA:N	2.14	0.41
1:B:91:ILE:C	1:B:91:ILE:CD1	2.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/150 (97%)	140 (97%)	5 (3%)	0	100	100
1	B	145/150 (97%)	128 (88%)	11 (8%)	6 (4%)	2	3
All	All	290/300 (97%)	268 (92%)	16 (6%)	6 (2%)	5	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	25	SER
1	B	28	ALA
1	B	26	GLU
1	B	30	ARG
1	B	7	ASP
1	B	27	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/126 (96%)	96 (79%)	25 (21%)	1	2
1	B	118/126 (94%)	93 (79%)	25 (21%)	1	2
All	All	239/252 (95%)	189 (79%)	50 (21%)	1	2

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	14	ARG
1	A	21	ARG
1	A	24	LEU
1	A	29	THR
1	A	38	VAL
1	A	44	ARG
1	A	45	LEU
1	A	58	ILE
1	A	61	GLU
1	A	67	LEU
1	A	71	VAL
1	A	74	THR
1	A	75	PRO
1	A	80	GLN
1	A	87	ARG
1	A	94	VAL
1	A	100	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	110	VAL
1	A	112	VAL
1	A	127	THR
1	A	128	THR
1	A	131	VAL
1	A	132	ARG
1	A	146	ARG
1	B	5	LEU
1	B	6	ASP
1	B	11	ILE
1	B	15	GLU
1	B	23	THR
1	B	24	LEU
1	B	25	SER
1	B	35	VAL
1	B	41	ARG
1	B	51	VAL
1	B	52	GLN
1	B	61	GLU
1	B	67	LEU
1	B	91	ILE
1	B	92	GLU
1	B	93	GLU
1	B	94	VAL
1	B	100	VAL
1	B	104	GLU
1	B	110	VAL
1	B	112	VAL
1	B	120	ASP
1	B	128	THR
1	B	130	ASN
1	B	146	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	B	39	GLN
1	B	59	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MET	A	1151	-	7,8,8	0.73	0	5,9,9	1.54	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MET	A	1151	-	-	1/8/8/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1151	MET	OXT-C-O	-3.33	116.52	124.08

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1151	MET	CB-CG-SD-CE

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1151	MET	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/150 (98%)	-0.32	2 (1%) 73 73	16, 36, 60, 77	0
1	B	147/150 (98%)	-0.05	8 (5%) 31 31	14, 41, 83, 97	0
All	All	294/300 (98%)	-0.19	10 (3%) 48 46	14, 38, 80, 97	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	4	ALA	4.3
1	B	48	ARG	3.5
1	A	81	PRO	3.4
1	B	24	LEU	3.2
1	A	4	ALA	3.0
1	B	8	ILE	2.6
1	B	26	GLU	2.5
1	B	44	ARG	2.4
1	B	35	VAL	2.3
1	B	30	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MET	A	1151	9/9	0.98	0.13	42,43,49,54	0

6.5 Other polymers

There are no such residues in this entry.