



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:02 PM UTC

PDB ID : 1VDG / pdb_00001vdg
Title : Crystal structure of LIR1.01, one of the alleles of LIR1
Authors : Shiroishi, M.; Rasubala, L.; Kuroki, K.; Amano, K.; Tsuchiya, N.; Tokunaga, K.; Kohda, D.; Maenaka, K.
Deposited on : 2004-03-22
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	 50% 40% 5% 5%
1	B	197	 44% 44% 7% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leukocyte immunoglobulin-like receptor subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1478	940	257	275	6			
1	B	187	Total	C	N	O	S	0	0	0
			1471	935	252	278	6			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	29	Total	O	0	0
			29	29		
2	B	32	Total	O	0	0
			32	32		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Leukocyte immunoglobulin-like receptor subfamily B member 1



- Molecule 1: Leukocyte immunoglobulin-like receptor subfamily B member 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.97Å 93.22Å 57.77Å 90.00° 107.82° 90.00°	Depositor
Resolution (Å)	19.98 – 2.80 19.98 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (19.98-2.80) 84.8 (19.98-2.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.06 (at 2.76Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.241 , 0.301 0.221 , 0.239	Depositor DCC
R_{free} test set	519 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	13.4	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 15.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.30$, $\langle L^2 \rangle = 0.13$	Xtriage
Estimated twinning fraction	0.226 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3010	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1520	1.33	25/2068 (1.2%)
1	B	0.70	2/1510 (0.1%)	1.14	13/2052 (0.6%)
All	All	0.62	2/3030 (0.1%)	1.24	38/4120 (0.9%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	106	ALA	CA-CB	10.88	1.70	1.53
1	B	107	GLN	CA-C	7.35	1.60	1.52

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	32	THR	N-CA-C	23.80	138.44	108.45
1	A	33	GLN	N-CA-C	10.97	134.16	110.80
1	A	32	THR	CA-C-N	10.59	141.77	121.54
1	A	32	THR	C-N-CA	10.59	141.77	121.54
1	B	155	SER	N-CA-C	9.14	130.27	110.80
1	A	20	SER	CA-C-N	8.34	128.48	120.31
1	A	20	SER	C-N-CA	8.34	128.48	120.31
1	A	109	SER	N-CA-C	7.79	117.09	108.14
1	B	62	ILE	CA-C-N	6.84	126.36	119.24
1	B	62	ILE	C-N-CA	6.84	126.36	119.24
1	B	86	GLU	N-CA-C	-6.67	102.21	110.41
1	A	148	GLN	N-CA-C	-6.51	102.50	110.31
1	A	50	ILE	CA-C-N	6.12	127.48	119.84
1	A	50	ILE	C-N-CA	6.12	127.48	119.84
1	A	32	THR	CB-CA-C	-5.97	99.35	113.02
1	B	107	GLN	C-N-CD	-5.91	107.59	120.60
1	A	144	CYS	N-CA-C	5.67	118.93	110.14
1	A	32	THR	O-C-N	5.54	129.87	122.73
1	A	34	GLU	N-CA-C	5.44	122.39	110.80
1	B	187	LEU	N-CA-C	-5.31	102.75	110.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	32	THR	CA-C-O	5.31	126.61	121.14
1	A	187	LEU	N-CA-C	-5.27	102.81	110.08
1	B	181	SER	CA-C-N	5.25	126.40	119.84
1	B	181	SER	C-N-CA	5.25	126.40	119.84
1	B	109	SER	CA-C-N	5.23	124.95	119.82
1	B	109	SER	C-N-CA	5.23	124.95	119.82
1	A	16	ILE	N-CA-C	5.14	115.31	108.11
1	A	165	SER	CA-C-N	5.12	124.73	119.05
1	A	165	SER	C-N-CA	5.12	124.73	119.05
1	A	31	GLU	CA-C-N	-5.10	113.43	122.20
1	A	31	GLU	C-N-CA	-5.10	113.43	122.20
1	B	89	ASP	N-CA-C	-5.08	103.08	110.08
1	A	181	SER	CA-C-N	5.06	126.16	119.84
1	A	181	SER	C-N-CA	5.06	126.16	119.84
1	A	189	SER	N-CA-C	-5.04	104.03	110.53
1	A	86	GLU	N-CA-C	-5.02	103.90	110.53
1	B	107	GLN	CA-C-N	5.00	139.01	127.00
1	B	107	GLN	C-N-CA	5.00	139.01	127.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1478	0	1440	85	0
1	B	1471	0	1427	100	0
2	A	29	0	0	0	0
2	B	32	0	0	1	0
All	All	3010	0	2867	164	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All (164) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:GLN:O	1:A:35:TYR:N	1.95	0.98
1:B:171:TRP:HB3	1:B:191:LEU:HD11	1.46	0.96
1:A:31:GLU:HG2	1:B:119:ILE:HD11	1.51	0.92
1:B:66:THR:H	1:B:69:HIS:HD2	1.17	0.90
1:B:66:THR:H	1:B:69:HIS:CD2	1.93	0.86
1:B:84:ARG:HG2	1:B:84:ARG:HH11	1.41	0.84
1:A:178:ASP:OD1	1:A:180:ASN:HB2	1.81	0.80
1:A:66:THR:H	1:A:69:HIS:HD2	1.33	0.76
1:A:31:GLU:C	1:A:32:THR:CG2	2.58	0.76
1:A:2:HIS:H	1:B:152:ARG:HH22	1.33	0.74
1:A:27:GLN:HG2	1:B:145:LEU:HD13	1.69	0.74
1:A:31:GLU:C	1:A:32:THR:HG22	2.13	0.73
1:A:34:GLU:O	1:A:77:TYR:HA	1.89	0.71
1:A:147:SER:HB2	1:B:29:GLY:HA2	1.72	0.71
1:B:6:PRO:HG3	1:B:77:TYR:CD2	2.26	0.70
1:B:104:LEU:HG	1:B:192:LEU:HD22	1.72	0.70
1:B:66:THR:N	1:B:69:HIS:HD2	1.89	0.70
1:A:171:TRP:CD1	1:A:193:GLU:HG3	2.26	0.70
1:B:170:TRP:HE3	1:B:172:TYR:HH	1.39	0.70
1:B:99:TYR:H	1:B:186:SER:HA	1.57	0.69
1:B:8:LEU:HD12	1:B:25:ARG:O	1.92	0.69
1:B:66:THR:HB	2:B:205:HOH:O	1.93	0.69
1:A:33:GLN:HA	1:A:56:LYS:HA	1.73	0.68
1:A:179:SER:O	1:A:182:PRO:HD3	1.93	0.68
1:B:84:ARG:HG2	1:B:84:ARG:NH1	2.09	0.68
1:A:171:TRP:NE1	1:A:193:GLU:HG3	2.09	0.67
1:B:149:PRO:HA	1:B:156:ARG:O	1.96	0.66
1:B:94:VAL:HG11	1:B:185:TRP:CD2	2.32	0.65
1:B:166:PRO:HG3	1:B:196:VAL:HG11	1.78	0.65
1:A:2:HIS:O	1:A:4:PRO:HD3	1.97	0.64
1:A:66:THR:H	1:A:69:HIS:CD2	2.15	0.64
1:A:28:GLY:O	1:B:146:ASN:O	2.16	0.63
1:A:32:THR:OG1	1:A:33:GLN:HG2	1.99	0.63
1:A:29:GLY:HA2	1:B:145:LEU:HD12	1.80	0.62
1:A:31:GLU:O	1:A:32:THR:HG22	1.98	0.62
1:A:136:GLU:OE2	1:A:173:ARG:HG3	2.00	0.62
1:A:2:HIS:O	1:B:152:ARG:NH2	2.33	0.61
1:A:91:LEU:C	1:A:91:LEU:HD23	2.25	0.61
1:B:6:PRO:HG3	1:B:77:TYR:HD2	1.66	0.61
1:A:173:ARG:HE	1:A:188:PRO:HB2	1.66	0.60
1:B:52:GLN:O	1:B:56:LYS:HG3	2.02	0.60
1:B:99:TYR:HB2	1:B:186:SER:CB	2.32	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:ASN:OD1	1:B:162:GLY:HA2	2.03	0.59
1:A:31:GLU:CG	1:B:119:ILE:HD11	2.30	0.58
1:A:170:TRP:HB2	1:A:194:LEU:HB2	1.86	0.58
1:A:2:HIS:H	1:B:152:ARG:NH2	2.01	0.58
1:A:72:ARG:NH2	1:A:90:PRO:HG3	2.19	0.58
1:A:171:TRP:HB3	1:A:191:LEU:HD11	1.86	0.57
1:B:106:ALA:HB1	1:B:109:SER:C	2.30	0.57
1:A:31:GLU:HG2	1:B:119:ILE:CD1	2.29	0.56
1:A:4:PRO:CD	1:B:148:GLN:HG2	2.35	0.56
1:A:36:ARG:HD2	1:A:76:TYR:CE1	2.41	0.55
1:A:66:THR:N	1:A:69:HIS:HD2	2.05	0.55
1:B:2:HIS:O	1:B:4:PRO:HD3	2.08	0.54
1:A:40:GLU:O	1:A:41:LYS:HB2	2.08	0.54
1:A:112:VAL:HG11	1:A:118:VAL:HB	1.89	0.54
1:B:149:PRO:HB2	1:B:152:ARG:HA	1.90	0.53
1:A:18:GLN:HG2	1:A:19:GLY:N	2.23	0.53
1:B:170:TRP:HB2	1:B:194:LEU:HB2	1.91	0.53
1:B:9:TRP:CE2	1:B:25:ARG:HB2	2.44	0.53
1:B:175:TYR:CE2	1:B:188:PRO:HB3	2.44	0.53
1:B:149:PRO:HB2	1:B:152:ARG:CA	2.39	0.52
1:A:16:ILE:HD11	1:A:65:ILE:HB	1.90	0.52
1:A:2:HIS:N	1:B:152:ARG:HH22	2.04	0.52
1:B:36:ARG:HG2	1:B:36:ARG:HH11	1.75	0.52
1:A:31:GLU:HB2	1:A:32:THR:HG23	1.91	0.52
1:A:53:GLU:H	1:A:53:GLU:CD	2.17	0.52
1:B:196:VAL:O	1:B:197:LEU:HD12	2.09	0.52
1:B:170:TRP:HE3	1:B:172:TYR:OH	1.92	0.52
1:B:24:LEU:HD11	1:B:93:LEU:HD22	1.91	0.51
1:B:6:PRO:HG3	1:B:77:TYR:CE2	2.45	0.51
1:A:30:GLN:O	1:A:31:GLU:C	2.54	0.51
1:A:31:GLU:HA	1:B:160:SER:CB	2.41	0.50
1:A:122:CYS:O	1:A:156:ARG:HA	2.11	0.50
1:B:111:VAL:HG22	1:B:195:LEU:HB2	1.93	0.50
1:B:36:ARG:HG2	1:B:36:ARG:NH1	2.27	0.50
1:A:4:PRO:HD3	1:B:148:GLN:HG2	1.93	0.50
1:A:34:GLU:HB3	1:A:78:GLY:H	1.76	0.50
1:B:111:VAL:HA	1:B:195:LEU:O	2.12	0.50
1:A:171:TRP:HE1	1:A:193:GLU:HG3	1.75	0.50
1:B:165:SER:H	1:B:170:TRP:HZ2	1.58	0.50
1:B:161:VAL:CG2	1:B:161:VAL:O	2.59	0.49
1:A:121:GLN:HG2	1:A:123:ASP:OD1	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:TYR:H	1:A:186:SER:HA	1.78	0.49
1:B:106:ALA:CB	1:B:110:PRO:N	2.75	0.49
1:A:39:ARG:HD2	1:A:71:GLY:HA3	1.94	0.49
1:A:34:GLU:HB3	1:A:78:GLY:N	2.28	0.49
1:A:148:GLN:NE2	1:B:33:GLN:HE22	2.10	0.48
1:B:97:GLY:HA2	1:B:187:LEU:HD23	1.95	0.48
1:B:162:GLY:O	1:B:163:PRO:C	2.57	0.48
1:B:99:TYR:HB2	1:B:186:SER:HB2	1.95	0.48
1:B:121:GLN:NE2	1:B:156:ARG:CZ	2.77	0.48
1:B:92:GLU:HB3	1:B:183:TYR:HE2	1.78	0.47
1:A:4:PRO:HD2	1:B:148:GLN:HG2	1.97	0.47
1:A:5:LYS:CG	1:A:88:SER:HA	2.45	0.47
1:A:154:SER:O	1:A:155:SER:CB	2.62	0.47
1:A:16:ILE:HD11	1:A:65:ILE:CB	2.45	0.47
1:A:112:VAL:HG23	1:A:164:VAL:HG21	1.97	0.47
1:A:26:CYS:O	1:A:58:GLY:HA3	2.15	0.46
1:B:28:GLY:N	1:B:35:TYR:OH	2.48	0.46
1:B:39:ARG:HH21	1:B:42:LYS:HG2	1.80	0.46
1:A:153:GLY:HA3	1:A:156:ARG:HD2	1.96	0.46
1:B:171:TRP:CD1	1:B:193:GLU:HG3	2.51	0.46
1:B:112:VAL:HG11	1:B:118:VAL:HB	1.98	0.46
1:A:16:ILE:HD11	1:A:65:ILE:CG2	2.46	0.46
1:B:182:PRO:HG2	1:B:183:TYR:CE1	2.51	0.46
1:B:46:TRP:CZ2	1:B:62:ILE:HD11	2.51	0.46
1:A:89:ASP:OD1	1:B:72:ARG:NH2	2.49	0.45
1:B:179:SER:O	1:B:182:PRO:HD3	2.17	0.45
1:B:181:SER:N	1:B:182:PRO:CD	2.77	0.45
1:A:36:ARG:N	1:A:76:TYR:O	2.44	0.45
1:B:16:ILE:CG2	1:B:20:SER:HB2	2.47	0.45
1:B:80:ASP:OD1	1:B:84:ARG:NE	2.50	0.45
1:B:22:VAL:HG11	1:B:65:ILE:HD12	1.98	0.45
1:B:51:PRO:O	1:B:55:VAL:HG23	2.16	0.45
1:B:166:PRO:HG3	1:B:196:VAL:CG1	2.45	0.45
1:A:31:GLU:HA	1:B:160:SER:HB3	1.97	0.44
1:B:36:ARG:HD3	1:B:76:TYR:OH	2.18	0.44
1:A:158:ILE:O	1:B:29:GLY:O	2.36	0.44
1:B:5:LYS:HG2	1:B:88:SER:HA	1.99	0.44
1:B:103:THR:HG22	1:B:123:ASP:HB3	1.99	0.44
1:A:5:LYS:HG2	1:A:88:SER:HA	1.98	0.44
1:A:112:VAL:CG2	1:A:164:VAL:HG21	2.47	0.44
1:B:2:HIS:CE1	1:B:4:PRO:HG3	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:GLU:OE1	1:B:72:ARG:HD3	2.18	0.43
1:A:2:HIS:ND1	1:B:179:SER:HB3	2.34	0.43
1:A:16:ILE:HD11	1:A:65:ILE:HG21	2.00	0.43
1:A:173:ARG:NH1	1:A:191:LEU:HB2	2.33	0.43
1:B:92:GLU:HB3	1:B:183:TYR:CE2	2.53	0.43
1:B:66:THR:N	1:B:69:HIS:CD2	2.72	0.43
1:B:148:GLN:HB3	1:B:152:ARG:HH21	1.83	0.43
1:B:109:SER:HA	1:B:110:PRO:HD3	1.85	0.42
1:A:16:ILE:HD13	1:A:22:VAL:CG1	2.49	0.42
1:A:30:GLN:O	1:A:32:THR:N	2.53	0.42
1:B:5:LYS:CG	1:B:88:SER:HA	2.49	0.42
1:B:77:TYR:O	1:B:84:ARG:HA	2.19	0.42
1:A:51:PRO:O	1:A:52:GLN:C	2.61	0.42
1:B:70:ALA:HB1	1:B:183:TYR:HB2	2.00	0.42
1:A:177:TYR:CD1	1:A:177:TYR:C	2.97	0.42
1:B:9:TRP:CD2	1:B:25:ARG:HB2	2.54	0.42
1:A:30:GLN:HB3	1:B:158:ILE:O	2.19	0.42
1:B:108:PRO:O	1:B:109:SER:HB3	2.20	0.42
1:B:13:GLY:O	1:B:16:ILE:HD11	2.20	0.41
1:A:9:TRP:HA	1:A:91:LEU:HD12	2.03	0.41
1:A:51:PRO:O	1:A:53:GLU:N	2.53	0.41
1:A:120:LEU:HD23	1:A:120:LEU:HA	1.89	0.41
1:B:20:SER:HB3	1:B:21:PRO:HD2	2.02	0.41
1:A:20:SER:HA	1:A:21:PRO:HD3	1.84	0.41
1:B:103:THR:CG2	1:B:123:ASP:HB3	2.50	0.41
1:B:106:ALA:HB3	1:B:110:PRO:CD	2.50	0.41
1:A:16:ILE:CG1	1:A:65:ILE:HG21	2.50	0.41
1:A:5:LYS:HG3	1:A:87:SER:O	2.21	0.41
1:A:32:THR:HB	1:A:33:GLN:H	1.35	0.41
1:A:147:SER:HB3	1:A:159:PHE:CZ	2.55	0.41
1:B:106:ALA:HB3	1:B:110:PRO:HD3	2.03	0.41
1:B:106:ALA:HB2	1:B:110:PRO:HG3	2.02	0.41
1:A:16:ILE:O	1:A:95:VAL:HA	2.22	0.40
1:A:154:SER:O	1:A:155:SER:HB3	2.20	0.40
1:A:182:PRO:HG2	1:A:183:TYR:CE1	2.56	0.40
1:B:106:ALA:CB	1:B:109:SER:C	2.92	0.40
1:B:116:GLY:O	1:B:164:VAL:HG23	2.21	0.40
1:B:178:ASP:OD1	1:B:180:ASN:HB2	2.21	0.40
1:A:40:GLU:OE2	1:A:74:ARG:NE	2.38	0.40
1:B:31:GLU:C	1:B:32:THR:HG22	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/197 (92%)	164 (91%)	14 (8%)	3 (2%)	7	25
1	B	177/197 (90%)	166 (94%)	8 (4%)	3 (2%)	7	25
All	All	358/394 (91%)	330 (92%)	22 (6%)	6 (2%)	7	25

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	GLU
1	A	151	ALA
1	B	155	SER
1	B	163	PRO
1	A	52	GLN
1	B	162	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/171 (95%)	152 (93%)	11 (7%)	15	42
1	B	163/171 (95%)	151 (93%)	12 (7%)	13	37
All	All	326/342 (95%)	303 (93%)	23 (7%)	13	39

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ILE
1	A	17	THR
1	A	32	THR
1	A	136	GLU
1	A	143	GLN
1	A	145	LEU
1	A	148	GLN
1	A	156	ARG
1	A	158	ILE
1	A	160	SER
1	A	169	ARG
1	B	32	THR
1	B	34	GLU
1	B	43	THR
1	B	79	SER
1	B	84	ARG
1	B	91	LEU
1	B	108	PRO
1	B	145	LEU
1	B	148	GLN
1	B	155	SER
1	B	160	SER
1	B	161	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	HIS
1	A	117	ASN
1	A	121	GLN
1	B	33	GLN
1	B	69	HIS
1	B	113	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	187/197 (94%)	-1.64	0 100 100	12, 28, 50, 57	0
1	B	187/197 (94%)	-1.60	0 100 100	10, 30, 57, 81	0
All	All	374/394 (94%)	-1.62	0 100 100	10, 29, 53, 81	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.