



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 06:47 PM UTC

PDB ID : 1VDK / pdb_00001vdk
Title : Crystal structure of fumarase from thermus thermophilus HB8
Authors : Mizutani, H.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2004-03-23
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	11% 79% 19%
1	B	466	8% 78% 20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate hydratase class II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	458	Total	C	N	O	S	0	0	0
			3515	2231	617	653	14			
1	B	457	Total	C	N	O	S	0	0	0
			3517	2234	617	652	14			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	357	Total	O	0	0
			357	357		
2	B	291	Total	O	0	0
			291	291		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fumarate hydratase class II



• Molecule 1: Fumarate hydratase class II



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	125.34Å 140.12Å 135.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 30.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-1.80) 99.9 (30.00-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 1.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.185 , 0.201 0.185 , 0.190	Depositor DCC
R_{free} test set	5490 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7680	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3580	0.90	9/4853 (0.2%)
1	B	0.35	0/3582	0.86	11/4855 (0.2%)
All	All	0.35	0/7162	0.88	20/9708 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	231	ALA	N-CA-C	8.24	119.89	111.07
1	B	231	ALA	N-CA-C	7.34	118.93	111.07
1	A	128	ALA	N-CA-C	-6.94	96.33	108.20
1	A	127	TYR	N-CA-C	-6.51	103.45	111.33
1	B	127	TYR	N-CA-C	-6.50	104.20	111.28
1	A	353	GLN	N-CA-C	6.15	119.73	112.72
1	B	128	ALA	N-CA-C	-5.94	98.05	108.20
1	B	140	SER	N-CA-C	-5.80	105.09	111.82
1	A	140	SER	N-CA-C	-5.76	105.14	111.82
1	B	353	GLN	N-CA-C	5.75	119.27	112.72
1	A	159	ARG	N-CA-C	5.65	119.95	112.72
1	B	461	LYS	CA-C-N	5.64	125.32	119.56
1	B	461	LYS	C-N-CA	5.64	125.32	119.56
1	B	37	ASP	N-CA-C	5.33	116.90	111.14
1	B	358	LEU	N-CA-C	5.24	116.58	107.93
1	A	39	PHE	N-CA-C	5.21	118.75	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ALA	N-CA-C	5.20	116.64	111.07
1	B	39	PHE	N-CA-C	5.18	119.79	111.81
1	A	106	ASN	N-CA-C	-5.10	105.72	111.28
1	B	230	THR	CB-CA-C	-5.07	110.71	116.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	367	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3515	0	3557	58	0
1	B	3517	0	3563	53	2
2	A	357	0	0	2	0
2	B	291	0	0	7	0
All	All	7680	0	7120	110	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:ARG:HG2	1:B:12:GLU:HG2	1.45	0.94
1:A:428:ALA:HB3	1:A:435:LEU:HD23	1.64	0.79
1:B:436:LYS:HD3	1:B:447:GLU:HG3	1.65	0.79
1:B:48:ALA:HA	1:B:51:MET:HE3	1.68	0.75
1:A:440:LEU:HD21	1:A:447:GLU:N	2.03	0.73
1:A:446:THR:HG23	1:A:449:GLU:H	1.55	0.72
1:B:113:ASN:HB3	1:B:123:LEU:HD22	1.72	0.71
1:B:145:PRO:HA	1:B:148:MET:HG2	1.73	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:PRO:HA	1:A:148:MET:HG2	1.75	0.68
1:B:184:VAL:HG13	1:B:407:MET:HE1	1.79	0.63
1:A:105:MET:O	1:A:109:GLU:HG3	1.99	0.62
1:A:196:THR:OG1	1:A:199:GLN:HG3	2.00	0.61
1:A:402:LEU:HG	1:A:408:LEU:HD11	1.81	0.61
1:A:434:THR:HG23	1:A:437:GLN:HB2	1.81	0.61
1:A:122:PRO:HG2	1:A:125:SER:HB3	1.81	0.61
1:A:420:LYS:N	1:A:420:LYS:HD2	2.15	0.61
1:B:248:LYS:O	1:B:252:GLU:HG3	2.01	0.60
1:A:24:GLN:NE2	1:A:105:MET:HE2	2.17	0.58
1:B:387:LEU:HD23	1:B:387:LEU:C	2.30	0.57
1:A:387:LEU:C	1:A:387:LEU:HD23	2.29	0.57
1:A:440:LEU:HD21	1:A:447:GLU:H	1.70	0.56
1:A:7:ARG:HB3	1:A:12:GLU:HG2	1.87	0.56
1:B:260:VAL:HG11	1:B:264:ARG:HE	1.70	0.56
1:B:303:TYR:HA	1:B:397:ARG:NH2	2.20	0.56
1:B:320:ILE:N	1:B:320:ILE:HD12	2.21	0.56
1:B:16:PRO:HB2	1:B:19:LYS:HG2	1.88	0.55
1:B:446:THR:C	1:B:448:GLU:H	2.13	0.55
1:A:408:LEU:HD22	1:A:456:PRO:HG3	1.89	0.55
1:A:172:THR:O	1:A:176:GLN:HG3	2.07	0.54
1:B:172:THR:O	1:B:176:GLN:HG3	2.07	0.54
1:A:229:GLY:HA3	2:A:696:HOH:O	2.08	0.54
1:B:264:ARG:O	1:B:268:LEU:HD13	2.08	0.53
1:B:105:MET:O	1:B:109:GLU:HG3	2.09	0.53
1:A:436:LYS:HE2	1:A:447:GLU:HB2	1.90	0.53
1:B:52:LEU:C	1:B:52:LEU:HD23	2.34	0.53
1:A:407:MET:HB3	1:A:459:LEU:HD13	1.91	0.52
1:B:282:ARG:HD2	2:B:583:HOH:O	2.10	0.52
1:A:114:ARG:O	1:A:118:ILE:HG13	2.10	0.52
1:A:236:LEU:O	1:A:237:ASN:HB2	2.10	0.52
1:B:260:VAL:CG1	1:B:264:ARG:HE	2.22	0.51
1:A:428:ALA:HB2	1:A:438:ALA:HB2	1.93	0.51
1:A:362:LYS:HB2	1:A:363:PRO:HD3	1.93	0.50
1:A:410:THR:HA	1:A:413:ASN:HD22	1.77	0.50
1:A:447:GLU:OE2	1:A:448:GLU:HG3	2.11	0.50
1:B:92:LEU:HD13	1:B:102:GLN:HB3	1.94	0.50
1:A:440:LEU:HD11	1:A:447:GLU:HB3	1.93	0.50
1:B:236:LEU:O	1:B:237:ASN:HB2	2.10	0.50
1:A:52:LEU:C	1:A:52:LEU:HD23	2.36	0.49
1:A:427:LYS:HE2	1:A:441:GLU:HB2	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:TYR:HB2	1:B:162:PRO:HD3	1.94	0.49
1:B:196:THR:OG1	1:B:199:GLN:HG3	2.11	0.49
1:B:446:THR:C	1:B:448:GLU:N	2.71	0.49
1:A:126:LYS:HA	2:A:693:HOH:O	2.13	0.48
1:B:362:LYS:HB2	1:B:363:PRO:HD3	1.94	0.48
1:B:427:LYS:HD2	1:B:442:LEU:HD21	1.95	0.48
1:A:324:LYS:HE2	1:B:191:ASP:O	2.14	0.48
1:B:306:ILE:HG22	1:B:308:GLU:HG3	1.96	0.48
1:B:420:LYS:HG3	2:B:694:HOH:O	2.14	0.48
1:A:180:GLN:HE21	1:A:180:GLN:HA	1.79	0.48
1:A:188:HIS:O	1:A:189:LEU:HB2	2.14	0.47
1:B:197:LEU:CD2	1:B:393:PRO:HB3	2.45	0.47
1:B:80:GLU:HG2	1:B:85:LYS:HD2	1.95	0.47
1:B:223:TYR:CD1	1:B:257:PRO:HG2	2.49	0.47
1:B:184:VAL:HG13	1:B:407:MET:CE	2.45	0.47
1:A:434:THR:CG2	1:A:437:GLN:HB2	2.46	0.46
1:A:436:LYS:CE	1:A:447:GLU:HB2	2.46	0.46
1:A:427:LYS:HD3	1:A:438:ALA:HB1	1.97	0.46
1:B:420:LYS:O	1:B:424:ILE:HG13	2.15	0.46
1:A:180:GLN:HA	1:A:180:GLN:NE2	2.30	0.46
1:B:105:MET:HG3	2:B:723:HOH:O	2.16	0.46
1:B:24:GLN:OE1	1:B:105:MET:HE2	2.16	0.46
1:B:187:THR:O	1:B:188:HIS:HB2	2.16	0.46
1:B:114:ARG:O	1:B:118:ILE:HG13	2.16	0.45
1:A:25:THR:OG1	1:A:106:ASN:HA	2.16	0.45
1:A:223:TYR:CD1	1:A:257:PRO:HG2	2.51	0.45
1:B:399:GLU:HG3	2:B:612:HOH:O	2.18	0.44
1:A:97:THR:HG21	1:A:101:THR:HB	1.99	0.44
1:A:306:ILE:HG22	1:A:308:GLU:HG3	2.00	0.44
1:A:455:VAL:HB	1:A:458:ARG:HD2	1.99	0.44
1:A:161:TYR:HB2	1:A:162:PRO:HD3	2.00	0.43
1:A:339:ARG:NH1	1:A:343:ASN:OD1	2.51	0.43
1:B:326:ASN:HB3	2:B:591:HOH:O	2.17	0.43
1:A:28:SER:HB2	1:A:102:GLN:HG2	2.01	0.43
1:B:428:ALA:HB1	1:B:433:LYS:O	2.19	0.43
1:A:446:THR:HG22	1:A:449:GLU:OE1	2.18	0.43
1:A:43:LEU:O	1:A:47:ARG:HG2	2.19	0.42
1:B:129:HIS:HA	1:B:130:PRO:HD3	1.91	0.42
1:B:145:PRO:HB3	1:B:271:HIS:CE1	2.55	0.42
1:B:66:LEU:HD11	2:B:531:HOH:O	2.19	0.42
1:A:434:THR:HG23	1:A:437:GLN:CB	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:SER:O	1:A:126:LYS:HB2	2.20	0.42
1:A:86:TRP:CZ2	1:A:114:ARG:HD3	2.54	0.42
1:B:125:SER:O	1:B:126:LYS:HB2	2.19	0.42
1:A:397:ARG:HD2	1:A:401:TYR:CZ	2.55	0.42
1:A:427:LYS:HE3	1:A:441:GLU:OE1	2.20	0.42
1:B:4:ARG:HG2	1:B:4:ARG:HH11	1.85	0.41
1:A:442:LEU:HB3	1:A:444:TYR:CD1	2.55	0.41
1:A:232:VAL:C	1:A:268:LEU:HD12	2.45	0.41
1:B:144:PHE:N	1:B:145:PRO:HD2	2.35	0.41
1:A:420:LYS:N	1:A:420:LYS:CD	2.84	0.41
1:A:187:THR:O	1:A:188:HIS:HB2	2.21	0.41
1:B:405:ASN:HA	1:B:406:PRO:HD3	1.86	0.41
1:B:106:ASN:C	1:B:106:ASN:HD22	2.29	0.41
1:A:399:GLU:OE2	1:A:403:GLN:NE2	2.53	0.41
1:A:48:ALA:HA	1:A:51:MET:HE3	2.03	0.41
1:B:126:LYS:HA	2:B:751:HOH:O	2.21	0.41
1:B:396:GLU:H	1:B:396:GLU:CD	2.29	0.41
1:A:455:VAL:HB	1:A:458:ARG:CD	2.51	0.40
1:B:226:ALA:HB1	1:B:268:LEU:HD12	2.02	0.40
1:B:401:TYR:O	1:B:405:ASN:HB2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:ARG:NH2	1:B:339:ARG:NH2[3_655]	1.75	0.45
1:B:452:ARG:NH1	1:B:452:ARG:NH1[4_555]	1.83	0.37

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/466 (97%)	431 (95%)	23 (5%)	0	100	100
1	B	453/466 (97%)	439 (97%)	14 (3%)	0	100	100
All	All	907/932 (97%)	870 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/367 (98%)	359 (99%)	2 (1%)	78	77
1	B	361/367 (98%)	358 (99%)	3 (1%)	73	70
All	All	722/734 (98%)	717 (99%)	5 (1%)	76	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	TYR
1	A	91	PRO
1	B	49	TYR
1	B	91	PRO
1	B	259	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	131	ASN
1	A	180	GLN
1	A	345	HIS
1	A	355	ASN
1	A	437	GLN
1	B	31	ASN
1	B	326	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	329	GLN
1	B	355	ASN
1	B	386	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/466 (98%)	0.35	51 (11%) 10 8	13, 23, 76, 80	0
1	B	457/466 (98%)	0.48	38 (8%) 17 15	15, 28, 58, 72	0
All	All	915/932 (98%)	0.41	89 (9%) 13 11	13, 26, 68, 80	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	435	LEU	7.8
1	A	428	ALA	7.4
1	A	319	SER	6.9
1	A	421	ALA	5.8
1	A	318	SER	5.8
1	A	429	LEU	5.3
1	A	438	ALA	5.2
1	B	316	PRO	5.1
1	A	440	LEU	5.1
1	A	444	TYR	4.8
1	A	320	ILE	4.7
1	A	416	ILE	4.5
1	A	439	ALA	4.4
1	A	442	LEU	4.3
1	A	410	THR	4.2
1	B	122	PRO	4.2
1	A	445	LEU	4.1
1	A	413	ASN	4.1
1	B	5	ILE	4.0
1	A	425	VAL	4.0
1	A	322	PRO	3.9
1	A	424	ILE	3.7
1	A	450	PHE	3.7
1	B	10	MET	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	321	MET	3.6
1	B	66	LEU	3.6
1	A	434	THR	3.5
1	A	453	ILE	3.4
1	A	463	HIS	3.4
1	A	411	ALA	3.4
1	A	422	ALA	3.4
1	B	123	LEU	3.3
1	A	446	THR	3.2
1	A	408	LEU	3.2
1	A	412	LEU	3.2
1	A	430	LYS	3.2
1	B	13	VAL	3.2
1	A	426	LYS	3.1
1	A	433	LYS	3.1
1	B	253	GLU	3.1
1	B	315	GLU	3.1
1	B	322	PRO	3.0
1	B	418	TYR	2.9
1	A	407	MET	2.9
1	B	445	LEU	2.8
1	A	427	LYS	2.8
1	A	441	GLU	2.8
1	A	418	TYR	2.8
1	A	317	GLY	2.8
1	B	9	THR	2.8
1	A	415	ALA	2.7
1	B	128	ALA	2.6
1	B	15	VAL	2.6
1	B	432	LYS	2.6
1	B	441	GLU	2.6
1	A	454	VAL	2.6
1	A	321	MET	2.5
1	B	448	GLU	2.5
1	B	450	PHE	2.5
1	B	14	ARG	2.5
1	B	446	THR	2.5
1	A	120	GLY	2.4
1	A	420	LYS	2.4
1	B	119	LEU	2.4
1	B	429	LEU	2.4
1	B	440	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	320	ILE	2.4
1	A	14	ARG	2.4
1	B	121	LYS	2.4
1	A	451	ASP	2.3
1	B	413	ASN	2.3
1	A	414	LYS	2.3
1	A	303	TYR	2.3
1	B	442	LEU	2.3
1	A	447	GLU	2.2
1	A	18	ASP	2.2
1	B	176	GLN	2.2
1	A	443	GLY	2.2
1	B	439	ALA	2.2
1	B	69	GLU	2.2
1	B	120	GLY	2.2
1	A	409	ALA	2.1
1	B	33	ARG	2.1
1	B	438	ALA	2.1
1	B	451	ASP	2.1
1	A	121	LYS	2.1
1	B	20	TYR	2.0
1	B	425	VAL	2.0
1	A	403	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.