



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:26 PM UTC

PDB ID : 2VDR / pdb_00002vdr
Title : Integrin AlphaIIbBeta3 Headpiece Bound to a chimeric Fibrinogen Gamma chain peptide, LGGAKQRGDV
Authors : Springer, T.A.; Zhu, J.; Xiao, T.
Deposited on : 2007-10-10
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	4% 96% .
2	B	461	24% 93% 6% .
3	C	10	40% 70% 30%
4	H	221	7% 94% 5% .
5	L	214	2% 93% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	D	5	
7	E	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MAN	D	3	X	-	-	-
7	MAN	E	3	X	-	-	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 11884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTEGRIN ALPHA-IIB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	9	0
			3539	2250	616	665	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	282	GLY	ALA	conflict	UNP P08514

- Molecule 2 is a protein called INTEGRIN BETA-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	455	Total	C	N	O	S	0	10	0
			3580	2236	607	704	33			

- Molecule 3 is a protein called FIBRINOGEN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	0	0	0
			54	31	12	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	408	ARG	ALA	engineered mutation	UNP Q53Y18


- Molecule 4 is a protein called MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	219	Total	C	N	O	S	0	3	0
			1678	1064	270	338	6			

- Molecule 5 is a protein called MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	L	214	Total	C	N	O	S	0	9	0
			1686	1051	273	351	11			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	E	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	4	Total	Ca	0	0
			4	4		
9	B	2	Total	Ca	0	0
			2	2		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	1	Total	Mg	0	0
			1	1		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	412	Total	O	0	0
			412	412		
12	B	256	Total	O	0	0
			256	256		
12	C	12	Total	O	0	0
			12	12		
12	H	214	Total	O	0	0
			214	214		
12	L	248	Total	O	0	0
			248	248		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: INTEGRIN ALPHA-IIB



• Molecule 2: INTEGRIN BETA-3




• Molecule 3: FIBRINOGEN



• Molecule 4: MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN





• Molecule 5: MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN



• Molecule 6: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 7: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	148.33Å 148.33Å 176.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.44 – 2.40 43.44 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.5 (43.44-2.40) 96.1 (43.44-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.148 , 0.193 0.154 , 0.198	Depositor DCC
R_{free} test set	4276 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	11884	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, GOL, NAG, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/3656	0.66	0/4980
2	B	0.45	0/3661	0.68	0/4960
3	C	0.49	0/53	0.71	0/67
4	H	0.44	0/1730	0.64	0/2369
5	L	0.44	0/1754	0.61	0/2377
All	All	0.45	0/10854	0.66	0/14753

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3539	0	3385	13	0
2	B	3580	0	3523	14	0
3	C	54	0	54	0	0
4	H	1678	0	1637	14	0
5	L	1686	0	1614	15	0
6	D	61	0	52	0	0
7	E	83	0	70	0	0
8	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	6	0	7	0	0
9	A	4	0	0	0	0
9	B	2	0	0	0	0
10	A	28	0	26	0	0
10	B	14	0	13	0	0
11	B	1	0	0	0	0
12	A	412	0	0	1	0
12	B	256	0	0	4	0
12	C	12	0	0	0	0
12	H	214	0	0	2	0
12	L	248	0	0	0	0
All	All	11884	0	10389	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:HD11	1.46	1.15
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:CD1	2.07	1.01
4:H:177[B]:GLN:CD	5:L:160[B]:LEU:CD1	2.68	0.66
4:H:177[B]:GLN:CD	5:L:160[B]:LEU:HD11	2.21	0.65
1:A:270:LEU:HD23	1:A:276[A]:ARG:HA	1.81	0.63
1:A:270:LEU:HD23	1:A:276[B]:ARG:HA	1.82	0.60
2:B:201[A]:THR:HG22	12:B:4097:HOH:O	2.02	0.59
5:L:4:MET:HE3	5:L:23:CYS:SG	2.42	0.59
5:L:48:ILE:HD13	5:L:54:LEU:HD12	1.85	0.58
5:L:136:LEU:HD21	5:L:146:VAL:HG22	1.84	0.58
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:HD12	1.99	0.57
4:H:177[A]:GLN:NE2	12:H:4197:HOH:O	2.32	0.54
1:A:280:LEU:CD1	1:A:306:LEU:HD23	2.41	0.51
4:H:30:LYS:NZ	12:H:4046:HOH:O	2.41	0.51
5:L:122:SER:O	5:L:126:THR:HG23	2.10	0.51
5:L:48:ILE:CD1	5:L:54:LEU:HD12	2.41	0.50
2:B:173:LEU:HD12	12:B:4068:HOH:O	2.11	0.50
2:B:88[B]:ILE:HD11	2:B:427:VAL:HG22	1.93	0.50
2:B:142:MET:HE1	2:B:345:VAL:HG23	1.94	0.50
5:L:136:LEU:HD12	5:L:136:LEU:N	2.27	0.48
4:H:133:VAL:HG12	4:H:134:CYS:N	2.29	0.48
1:A:280:LEU:CD1	1:A:306:LEU:CD2	2.93	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:141:GLN:NE2	12:B:4053:HOH:O	2.46	0.46
2:B:308:PHE:CE2	2:B:328:THR:HG21	2.50	0.46
2:B:201[A]:THR:HG23	12:B:4095:HOH:O	2.15	0.46
5:L:37:GLN:HB2	5:L:47:LEU:HD11	1.98	0.46
1:A:122:ALA:O	1:A:123:GLU:HB2	2.14	0.46
1:A:278[A]:HIS:CD2	1:A:339:ALA:HB1	2.51	0.46
1:A:314:MET:HE1	2:B:324:LEU:HB3	1.98	0.46
5:L:105[A]:GLU:HG3	5:L:173:TYR:OH	2.17	0.45
1:A:235:TRP:CZ2	1:A:270:LEU:HD11	2.52	0.45
2:B:223:PHE:CZ	2:B:254:THR:HG21	2.52	0.44
1:A:90[A]:ARG:CZ	12:A:4053:HOH:O	2.64	0.44
4:H:194:TRP:CG	4:H:195:PRO:HA	2.52	0.44
1:A:5:PRO:HA	1:A:8:LEU:HD21	2.00	0.44
4:H:208:SER:O	4:H:210:THR:HG23	2.18	0.43
2:B:443:PRO:HA	2:B:454:THR:HG22	2.01	0.43
1:A:278[A]:HIS:NE2	1:A:339:ALA:HB1	2.34	0.43
2:B:245:LEU:HD23	2:B:305:ASN:HB2	2.00	0.43
4:H:172:PHE:CD1	5:L:164:THR:HG23	2.54	0.42
1:A:297:ASP:O	1:A:372:ASN:HB2	2.20	0.42
1:A:262:TRP:HB3	2:B:317:LEU:HD13	2.01	0.42
4:H:158:LEU:C	4:H:158:LEU:HD23	2.45	0.42
2:B:15:GLN:O	2:B:19:VAL:HG23	2.20	0.41
4:H:177[B]:GLN:HB3	5:L:160[B]:LEU:HD11	2.02	0.41
4:H:35:HIS:O	4:H:96:CYS:HA	2.21	0.41
2:B:176:PRO:HB2	2:B:214:ARG:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	459/452 (102%)	450 (98%)	8 (2%)	1 (0%)	43 58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	460/461 (100%)	440 (96%)	17 (4%)	3 (1%)	18	28
3	C	5/10 (50%)	4 (80%)	1 (20%)	0	100	100
4	H	218/221 (99%)	213 (98%)	5 (2%)	0	100	100
5	L	223/214 (104%)	218 (98%)	5 (2%)	0	100	100
All	All	1365/1358 (100%)	1325 (97%)	36 (3%)	4 (0%)	36	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	GLU
2	B	9	GLY
2	B	10	VAL
2	B	80	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/360 (102%)	366 (99%)	3 (1%)	73	86
2	B	414/409 (101%)	412 (100%)	2 (0%)	81	91
3	C	5/6 (83%)	5 (100%)	0	100	100
4	H	192/190 (101%)	192 (100%)	0	100	100
5	L	199/188 (106%)	197 (99%)	2 (1%)	68	84
All	All	1179/1153 (102%)	1172 (99%)	7 (1%)	81	89

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LEU
1	A	166	TYR
1	A	288	TYR
2	B	215	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	277	SER
5	L	105[A]	GLU
5	L	105[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	GLN
1	A	219	GLN
1	A	388	GLN
2	B	133	ASN
2	B	141	GLN
2	B	199	GLN
2	B	301	GLN
2	B	316	ASN
2	B	449	ASN
2	B	450	ASN
3	C	407	GLN
4	H	65	GLN
5	L	27	GLN
5	L	190	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	D	1	2,6	14,14,15	0.57	0	17,19,21	0.90	1 (5%)
6	NAG	D	2	6	14,14,15	0.68	0	17,19,21	0.80	0
6	MAN	D	3	6	11,11,12	0.52	0	15,15,17	0.58	0
6	MAN	D	4	6	11,11,12	0.53	0	15,15,17	0.60	0
6	MAN	D	5	6	11,11,12	0.61	0	15,15,17	1.07	2 (13%)
7	NAG	E	1	2,7	14,14,15	0.71	0	17,19,21	1.02	1 (5%)
7	NAG	E	2	7	14,14,15	0.56	0	17,19,21	0.82	0
7	MAN	E	3	7	11,11,12	0.64	0	15,15,17	0.65	0
7	MAN	E	4	7	11,11,12	0.48	0	15,15,17	1.16	1 (6%)
7	MAN	E	5	7	11,11,12	0.51	0	15,15,17	0.93	1 (6%)
7	MAN	E	6	7	11,11,12	0.57	0	15,15,17	0.72	1 (6%)
7	MAN	E	7	7	11,11,12	0.57	0	15,15,17	0.76	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	D	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	D	2	6	-	0/6/23/26	0/1/1/1
6	MAN	D	3	6	1/1/4/5	0/2/19/22	0/1/1/1
6	MAN	D	4	6	-	2/2/19/22	0/1/1/1
6	MAN	D	5	6	-	2/2/19/22	0/1/1/1
7	NAG	E	1	2,7	-	0/6/23/26	0/1/1/1
7	NAG	E	2	7	-	2/6/23/26	0/1/1/1
7	MAN	E	3	7	1/1/4/5	0/2/19/22	0/1/1/1
7	MAN	E	4	7	-	0/2/19/22	0/1/1/1
7	MAN	E	5	7	-	0/2/19/22	0/1/1/1
7	MAN	E	6	7	-	2/2/19/22	0/1/1/1
7	MAN	E	7	7	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	4	MAN	C1-O5-C5	3.64	117.06	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	5	MAN	C1-C2-C3	2.80	113.72	109.64
7	E	5	MAN	C1-O5-C5	2.59	115.66	112.19
7	E	7	MAN	C1-O5-C5	2.18	115.11	112.19
7	E	6	MAN	C1-O5-C5	2.09	114.99	112.19
6	D	1	NAG	O5-C1-C2	-2.09	108.06	111.29
6	D	5	MAN	C1-O5-C5	2.04	114.92	112.19
7	E	1	NAG	O5-C1-C2	-2.03	108.16	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	D	3	MAN	C1
7	E	3	MAN	C1


All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	D	5	MAN	O5-C5-C6-O6
7	E	6	MAN	O5-C5-C6-O6
7	E	6	MAN	C4-C5-C6-O6
7	E	2	NAG	O5-C5-C6-O6
6	D	5	MAN	C4-C5-C6-O6
6	D	4	MAN	C4-C5-C6-O6
7	E	2	NAG	C4-C5-C6-O6
6	D	4	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	NAG	A	3015	1	14,14,15	0.47	0	17,19,21	0.74	0
10	NAG	A	3249	1	14,14,15	0.48	0	17,19,21	0.97	1 (5%)
10	NAG	B	3099	2	14,14,15	0.49	0	17,19,21	0.87	1 (5%)
8	GOL	B	1462	9	5,5,5	0.41	0	5,5,5	0.35	0
8	GOL	A	1453	-	5,5,5	0.44	0	5,5,5	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	A	3015	1	-	0/6/23/26	0/1/1/1
10	NAG	A	3249	1	-	0/6/23/26	0/1/1/1
10	NAG	B	3099	2	-	2/6/23/26	0/1/1/1
8	GOL	B	1462	9	-	2/4/4/4	-
8	GOL	A	1453	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	3249	NAG	O5-C1-C2	-2.37	107.63	111.29
10	B	3099	NAG	O5-C1-C2	-2.15	107.96	111.29

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	3099	NAG	C4-C5-C6-O6
10	B	3099	NAG	O5-C5-C6-O6
8	B	1462	GOL	O1-C1-C2-C3
8	B	1462	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	452/452 (100%)	0.66	18 (3%) 42 38	23, 48, 66, 90	9 (1%)
2	B	455/461 (98%)	1.35	111 (24%) 2 1	23, 55, 95, 109	10 (2%)
3	C	7/10 (70%)	2.41	4 (57%) 0 0	39, 49, 108, 110	0
4	H	219/221 (99%)	0.67	16 (7%) 21 17	31, 48, 76, 106	3 (1%)
5	L	214/214 (100%)	0.49	4 (1%) 66 62	22, 47, 61, 99	9 (4%)
All	All	1347/1358 (99%)	0.88	153 (11%) 10 7	22, 50, 87, 110	31 (2%)

All (153) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	457	CYS	7.7
2	B	456	GLU	7.5
2	B	441	ALA	6.6
2	B	72	LYS	5.8
2	B	458	GLY	5.4
2	B	461	ARG	5.4
2	B	444	ASN	5.2
2	B	442	GLU	4.8
2	B	1	GLY	4.8
2	B	455	PHE	4.4
3	C	406	LYS	4.3
2	B	379	VAL	4.3
2	B	9	GLY	4.2
4	H	138	THR	4.0
2	B	10	VAL	4.0
3	C	405	ALA	4.0
2	B	395	VAL	3.9
2	B	405	GLY	3.9
4	H	137	THR	3.8
2	B	453	GLY	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	450	ASN	3.7
2	B	447	ARG	3.7
1	A	1	LEU	3.7
2	B	451	GLY	3.4
2	B	11	SER	3.4
2	B	24	ALA	3.4
2	B	460	CYS	3.4
2	B	437	CYS	3.4
4	H	134	CYS	3.4
2	B	397	PHE	3.3
2	B	452	ASN	3.3
4	H	221	PRO	3.3
2	B	422	LYS	3.3
2	B	51	PRO	3.3
2	B	436	ALA	3.3
2	B	443	PRO	3.3
2	B	49	CYS	3.2
2	B	446	HIS	3.1
2	B	91	ARG	3.1
2	B	392	GLY	3.1
2	B	22	MET	3.0
2	B	393	ASP	3.0
2	B	40	LEU	3.0
3	C	407	GLN	3.0
2	B	123	SER	3.0
2	B	34	GLY	3.0
2	B	459	VAL	3.0
4	H	195	PRO	3.0
2	B	372	ALA	2.9
2	B	32	PRO	2.9
2	B	366	LEU	2.9
2	B	374	CYS	2.9
2	B	454	THR	2.9
5	L	121	SER	2.9
2	B	380	ILE	2.9
2	B	435	CYS	2.9
4	H	133	VAL	2.9
2	B	33	LEU	2.9
2	B	56	PHE	2.9
2	B	61	ALA	2.9
2	B	433	CYS	2.8
2	B	375	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	35	SER	2.8
2	B	83	VAL	2.8
4	H	194	TRP	2.8
2	B	430	THR	2.8
2	B	31	LEU	2.8
1	A	8	LEU	2.8
2	B	59	SER	2.7
1	A	328	VAL	2.7
2	B	79	GLN	2.7
2	B	69	LEU	2.7
2	B	387[A]	MET	2.7
1	A	326	GLY	2.7
2	B	403	VAL	2.7
2	B	382	GLY	2.6
1	A	337	PRO	2.6
2	B	52	GLU	2.6
2	B	373	THR	2.6
2	B	8	ARG	2.6
2	B	53	SER	2.6
2	B	396	SER	2.6
2	B	445	SER	2.6
1	A	325	VAL	2.6
2	B	36	PRO	2.6
2	B	92	LEU	2.6
2	B	58	VAL	2.6
5	L	213	GLU	2.5
2	B	121	SER	2.5
4	H	177[A]	GLN	2.5
1	A	248	LEU	2.5
2	B	253	LYS	2.5
2	B	449	ASN	2.5
1	A	323	ALA	2.5
2	B	23	CYS	2.5
1	A	335[A]	ARG	2.5
4	H	193	THR	2.5
2	B	80	VAL	2.4
2	B	398	SER	2.4
5	L	212	ASN	2.4
2	B	391	ILE	2.4
2	B	67	ARG	2.4
3	C	411	VAL	2.4
1	A	324	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	67	SER	2.3
4	H	196	SER	2.3
2	B	394	THR	2.3
2	B	448	CYS	2.3
2	B	419	VAL	2.3
2	B	409	GLU	2.3
2	B	30	ALA	2.3
4	H	186	SER	2.3
4	H	220	GLY	2.3
2	B	46	LYS	2.3
2	B	65	GLU	2.3
2	B	2	PRO	2.3
2	B	390	LYS	2.3
2	B	406	CYS	2.3
2	B	82	GLN	2.2
4	H	144	LEU	2.2
1	A	162[A]	TRP	2.2
2	B	81	THR	2.2
1	A	130	CYS	2.2
2	B	319[A]	GLN	2.2
2	B	124	MET	2.2
2	B	45	LEU	2.2
2	B	84	SER	2.2
2	B	28	ASP	2.2
2	B	39	ASP	2.2
2	B	241	ASP	2.2
2	B	48	ASN	2.2
2	B	440	GLN	2.2
2	B	5	CYS	2.2
1	A	400	ARG	2.2
4	H	175	VAL	2.1
2	B	13	CYS	2.1
2	B	389	LEU	2.1
2	B	431	PHE	2.1
2	B	44	LEU	2.1
2	B	215	ASN	2.1
1	A	327	ARG	2.1
1	A	336	GLY	2.1
2	B	412[A]	LYS	2.1
2	B	162	SER	2.1
4	H	65	GLN	2.0
2	B	54	ILE	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	219	PRO	2.0
5	L	91	TYR	2.0
1	A	338	HIS	2.0
2	B	62	ARG	2.0
1	A	50	THR	2.0
2	B	88[A]	ILE	2.0
4	H	208	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	D	1	14/15	-	-	44,54,61,69	0
6	NAG	D	2	14/15	-	-	74,80,88,96	0
6	MAN	D	3	11/12	-	-	103,107,112,115	0
6	MAN	D	4	11/12	-	-	101,105,108,108	0
6	MAN	D	5	11/12	-	-	115,116,117,117	0
7	NAG	E	1	14/15	-	-	66,74,78,80	0
7	NAG	E	2	14/15	-	-	68,82,90,91	0
7	MAN	E	3	11/12	-	-	80,91,93,96	0
7	MAN	E	4	11/12	-	-	67,73,80,86	0
7	MAN	E	5	11/12	-	-	65,67,71,73	0
7	MAN	E	6	11/12	-	-	90,93,95,95	0
7	MAN	E	7	11/12	-	-	95,99,100,100	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	NAG	A	3249	14/15	0.42	0.25	91,93,94,94	14
10	NAG	A	3015	14/15	0.51	0.28	89,91,93,94	14
10	NAG	B	3099	14/15	0.65	0.21	103,108,110,111	0
8	GOL	B	1462	6/6	0.87	0.20	62,62,63,65	0
8	GOL	A	1453	6/6	0.87	0.21	60,68,72,75	0
9	CA	B	2002	1/1	0.98	0.17	40,40,40,40	0
11	MG	B	2001	1/1	0.98	0.15	28,28,28,28	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	CA	A	2005	1/1	0.99	0.17	51,51,51,51	0
9	CA	A	2006	1/1	0.99	0.13	61,61,61,61	0
9	CA	A	2007	1/1	0.99	0.15	57,57,57,57	0
9	CA	A	2004	1/1	0.99	0.14	55,55,55,55	0
9	CA	B	2003	1/1	1.00	0.19	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.