



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:50 PM UTC

PDB ID : 3VDI / pdb_00003vdi
Title : Structure of the FMO protein from Pelodictyon phaeum
Authors : Tronrud, D.E.; Larson, C.R.; Seng, C.O.; Lauman, L.; Matthies, H.J.; Wen, J.; Blankenship, R.E.; Allen, J.P.
Deposited on : 2012-01-05
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called bacteriochlorophyll A protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2782	1771	486	518	7	0	2	0

- Molecule 2 is BACTERIOCHLOROPHYLL A (CCD ID: BCL) (formula: C₅₅H₇₄MgN₄O₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Mg	N	O		
2	A	1	Total 66	55	1	4	6	0	0
2	A	1	Total 66	55	1	4	6	0	0
2	A	1	Total 66	55	1	4	6	0	0
2	A	1	Total 66	55	1	4	6	0	0
2	A	1	Total 66	55	1	4	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
2	A	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			8	5	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	203	Total	O	0	2
			203	203		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: bacteriochlorophyll A protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	83.97Å 83.97Å 115.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 1.99 40.00 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-1.99) 94.5 (40.00-1.99)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.162 , 0.190 0.165 , 0.196	Depositor DCC
R_{free} test set	1523 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.39$	Xtriage
Estimated twinning fraction	0.056 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3485	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/2857 (0.0%)	1.11	3/3874 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	289	ILE	CA-CB	5.17	1.56	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	ILE	N-CA-C	-8.05	98.26	109.21
1	A	35	ALA	N-CA-C	-5.73	102.82	110.55
1	A	301	ASN	CA-CB-CG	5.40	118.00	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2782	0	2696	23	0
2	A	462	0	518	15	0
3	A	38	0	47	0	0
4	A	203	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3485	0	3261	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:HIS:HE1	1:A:354:GLY:H	1.25	0.81
1:A:146:LYS:HG2	1:A:216[A]:VAL:HG12	1.71	0.71
2:A:402:BCL:HBB2	2:A:407:BCL:H193	1.73	0.69
1:A:77:LEU:HD22	2:A:407:BCL:H191	1.80	0.63
2:A:403:BCL:HBB3	2:A:403:BCL:HHC	1.83	0.61
1:A:158[A]:TRP:CE3	2:A:401:BCL:HBB2	2.43	0.53
1:A:347:HIS:CE1	1:A:354:GLY:H	2.16	0.53
1:A:307:THR:HG21	1:A:345:VAL:HG21	1.91	0.52
1:A:40:LEU:O	4:A:1133:HOH:O	2.19	0.51
2:A:407:BCL:CBB	2:A:407:BCL:HMB1	2.41	0.50
2:A:404:BCL:H2C	2:A:404:BCL:H92	1.93	0.49
1:A:63:THR:HG22	1:A:80:GLU:HG3	1.94	0.49
2:A:401:BCL:HBB2	2:A:401:BCL:HMB3	1.96	0.48
2:A:406:BCL:HMB1	2:A:406:BCL:CBB	2.43	0.48
1:A:329:GLN:NE2	4:A:1192:HOH:O	2.37	0.48
1:A:158[A]:TRP:CZ2	1:A:205:GLY:O	2.67	0.48
1:A:101:VAL:HB	2:A:401:BCL:HED2	1.95	0.47
1:A:336:TRP:HB3	1:A:341:TYR:CG	2.49	0.47
1:A:81:VAL:HG11	2:A:406:BCL:O1A	2.15	0.47
1:A:231:SER:HA	1:A:238:LEU:CD2	2.45	0.47
1:A:181:PHE:HE1	2:A:403:BCL:H142	1.81	0.46
1:A:147:VAL:HB	2:A:401:BCL:HBB3	1.97	0.46
2:A:402:BCL:H18	2:A:406:BCL:HMB2	1.97	0.46
1:A:22:TRP:HA	1:A:270:PRO:HA	1.96	0.46
1:A:11:TYR:CE1	2:A:404:BCL:H43	2.52	0.45
2:A:407:BCL:HMB1	2:A:407:BCL:HBB2	1.98	0.45
1:A:77:LEU:HD13	1:A:99:VAL:HG22	1.98	0.45
1:A:94:VAL:HG22	1:A:111:GLU:HG2	1.99	0.44
1:A:240:PRO:HG2	1:A:270:PRO:HG2	2.01	0.42
1:A:347:HIS:CE1	1:A:352:GLY:HA3	2.56	0.41
1:A:24:GLN:HE21	1:A:26:LYS:HE3	1.84	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/367 (96%)	349 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/304 (96%)	289 (99%)	3 (1%)	68	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	LEU
1	A	81	VAL
1	A	264	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	308	GLN
1	A	347	HIS
1	A	355	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BCL	A	406	-	69,74,74	1.19	5 (7%)	79,115,115	1.41	8 (10%)
2	BCL	A	405	1	69,74,74	1.28	5 (7%)	79,115,115	1.35	8 (10%)
2	BCL	A	401	-	69,74,74	1.30	7 (10%)	79,115,115	1.47	7 (8%)
2	BCL	A	404	-	69,74,74	1.20	7 (10%)	79,115,115	1.30	7 (8%)
3	PG4	A	409	-	9,9,12	0.63	0	8,8,11	0.41	0
3	PG4	A	410	-	6,6,12	0.47	0	5,5,11	0.52	0
2	BCL	A	403	-	69,74,74	1.17	6 (8%)	79,115,115	1.33	8 (10%)
3	PG4	A	411	-	7,7,12	0.80	0	6,6,11	0.60	0
2	BCL	A	402	4	69,74,74	1.32	7 (10%)	79,115,115	1.37	8 (10%)
2	BCL	A	407	-	69,74,74	1.22	5 (7%)	79,115,115	1.47	13 (16%)
3	PG4	A	408	-	12,12,12	0.69	0	11,11,11	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BCL	A	406	-	-	5/41/137/137	-
2	BCL	A	405	1	-	3/41/137/137	-
2	BCL	A	401	-	-	2/41/137/137	-
2	BCL	A	404	-	-	1/41/137/137	-
3	PG4	A	409	-	-	1/7/7/10	-
3	PG4	A	410	-	-	3/4/4/10	-
2	BCL	A	403	-	-	5/41/137/137	-
3	PG4	A	411	-	-	3/5/5/10	-
2	BCL	A	402	4	-	1/41/137/137	-
2	BCL	A	407	-	-	3/41/137/137	-
3	PG4	A	408	-	-	3/10/10/10	-

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	BCL	MG-NA	5.44	2.19	2.06
2	A	402	BCL	MG-NA	5.31	2.18	2.06
2	A	407	BCL	MG-NA	5.30	2.18	2.06
2	A	403	BCL	MG-NA	5.01	2.18	2.06
2	A	406	BCL	MG-NA	4.66	2.17	2.06
2	A	404	BCL	MG-NA	4.66	2.17	2.06
2	A	405	BCL	MG-NC	4.41	2.16	2.06
2	A	402	BCL	MG-NC	4.40	2.16	2.06
2	A	405	BCL	MG-NA	4.38	2.16	2.06
2	A	401	BCL	MG-NC	3.41	2.14	2.06
2	A	406	BCL	MG-NC	3.35	2.14	2.06
2	A	407	BCL	MG-NC	3.33	2.14	2.06
2	A	404	BCL	C3B-C4B	3.12	1.47	1.41
2	A	406	BCL	C3B-C4B	3.11	1.47	1.41
2	A	404	BCL	MG-NC	2.94	2.13	2.06
2	A	402	BCL	OBD-CAD	2.81	1.27	1.22
2	A	405	BCL	C1-C2	2.79	1.57	1.49
2	A	401	BCL	C1D-ND	2.75	1.41	1.37
2	A	402	BCL	CHD-C1D	2.71	1.43	1.38
2	A	407	BCL	C3B-C4B	2.63	1.46	1.41
2	A	405	BCL	CHD-C1D	2.61	1.43	1.38
2	A	406	BCL	C1D-ND	2.60	1.41	1.37
2	A	407	BCL	C1D-ND	2.56	1.41	1.37
2	A	401	BCL	C3B-C4B	2.50	1.46	1.41
2	A	405	BCL	C1D-ND	2.49	1.41	1.37
2	A	403	BCL	CHD-C1D	2.49	1.43	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	404	BCL	CHD-C1D	2.43	1.43	1.38
2	A	403	BCL	MG-NC	2.40	2.12	2.06
2	A	402	BCL	C3B-C4B	2.38	1.45	1.41
2	A	401	BCL	CHD-C1D	2.37	1.43	1.38
2	A	403	BCL	C3D-C4D	-2.32	1.39	1.44
2	A	407	BCL	C3D-C4D	-2.29	1.39	1.44
2	A	402	BCL	C1D-ND	2.21	1.40	1.37
2	A	404	BCL	C1D-ND	2.20	1.40	1.37
2	A	406	BCL	CHD-C1D	2.16	1.42	1.38
2	A	403	BCL	C1D-ND	2.10	1.40	1.37
2	A	403	BCL	C3B-C4B	2.10	1.45	1.41
2	A	404	BCL	C1-C2	2.06	1.55	1.49
2	A	402	BCL	C3D-C4D	-2.06	1.39	1.44
2	A	404	BCL	C3D-C4D	-2.03	1.39	1.44
2	A	401	BCL	CAA-CBA	2.01	1.58	1.52
2	A	401	BCL	C3D-C4D	-2.00	1.39	1.44

All (59) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	405	BCL	C4D-CHA-C1A	5.55	127.86	121.24
2	A	401	BCL	C4D-CHA-C1A	5.52	127.83	121.24
2	A	406	BCL	C4D-CHA-C1A	5.47	127.76	121.24
2	A	407	BCL	C4D-CHA-C1A	5.13	127.36	121.24
2	A	402	BCL	C4D-CHA-C1A	5.00	127.21	121.24
2	A	404	BCL	C4D-CHA-C1A	4.82	126.99	121.24
2	A	401	BCL	C4A-NA-C1A	4.25	108.62	106.68
2	A	403	BCL	C4D-CHA-C1A	4.24	126.31	121.24
2	A	402	BCL	C1D-ND-C4D	-4.18	103.38	106.31
2	A	401	BCL	C1-C2-C3	-4.03	119.59	126.20
2	A	407	BCL	CHD-C1D-ND	-3.93	119.27	124.80
2	A	406	BCL	C4A-NA-C1A	3.88	108.45	106.68
2	A	403	BCL	C1D-ND-C4D	-3.87	103.60	106.31
2	A	402	BCL	CHD-C1D-ND	-3.86	119.37	124.80
2	A	403	BCL	C4A-NA-C1A	3.85	108.44	106.68
2	A	405	BCL	C1D-ND-C4D	-3.84	103.62	106.31
2	A	405	BCL	CHD-C1D-ND	-3.83	119.41	124.80
2	A	407	BCL	C1D-ND-C4D	-3.81	103.64	106.31
2	A	406	BCL	CHD-C1D-ND	-3.80	119.46	124.80
2	A	406	BCL	C1D-ND-C4D	-3.80	103.65	106.31
2	A	404	BCL	C1D-ND-C4D	-3.78	103.66	106.31
2	A	401	BCL	C1D-ND-C4D	-3.74	103.69	106.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	404	BCL	CHD-C1D-ND	-3.66	119.65	124.80
2	A	403	BCL	CHD-C1D-ND	-3.64	119.67	124.80
2	A	401	BCL	CHD-C1D-ND	-3.55	119.81	124.80
2	A	401	BCL	CHA-C1A-NA	-3.36	118.78	126.39
2	A	402	BCL	C4A-NA-C1A	3.35	108.21	106.68
2	A	407	BCL	O2D-CGD-CBD	3.20	116.83	111.23
2	A	403	BCL	C1-C2-C3	-3.16	121.02	126.20
2	A	405	BCL	C4A-NA-C1A	3.13	108.11	106.68
2	A	406	BCL	CHA-C1A-NA	-3.09	119.39	126.39
2	A	406	BCL	O2D-CGD-CBD	3.07	116.60	111.23
2	A	401	BCL	C2A-C1A-CHA	3.06	129.17	123.87
2	A	402	BCL	CHA-C1A-NA	-3.04	119.52	126.39
2	A	407	BCL	C16-C15-C13	3.03	126.05	115.97
2	A	402	BCL	C1C-NC-C4C	3.00	108.05	106.68
2	A	407	BCL	C4A-NA-C1A	2.88	107.99	106.68
2	A	404	BCL	C4A-NA-C1A	2.83	107.97	106.68
2	A	405	BCL	CHA-C1A-NA	-2.82	120.00	126.39
2	A	403	BCL	CHA-C1A-NA	-2.81	120.03	126.39
2	A	403	BCL	C2A-C1A-CHA	2.80	128.73	123.87
2	A	404	BCL	CHA-C1A-NA	-2.78	120.10	126.39
2	A	407	BCL	CHA-C1A-NA	-2.78	120.10	126.39
2	A	405	BCL	C2A-C1A-CHA	2.76	128.66	123.87
2	A	404	BCL	C2A-C1A-CHA	2.64	128.44	123.87
2	A	402	BCL	C2A-C1A-CHA	2.62	128.41	123.87
2	A	405	BCL	CED-O2D-CGD	2.45	121.48	115.92
2	A	407	BCL	C1C-NC-C4C	2.39	107.77	106.68
2	A	406	BCL	C2A-C1A-CHA	2.38	127.99	123.87
2	A	407	BCL	C2A-C1A-CHA	2.36	127.95	123.87
2	A	407	BCL	C1-C2-C3	-2.35	122.34	126.20
2	A	403	BCL	CED-O2D-CGD	2.34	121.23	115.92
2	A	407	BCL	CED-O2D-CGD	2.33	121.21	115.92
2	A	405	BCL	C1-C2-C3	-2.26	122.50	126.20
2	A	407	BCL	C17-C16-C15	-2.22	103.31	113.28
2	A	407	BCL	O2D-CGD-O1D	-2.15	119.65	123.85
2	A	406	BCL	C1C-NC-C4C	2.13	107.65	106.68
2	A	404	BCL	C1C-NC-C4C	2.07	107.62	106.68
2	A	402	BCL	C1-C2-C3	-2.03	122.88	126.20

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	403	BCL	C4C-C3C-CAC-CBC
2	A	407	BCL	CHA-CBD-CGD-O2D
3	A	411	PG4	O2-C3-C4-O3
3	A	409	PG4	O4-C7-C8-O5
3	A	410	PG4	O2-C3-C4-O3
3	A	408	PG4	O2-C3-C4-O3
2	A	403	BCL	C16-C17-C18-C19
3	A	411	PG4	O3-C5-C6-O4
3	A	410	PG4	O3-C5-C6-O4
2	A	406	BCL	C10-C11-C12-C13
2	A	403	BCL	C2C-C3C-CAC-CBC
2	A	403	BCL	C16-C17-C18-C20
2	A	402	BCL	C13-C15-C16-C17
2	A	406	BCL	C4-C3-C5-C6
2	A	406	BCL	C2-C3-C5-C6
2	A	406	BCL	C4C-C3C-CAC-CBC
3	A	410	PG4	C6-C5-O3-C4
3	A	408	PG4	C3-C4-O3-C5
2	A	407	BCL	CHA-CBD-CGD-O1D
2	A	407	BCL	CAA-CBA-CGA-O2A
2	A	405	BCL	CAA-CBA-CGA-O2A
2	A	406	BCL	C2-C1-O2A-CGA
2	A	401	BCL	CAA-CBA-CGA-O2A
2	A	405	BCL	C2-C3-C5-C6
2	A	405	BCL	C4-C3-C5-C6
2	A	404	BCL	C2A-CAA-CBA-CGA
3	A	408	PG4	O1-C1-C2-O2
3	A	411	PG4	C6-C5-O3-C4
2	A	401	BCL	CAA-CBA-CGA-O1A
2	A	403	BCL	CAA-CBA-CGA-O2A


There are no ring outliers.




6 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	406	BCL	3	0
2	A	401	BCL	4	0
2	A	404	BCL	2	0
2	A	403	BCL	2	0
2	A	402	BCL	2	0
2	A	407	BCL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/367 (97%)	-0.08	8 (2%) 62 61	11, 21, 37, 66	2 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	208	SER	6.0
1	A	362	GLN	3.5
1	A	53	LEU	2.6
1	A	56	GLN	2.6
1	A	57	LYS	2.4
1	A	55	ALA	2.2
1	A	212	GLY	2.2
1	A	266	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PG4	A	411	8/13	0.76	0.20	49,49,50,51	0
3	PG4	A	410	7/13	0.88	0.14	43,43,44,44	0
3	PG4	A	408	13/13	0.90	0.10	23,24,27,27	13
3	PG4	A	409	10/13	0.91	0.12	39,39,41,42	0
2	BCL	A	406	66/66	0.95	0.07	16,19,28,28	0
2	BCL	A	402	66/66	0.96	0.07	12,15,32,32	0
2	BCL	A	405	66/66	0.96	0.06	12,17,24,28	0
2	BCL	A	401	66/66	0.96	0.06	13,18,20,22	0
2	BCL	A	407	66/66	0.96	0.06	12,15,27,34	0
2	BCL	A	404	66/66	0.97	0.05	12,15,18,20	0
2	BCL	A	403	66/66	0.97	0.06	11,16,33,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BCL A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL A 405:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL A 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BCL A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BCL A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.