



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 01:09 AM UTC

PDB ID : 7VF3 / pdb_00007vf3
Title : Plexin B1 extracellular fragment in complex with lasso-grafted PB1m7 peptide
Authors : Sugano, N.N.; Hirata, K.; Yamashita, K.; Yamamoto, M.; Arimori, T.; Takagi, J.
Deposited on : 2021-09-10
Resolution : 2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	525	6% 87% 7% 6%
1	C	525	7% 87% 7% 6%
2	B	155	2% 10% 90%
2	D	155	3% 9% 90%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7997 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plexin-B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	22	0	0
			3751	2357	656	716	22			
1	C	494	Total	C	N	O	S	3	0	0
			3745	2354	653	716	22			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	SER	THR	engineered mutation	UNP O43157
A	536	THR	-	expression tag	UNP O43157
A	537	ARG	-	expression tag	UNP O43157
A	538	GLU	-	expression tag	UNP O43157
A	539	ASN	-	expression tag	UNP O43157
A	540	LEU	-	expression tag	UNP O43157
A	541	TYR	-	expression tag	UNP O43157
A	542	PHE	-	expression tag	UNP O43157
A	543	GLN	-	expression tag	UNP O43157
C	19	SER	THR	engineered mutation	UNP O43157
C	536	THR	-	expression tag	UNP O43157
C	537	ARG	-	expression tag	UNP O43157
C	538	GLU	-	expression tag	UNP O43157
C	539	ASN	-	expression tag	UNP O43157
C	540	LEU	-	expression tag	UNP O43157
C	541	TYR	-	expression tag	UNP O43157
C	542	PHE	-	expression tag	UNP O43157
C	543	GLN	-	expression tag	UNP O43157

- Molecule 2 is a protein called Uteroglobin,PB1m7 peptide,Uteroglobin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	S	0	0	0
			129	83	20	24	2			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	15	Total	C	N	O	S	0	0	0
			129	83	20	24	2			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	29	SER	ALA	engineered mutation	UNP P11684
D	29	SER	ALA	engineered mutation	UNP P11684

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			13	8	5		
6	C	1	Total	C	O	0	0
			13	8	5		
6	C	1	Total	C	O	0	0
			13	8	5		
6	C	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	64	Total	O	0	0
			64	64		
7	B	2	Total	O	0	0
			2	2		
7	C	77	Total	O	0	0
			77	77		
7	D	3	Total	O	0	0
			3	3		

THR
LEU
PRO
GLN
LYS
PRO
ARG
GLU
SER
ILE
ILE
LYS
LEU
MET
GLU
LYS
ILE
ALA
GLN
SER
SER
SER
LEU
CYS
ASN

● Molecule 2: Uteroglobin,PB1m7 peptide,Uteroglobin



SER
GLU
ILE
CYS
PRO
SER
PHE
GLN
ARG
VAL
GLY
ILE
ILE
GLU
THR
LEU
LEU
MET
ASP
THR
PRO
SER
SER
TYR
GLU
SER
ALA
ALA
MET
GLU
PHE
SER
PRO
ASP
GLN
MET
THR
ARG
GLU
ALA
GLY
ALA
GLN
LEU
LYS
LYS
LEU
VAL
ASP
THR
PRO
GLN
LYS
PRO
ARG
GLU
SER
ILE
ILE
LYS
LEU

MET
GLU
LYS
ILE
ALA
GLN
SER
SER
SER
LEU
SER
GLY
C100
N101
S102
N103
W112
Y113
C114
GLY
SER
SER
PRO
ALA
GLN
PHE
SER
GLN
ARG
VAL
ILE
GLU
THR
LEU
LEU
MET
ASP
THR
THR
PRO
SER
SER
TYR
GLU
ALA
ALA
MET
GLU
LEU
PHE
SER
SER
PRO
ASP
GLN
ASP
MET
ARG
GLU
ALA
GLY
ALA
GLN
LYS

LYS
LEU
VAL
ASP
THR
LEU
PRO
GLN
LYS
PRO
ARG
GLU
SER
SER
ILE
ILE
LYS
LEU
MET
GLU
LYS
ILE
ALA
GLN
SER
SER
LEU
CYS
ASN

4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	263.59Å 263.59Å 108.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.41 – 2.29 47.41 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.41-2.29) 99.7 (47.41-2.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.196 , 0.226 0.196 , 0.226	Depositor DCC
R_{free} test set	4233 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	44.9	Xtriage
Anisotropy	0.446	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7997	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PEG, PG4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/3846	0.52	0/5252
1	C	0.34	0/3840	0.54	0/5245
2	B	0.27	0/134	0.40	0/184
2	D	0.33	0/134	0.37	0/184
All	All	0.33	0/7954	0.53	0/10865

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3751	0	3601	22	0
1	C	3745	0	3593	23	0
2	B	129	0	107	0	0
2	D	129	0	107	1	0
3	A	14	0	13	0	0
3	C	14	0	13	0	0
4	A	7	0	10	0	0
5	C	10	0	14	1	0
6	C	52	0	72	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	64	0	0	1	0
7	B	2	0	0	0	0
7	C	77	0	0	0	0
7	D	3	0	0	0	0
All	All	7997	0	7530	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:503:VAL:HG21	1:C:522:GLN:HE21	1.40	0.86
1:A:77:ARG:NH1	1:A:146:ASN:O	2.19	0.76
1:C:183:PRO:HD3	1:C:211:LEU:HD11	1.67	0.74
1:C:346:ARG:HE	5:C:602:PGE:H32	1.53	0.72
1:A:363:ASP:OD2	1:C:123:ARG:NH2	2.25	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/525 (93%)	476 (98%)	10 (2%)	0	100	100
1	C	486/525 (93%)	478 (98%)	8 (2%)	0	100	100
2	B	13/155 (8%)	13 (100%)	0	0	100	100
2	D	13/155 (8%)	12 (92%)	1 (8%)	0	100	100
All	All	998/1360 (73%)	979 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/424 (95%)	396 (99%)	5 (1%)	63	79
1	C	400/424 (94%)	397 (99%)	3 (1%)	73	86
2	B	15/141 (11%)	15 (100%)	0	100	100
2	D	15/141 (11%)	15 (100%)	0	100	100
All	All	831/1130 (74%)	823 (99%)	8 (1%)	68	82

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	534	LEU
1	C	527	PHE
1	A	527	PHE
1	A	414	ASP
1	C	468	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	427	GLN
1	C	522	GLN
1	C	57	GLN
1	C	63	GLN
1	C	256	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PG4	C	604	-	12,12,12	0.19	0	11,11,11	0.59	0
6	PG4	C	605	-	12,12,12	0.25	0	11,11,11	0.48	0
5	PGE	C	602	-	9,9,9	0.40	0	8,8,8	0.33	0
6	PG4	C	606	-	12,12,12	0.26	0	11,11,11	0.51	0
4	PEG	A	602	-	6,6,6	0.24	0	5,5,5	0.06	0
6	PG4	C	603	-	12,12,12	0.24	0	11,11,11	0.55	0
3	NAG	A	601	1	14,14,15	0.72	1 (7%)	17,19,21	0.67	1 (5%)
3	NAG	C	601	1	14,14,15	0.50	0	17,19,21	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PG4	C	604	-	-	7/10/10/10	-
6	PG4	C	605	-	-	5/10/10/10	-
5	PGE	C	602	-	-	4/7/7/7	-
6	PG4	C	606	-	-	5/10/10/10	-
4	PEG	A	602	-	-	2/4/4/4	-
6	PG4	C	603	-	-	1/10/10/10	-
3	NAG	A	601	1	-	2/6/23/26	0/1/1/1
3	NAG	C	601	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	NAG	O5-C1	2.31	1.47	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	NAG	C1-O5-C5	2.24	115.18	112.19

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	NAG	O5-C5-C6-O6
4	A	602	PEG	O1-C1-C2-O2
6	C	605	PG4	O2-C3-C4-O3
6	C	605	PG4	O3-C5-C6-O4
3	A	601	NAG	C4-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	604	PG4	2	0
6	C	605	PG4	1	0
5	C	602	PGE	1	0
6	C	603	PG4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	494/525 (94%)	0.22	34 (6%)	23 25	33, 46, 81, 113	5 (1%)
1	C	494/525 (94%)	0.22	36 (7%)	21 23	32, 47, 80, 116	1 (0%)
2	B	15/155 (9%)	0.85	3 (20%)	3 3	40, 49, 114, 123	0
2	D	15/155 (9%)	1.04	5 (33%)	1 1	39, 47, 105, 117	0
All	All	1018/1360 (74%)	0.24	78 (7%)	19 21	32, 47, 82, 123	6 (0%)

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	520	PRO	9.1
1	C	534	LEU	9.0
1	A	177	VAL	7.0
1	A	24	PRO	6.5
2	B	100	CYS	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PG4	C	606	13/13	0.74	0.29	53,78,89,91	0
6	PG4	C	604	13/13	0.77	0.23	63,72,84,84	0
6	PG4	C	605	13/13	0.83	0.18	55,69,77,80	0
4	PEG	A	602	7/7	0.87	0.18	60,63,71,79	0
5	PGE	C	602	10/10	0.88	0.17	52,61,72,74	0
6	PG4	C	603	13/13	0.91	0.14	45,57,64,67	0
3	NAG	A	601	14/15	0.92	0.12	46,53,70,75	0
3	NAG	C	601	14/15	0.92	0.10	41,50,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.