



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 10:59 PM UTC

PDB ID : 2VHS / pdb\_00002vhs  
Title : Cathsilicatein, a chimera  
Authors : Fairhead, M.; Kowatz, T.; McMahon, S.A.; Carter, L.G.; Oke, M.; Johnson, K.A.; Liu, H.; Naismith, J.H.; Wal, C.F.V.D.  
Deposited on : 2007-11-24  
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	 29% 90% 9%
1	B	217	 34% 93% 6%
1	C	217	 23% 93% 6%
1	D	217	 24% 91% 8%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATHSILICATEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	1661	1047	275	326	13	0	3	0
1	B	215	1657	1043	275	326	13	0	2	0
1	C	215	1661	1047	275	326	13	0	3	0
1	D	215	1657	1043	275	326	13	0	2	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	VAL	THR	conflict	UNP P07711
B	110	VAL	THR	conflict	UNP P07711
C	110	VAL	THR	conflict	UNP P07711
D	110	VAL	THR	conflict	UNP P07711
A	24	ALA	SER	engineered mutation	UNP P07711
A	25	SER	CYS	engineered mutation	UNP P07711
A	26	TYR	TRP	engineered mutation	UNP P07711
A	153	SER	GLU	engineered mutation	UNP P07711
A	154	SER	PRO	engineered mutation	UNP P07711
A	159	SER	GLU	engineered mutation	UNP P07711
A	160	SER	ASP	engineered mutation	UNP P07711
A	161	LEU	MET	engineered mutation	UNP P07711
A	162	ASN	ASP	engineered mutation	UNP P07711
A	164	ALA	GLY	engineered mutation	UNP P07711
A	165	MET	VAL	engineered mutation	UNP P07711
B	24	ALA	SER	engineered mutation	UNP P07711
B	25	SER	CYS	engineered mutation	UNP P07711
B	26	TYR	TRP	engineered mutation	UNP P07711
B	153	SER	GLU	engineered mutation	UNP P07711
B	154	SER	PRO	engineered mutation	UNP P07711
B	159	SER	GLU	engineered mutation	UNP P07711

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	160	SER	ASP	engineered mutation	UNP P07711
B	161	LEU	MET	engineered mutation	UNP P07711
B	162	ASN	ASP	engineered mutation	UNP P07711
B	164	ALA	GLY	engineered mutation	UNP P07711
B	165	MET	VAL	engineered mutation	UNP P07711
C	24	ALA	SER	engineered mutation	UNP P07711
C	25	SER	CYS	engineered mutation	UNP P07711
C	26	TYR	TRP	engineered mutation	UNP P07711
C	153	SER	GLU	engineered mutation	UNP P07711
C	154	SER	PRO	engineered mutation	UNP P07711
C	159	SER	GLU	engineered mutation	UNP P07711
C	160	SER	ASP	engineered mutation	UNP P07711
C	161	LEU	MET	engineered mutation	UNP P07711
C	162	ASN	ASP	engineered mutation	UNP P07711
C	164	ALA	GLY	engineered mutation	UNP P07711
C	165	MET	VAL	engineered mutation	UNP P07711
D	24	ALA	SER	engineered mutation	UNP P07711
D	25	SER	CYS	engineered mutation	UNP P07711
D	26	TYR	TRP	engineered mutation	UNP P07711
D	153	SER	GLU	engineered mutation	UNP P07711
D	154	SER	PRO	engineered mutation	UNP P07711
D	159	SER	GLU	engineered mutation	UNP P07711
D	160	SER	ASP	engineered mutation	UNP P07711
D	161	LEU	MET	engineered mutation	UNP P07711
D	162	ASN	ASP	engineered mutation	UNP P07711
D	164	ALA	GLY	engineered mutation	UNP P07711
D	165	MET	VAL	engineered mutation	UNP P07711

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	303	Total O 303 303	0	0
3	B	264	Total O 264 264	0	0
3	C	307	Total O 307 307	0	0
3	D	298	Total O 298 298	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CATHSILICATEIN




- Molecule 1: CATHSILICATEIN




- Molecule 1: CATHSILICATEIN



- Molecule 1: CATHSILICATEIN





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.83Å 58.10Å 70.21Å 105.67° 104.97° 105.05°	Depositor
Resolution (Å)	63.25 – 1.50 63.24 – 1.50	Depositor EDS
% Data completeness (in resolution range)	91.7 (63.25-1.50) 80.0 (63.24-1.50)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.188 , 0.220 0.196 , 0.233	Depositor DCC
$R_{free}$ test set	5042 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtrriage
Anisotropy	0.097	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.036 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7848	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 85.58 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2083e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/1709	0.64	0/2306
1	B	0.41	0/1702	0.62	0/2296
1	C	0.43	0/1709	0.63	0/2306
1	D	0.42	0/1702	0.63	0/2296
All	All	0.42	0/6822	0.63	0/9204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1661	0	1576	10	0
1	B	1657	0	1567	7	0
1	C	1661	0	1576	7	0
1	D	1657	0	1567	11	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
3	A	303	0	0	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	264	0	0	0	0
3	C	307	0	0	1	0
3	D	298	0	0	2	0
All	All	7848	0	6286	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:TYR:OH	1:B:213:SER:OG	2.10	0.69
1:A:202:ALA:HB1	1:A:205:ARG:HD3	1.78	0.66
1:B:123:MET:HE3	1:B:170:TYR:CE1	2.37	0.60
1:C:202:ALA:HB1	1:C:205:ARG:HD3	1.85	0.57
1:D:165:MET:HE2	1:D:185:VAL:HG12	1.89	0.54

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/217 (99%)	211 (99%)	3 (1%)	0	100	100
1	B	213/217 (98%)	209 (98%)	4 (2%)	0	100	100
1	C	214/217 (99%)	211 (99%)	3 (1%)	0	100	100
1	D	213/217 (98%)	209 (98%)	4 (2%)	0	100	100
All	All	854/868 (98%)	840 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/175 (101%)	171 (97%)	5 (3%)	38	11
1	B	175/175 (100%)	174 (99%)	1 (1%)	78	62
1	C	176/175 (101%)	175 (99%)	1 (1%)	78	62
1	D	175/175 (100%)	174 (99%)	1 (1%)	78	62
All	All	702/700 (100%)	694 (99%)	8 (1%)	63	41

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	156	CYS
1	C	156	CYS
1	A	211	ILE
1	A	174	SER
1	B	156	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	78	GLN
1	D	78	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	C	1221	-	4,4,4	0.18	0	6,6,6	0.26	0
2	SO4	B	1221	-	4,4,4	0.25	0	6,6,6	0.16	0
2	SO4	D	1221	-	4,4,4	0.28	0	6,6,6	0.29	0
2	SO4	C	1222	-	4,4,4	0.22	0	6,6,6	0.13	0
2	SO4	A	1221	-	4,4,4	0.23	0	6,6,6	0.26	0
2	SO4	B	1222	-	4,4,4	0.23	0	6,6,6	0.12	0
2	SO4	D	1222	-	4,4,4	0.26	0	6,6,6	0.09	0
2	SO4	A	1222	-	4,4,4	0.23	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/217 (99%)	1.61	63 (29%) <b>1</b> <b>1</b>	14, 24, 29, 31	3 (1%)
1	B	215/217 (99%)	1.71	73 (33%) <b>1</b> <b>1</b>	14, 24, 29, 31	2 (0%)
1	C	215/217 (99%)	1.47	50 (23%) <b>2</b> <b>2</b>	14, 24, 29, 31	3 (1%)
1	D	215/217 (99%)	1.56	53 (24%) <b>2</b> <b>2</b>	14, 23, 29, 31	2 (0%)
All	All	860/868 (99%)	1.59	239 (27%) <b>1</b> <b>1</b>	14, 24, 29, 31	10 (1%)

The worst 5 of 239 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	139	GLY	5.4
1	D	174	SER	5.4
1	A	152	PHE	5.0
1	B	1	ALA	4.9
1	C	174	SER	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	1222	5/5	0.79	0.14	52,52,52,52	0
2	SO4	A	1222	5/5	0.84	0.16	45,45,46,46	0
2	SO4	C	1222	5/5	0.87	0.14	38,38,39,39	0
2	SO4	D	1222	5/5	0.87	0.15	41,41,42,43	0
2	SO4	D	1221	5/5	0.94	0.10	26,27,30,31	0
2	SO4	B	1221	5/5	0.95	0.11	26,27,29,31	0
2	SO4	C	1221	5/5	0.96	0.10	25,28,29,31	0
2	SO4	A	1221	5/5	0.96	0.09	25,28,28,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.