



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 05:54 AM UTC

PDB ID : 1VIB / pdb_00001vib
Title : NMR SOLUTION STRUCTURE OF THE NEUROTOXIN B-IV, 20 STRUCTURES
Authors : Barnham, K.J.; Dyke, T.R.; Kem, W.R.; Norton, R.S.
Deposited on : 1996-11-25

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	55	

2 Ensemble composition and analysis

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:13-A:50 (38)	0.51	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 4, 5, 6, 11, 12, 13
2	7, 9, 15, 19
3	2, 8, 10, 16
4	14, 18
Single-model clusters	3; 17; 20

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 845 atoms, of which 422 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called NEUROTOXIN B-IV.

Mol	Chain	Residues	Atoms						Trace
1	A	55	Total	C	H	N	O	S	0
			845	258	422	84	73	8	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	10	HYP	PRO	conflict	UNP P01525

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: NEUROTOXIN B-IV




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: NEUROTOXIN B-IV




4.2.2 Score per residue for model 2

- Molecule 1: NEUROTOXIN B-IV




4.2.3 Score per residue for model 3

- Molecule 1: NEUROTOXIN B-IV




4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: NEUROTOXIN B-IV




4.2.5 Score per residue for model 5

- Molecule 1: NEUROTOXIN B-IV




4.2.6 Score per residue for model 6

- Molecule 1: NEUROTOXIN B-IV




4.2.7 Score per residue for model 7

- Molecule 1: NEUROTOXIN B-IV




4.2.8 Score per residue for model 8

- Molecule 1: NEUROTOXIN B-IV




4.2.9 Score per residue for model 9

- Molecule 1: NEUROTOXIN B-IV




4.2.10 Score per residue for model 10

- Molecule 1: NEUROTOXIN B-IV




4.2.11 Score per residue for model 11

- Molecule 1: NEUROTOXIN B-IV




4.2.12 Score per residue for model 12

- Molecule 1: NEUROTOXIN B-IV




4.2.13 Score per residue for model 13

- Molecule 1: NEUROTOXIN B-IV




4.2.14 Score per residue for model 14

- Molecule 1: NEUROTOXIN B-IV




4.2.15 Score per residue for model 15

- Molecule 1: NEUROTOXIN B-IV




4.2.16 Score per residue for model 16

- Molecule 1: NEUROTOXIN B-IV




4.2.17 Score per residue for model 17

- Molecule 1: NEUROTOXIN B-IV




4.2.18 Score per residue for model 18

- Molecule 1: NEUROTOXIN B-IV




4.2.19 Score per residue for model 19

- Molecule 1: NEUROTOXIN B-IV



4.2.20 Score per residue for model 20

- Molecule 1: NEUROTOXIN B-IV



5 Refinement protocol and experimental data overview ⓘ

Of the ? calculated structures, 20 were deposited, based on the following criterion: ?.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.84±0.02	4±1/302 (1.2± 0.2%)	1.83±0.07	7±2/399 (1.8± 0.5%)
All	All	1.84	74/6040 (1.2%)	1.83	142/7980 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±0.2
All	All	0	59

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	40	HIS	CG-ND1	-7.42	1.30	1.38	16	20
1	A	40	HIS	CD2-NE2	-6.31	1.30	1.37	14	20
1	A	30	TRP	NE1-CE2	-6.09	1.30	1.37	1	20
1	A	37	CYS	N-CA	5.46	1.52	1.46	6	9
1	A	37	CYS	CA-CB	5.23	1.61	1.53	7	2
1	A	24	ILE	CA-CB	5.10	1.61	1.54	7	2
1	A	40	HIS	CB-CG	5.09	1.57	1.50	20	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	25	ARG	N-CA-C	-10.41	97.53	110.65	9	16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	40	HIS	ND1-CE1-NE2	9.72	118.12	108.40	14	20
1	A	33	LYS	N-CA-C	-9.57	101.81	112.72	10	15
1	A	35	GLY	N-CA-C	-9.07	103.87	113.58	10	1
1	A	18	LYS	N-CA-C	-9.06	101.38	111.07	2	13
1	A	13	GLU	N-CA-C	-7.87	102.47	112.93	12	4
1	A	17	ARG	N-CA-C	-7.37	101.73	111.24	12	9
1	A	47	ASN	N-CA-C	-7.21	102.82	111.69	5	3
1	A	40	HIS	CE1-NE2-CD2	-7.18	101.82	109.00	14	20
1	A	40	HIS	CA-CB-CG	-6.34	107.46	113.80	6	8
1	A	42	ILE	N-CA-CB	-6.26	106.50	111.64	18	2
1	A	45	LYS	N-CA-C	-6.24	105.82	113.50	2	1
1	A	44	GLN	N-CA-C	-6.14	103.32	111.24	18	1
1	A	14	ASN	CA-CB-CG	-5.89	106.71	112.60	19	2
1	A	46	ASN	OD1-CG-ND2	-5.89	116.71	122.60	19	3
1	A	36	LYS	N-CA-C	-5.89	105.76	113.12	10	2
1	A	30	TRP	N-CA-C	-5.82	104.58	112.26	8	1
1	A	15	ASN	OD1-CG-ND2	-5.77	116.83	122.60	9	8
1	A	27	GLN	N-CA-C	-5.72	103.26	108.75	4	2
1	A	48	CYS	N-CA-C	-5.58	104.59	111.40	18	1
1	A	42	ILE	N-CA-C	-5.48	107.29	111.62	18	1
1	A	44	GLN	OE1-CD-NE2	-5.36	117.24	122.60	1	2
1	A	32	GLY	N-CA-C	-5.32	108.15	114.48	9	1
1	A	37	CYS	N-CA-C	-5.22	105.59	111.28	7	2
1	A	46	ASN	CA-CB-CG	-5.12	107.48	112.60	15	2
1	A	47	ASN	CA-CB-CG	-5.09	107.51	112.60	3	1
1	A	24	ILE	CA-C-O	5.03	127.07	120.78	2	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	17	ARG	Sidechain	20
1	A	34	ARG	Sidechain	20
1	A	25	ARG	Sidechain	19

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes

averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	298	301	301	24±3
All	All	5960	6020	6020	470

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:40:HIS:O	1:A:43:ILE:HG22	0.73	1.83	11	1
1:A:24:ILE:HG23	1:A:26:CYS:SG	0.73	2.23	6	18
1:A:23:CYS:C	1:A:41:CYS:SG	0.68	2.77	19	10
1:A:24:ILE:HG23	1:A:37:CYS:SG	0.66	2.30	9	3
1:A:30:TRP:CG	1:A:37:CYS:HB2	0.64	2.28	14	5
1:A:34:ARG:HD3	1:A:37:CYS:SG	0.64	2.32	20	1
1:A:30:TRP:CE3	1:A:37:CYS:HA	0.63	2.28	17	20
1:A:26:CYS:HA	1:A:30:TRP:HB2	0.63	1.71	7	19
1:A:15:ASN:ND2	1:A:48:CYS:SG	0.63	2.72	3	1
1:A:37:CYS:SG	1:A:38:ALA:N	0.62	2.71	20	11
1:A:30:TRP:CD2	1:A:37:CYS:HB2	0.62	2.29	14	5
1:A:26:CYS:SG	1:A:34:ARG:HD2	0.62	2.34	7	5
1:A:43:ILE:HG22	1:A:47:ASN:ND2	0.62	2.09	14	1
1:A:43:ILE:HG23	1:A:44:GLN:NE2	0.61	2.11	6	2
1:A:24:ILE:HG22	1:A:24:ILE:O	0.60	1.97	12	17
1:A:27:GLN:HE21	1:A:27:GLN:N	0.60	1.95	17	17
1:A:20:TYR:CE1	1:A:24:ILE:HD11	0.59	2.33	3	4
1:A:34:ARG:HD2	1:A:37:CYS:SG	0.58	2.38	8	7
1:A:40:HIS:HA	1:A:43:ILE:HD12	0.57	1.76	16	7
1:A:14:ASN:HD22	1:A:14:ASN:N	0.57	1.97	7	5
1:A:13:GLU:HA	1:A:48:CYS:SG	0.57	2.39	11	5
1:A:20:TYR:CD2	1:A:24:ILE:HD13	0.56	2.35	20	1
1:A:43:ILE:HG23	1:A:44:GLN:HE22	0.56	1.59	6	2
1:A:14:ASN:ND2	1:A:15:ASN:H	0.55	2.00	3	1
1:A:34:ARG:HA	1:A:37:CYS:SG	0.55	2.41	1	7
1:A:27:GLN:N	1:A:27:GLN:HE21	0.54	2.00	8	2
1:A:25:ARG:C	1:A:27:GLN:H	0.53	2.12	10	12
1:A:15:ASN:OD1	1:A:15:ASN:N	0.53	2.42	18	7
1:A:30:TRP:CE3	1:A:37:CYS:CA	0.52	2.92	2	20
1:A:30:TRP:CD2	1:A:37:CYS:HA	0.52	2.40	20	14
1:A:23:CYS:O	1:A:25:ARG:N	0.52	2.43	7	9
1:A:30:TRP:CZ3	1:A:37:CYS:HA	0.52	2.39	17	9

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:TRP:O	1:A:32:GLY:N	0.52	2.43	7	16
1:A:38:ALA:O	1:A:39:ALA:C	0.52	2.53	20	15
1:A:34:ARG:O	1:A:37:CYS:SG	0.51	2.68	1	1
1:A:26:CYS:SG	1:A:34:ARG:CD	0.51	2.99	14	2
1:A:15:ASN:O	1:A:19:LYS:HB2	0.51	2.06	2	2
1:A:19:LYS:O	1:A:23:CYS:N	0.50	2.44	18	11
1:A:15:ASN:CG	1:A:16:CYS:N	0.50	2.68	7	4
1:A:27:GLN:O	1:A:28:GLY:C	0.50	2.54	20	1
1:A:40:HIS:ND1	1:A:40:HIS:C	0.50	2.69	14	10
1:A:40:HIS:C	1:A:40:HIS:HD1	0.49	2.15	20	6
1:A:27:GLN:N	1:A:27:GLN:NE2	0.49	2.61	15	5
1:A:25:ARG:C	1:A:27:GLN:N	0.49	2.69	16	18
1:A:42:ILE:CG2	1:A:43:ILE:N	0.49	2.76	6	6
1:A:30:TRP:CD2	1:A:37:CYS:CB	0.48	2.96	14	4
1:A:46:ASN:O	1:A:47:ASN:C	0.48	2.56	7	6
1:A:29:LYS:HG2	1:A:30:TRP:CD1	0.48	2.44	15	2
1:A:39:ALA:O	1:A:43:ILE:HG22	0.47	2.09	6	1
1:A:36:LYS:O	1:A:37:CYS:C	0.47	2.57	2	9
1:A:27:GLN:O	1:A:29:LYS:N	0.47	2.48	18	7
1:A:24:ILE:N	1:A:41:CYS:SG	0.46	2.88	7	5
1:A:24:ILE:O	1:A:24:ILE:CG2	0.46	2.63	12	7
1:A:20:TYR:CD1	1:A:21:ASP:N	0.46	2.84	20	1
1:A:34:ARG:CD	1:A:37:CYS:SG	0.46	3.04	1	2
1:A:30:TRP:CZ3	1:A:37:CYS:N	0.46	2.84	12	3
1:A:24:ILE:HG13	1:A:41:CYS:SG	0.46	2.51	4	3
1:A:44:GLN:HE21	1:A:44:GLN:N	0.46	2.09	18	2
1:A:20:TYR:CD1	1:A:24:ILE:HD12	0.45	2.47	14	1
1:A:33:LYS:O	1:A:34:ARG:C	0.45	2.59	5	11
1:A:48:CYS:O	1:A:49:LYS:C	0.45	2.59	18	1
1:A:26:CYS:N	1:A:37:CYS:SG	0.45	2.90	19	1
1:A:47:ASN:OD1	1:A:48:CYS:N	0.45	2.50	2	1
1:A:37:CYS:O	1:A:41:CYS:SG	0.45	2.75	17	1
1:A:14:ASN:N	1:A:14:ASN:ND2	0.45	2.63	7	2
1:A:36:LYS:C	1:A:38:ALA:N	0.44	2.75	11	4
1:A:14:ASN:CG	1:A:15:ASN:N	0.44	2.75	14	1
1:A:42:ILE:HG22	1:A:43:ILE:N	0.44	2.28	3	3
1:A:20:TYR:CD1	1:A:20:TYR:C	0.44	2.96	17	1
1:A:42:ILE:C	1:A:44:GLN:N	0.44	2.75	18	4
1:A:20:TYR:CD2	1:A:24:ILE:CD1	0.44	3.01	20	1
1:A:13:GLU:O	1:A:13:GLU:CD	0.43	2.61	15	1
1:A:18:LYS:O	1:A:19:LYS:C	0.43	2.60	17	3

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:GLN:N	1:A:44:GLN:NE2	0.43	2.67	9	4
1:A:24:ILE:CG2	1:A:26:CYS:SG	0.43	3.06	3	1
1:A:44:GLN:O	1:A:45:LYS:C	0.43	2.61	18	1
1:A:25:ARG:O	1:A:27:GLN:N	0.43	2.52	2	1
1:A:20:TYR:CD1	1:A:42:ILE:HA	0.43	2.48	17	1
1:A:30:TRP:O	1:A:33:LYS:N	0.43	2.52	9	1
1:A:20:TYR:CE1	1:A:24:ILE:HD12	0.42	2.49	14	1
1:A:24:ILE:C	1:A:25:ARG:HE	0.42	2.23	6	1
1:A:26:CYS:HB3	1:A:34:ARG:CD	0.42	2.43	10	2
1:A:18:LYS:O	1:A:22:LEU:N	0.42	2.53	16	1
1:A:20:TYR:CE1	1:A:42:ILE:HG13	0.42	2.49	12	4
1:A:27:GLN:NE2	1:A:27:GLN:N	0.42	2.68	1	1
1:A:44:GLN:N	1:A:44:GLN:HE21	0.42	2.13	9	2
1:A:34:ARG:O	1:A:35:GLY:C	0.42	2.62	20	1
1:A:33:LYS:N	1:A:33:LYS:CD	0.42	2.82	5	2
1:A:44:GLN:HE21	1:A:44:GLN:CA	0.42	2.28	5	1
1:A:23:CYS:HA	1:A:25:ARG:CZ	0.42	2.44	9	1
1:A:40:HIS:C	1:A:40:HIS:ND1	0.42	2.77	5	1
1:A:42:ILE:O	1:A:46:ASN:ND2	0.41	2.53	17	1
1:A:14:ASN:N	1:A:14:ASN:HD22	0.41	2.14	3	2
1:A:24:ILE:CG1	1:A:38:ALA:HA	0.41	2.45	20	1
1:A:28:GLY:O	1:A:29:LYS:C	0.41	2.64	18	1
1:A:13:GLU:O	1:A:14:ASN:C	0.41	2.64	12	1
1:A:19:LYS:O	1:A:20:TYR:C	0.40	2.64	18	1
1:A:34:ARG:O	1:A:36:LYS:N	0.40	2.54	20	1
1:A:33:LYS:HD3	1:A:33:LYS:H	0.40	1.76	5	1
1:A:42:ILE:O	1:A:43:ILE:C	0.40	2.64	3	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	38/55 (69%)	26±2 (69±6%)	8±2 (22±6%)	4±1 (9±3%)	1	10
All	All	760/1100 (69%)	522 (69%)	166 (22%)	72 (9%)	1	10

All 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	31	ALA	18
1	A	28	GLY	15
1	A	35	GLY	11
1	A	38	ALA	10
1	A	24	ILE	9
1	A	26	CYS	5
1	A	34	ARG	1
1	A	23	CYS	1
1	A	15	ASN	1
1	A	50	GLY	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	31/41 (76%)	19±2 (62±7%)	12±2 (38±7%)	0	7
All	All	620/820 (76%)	387 (62%)	233 (38%)	0	7

All 26 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	27	GLN	20
1	A	42	ILE	19
1	A	44	GLN	19
1	A	20	TYR	18
1	A	48	CYS	16
1	A	45	LYS	14
1	A	24	ILE	12
1	A	25	ARG	11
1	A	34	ARG	11
1	A	19	LYS	10
1	A	23	CYS	10
1	A	21	ASP	9
1	A	36	LYS	8
1	A	14	ASN	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	17	ARG	7
1	A	46	ASN	6
1	A	18	LYS	5
1	A	33	LYS	5
1	A	37	CYS	5
1	A	49	LYS	5
1	A	29	LYS	4
1	A	13	GLU	3
1	A	41	CYS	3
1	A	16	CYS	2
1	A	43	ILE	2
1	A	15	ASN	2

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	HYP	A	10	1	7,8,9	0.65±0.08	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Counts	Bond angles	
						RMSZ	#Z>2
1	HYP	A	10	1	5,10,12	2.20±0.40	2±1 (45±16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HYP	A	10	1	-	0±0,0,11,13	0±0,1,1,1

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	10	HYP	CB-CG-CD	5.18	108.92	103.16	19	11
1	A	10	HYP	CG-CB-CA	4.14	98.97	103.75	18	4
1	A	10	HYP	OD1-CG-CB	3.19	102.36	110.10	20	20
1	A	10	HYP	OD1-CG-CD	3.16	103.84	110.30	17	10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided