



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:35 PM UTC

PDB ID : 1VIQ / pdb_00001viq
Title : Crystal structure of putative ADP ribose pyrophosphatase
Authors : Structural GenomiX
Deposited on : 2003-12-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	
1	B	220	
1	C	220	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribose pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	0	0
			1597	1010	281	302	4			
1	B	201	Total	C	N	O	S	0	1	0
			1595	1006	285	300	4			
1	C	194	Total	C	N	O	S	0	0	0
			1533	969	270	290	4			

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	cloning artifact	UNP Q93K97
A	0	SER	-	cloning artifact	UNP Q93K97
A	28	SER	PHE	variant	UNP Q93K97
A	209	GLU	-	cloning artifact	UNP Q93K97
A	210	GLY	-	cloning artifact	UNP Q93K97
A	211	GLY	-	cloning artifact	UNP Q93K97
A	212	SER	-	cloning artifact	UNP Q93K97
A	213	HIS	-	cloning artifact	UNP Q93K97
A	214	HIS	-	cloning artifact	UNP Q93K97
A	215	HIS	-	cloning artifact	UNP Q93K97
A	216	HIS	-	cloning artifact	UNP Q93K97
A	217	HIS	-	cloning artifact	UNP Q93K97
A	218	HIS	-	cloning artifact	UNP Q93K97
B	-1	MET	-	cloning artifact	UNP Q93K97
B	0	SER	-	cloning artifact	UNP Q93K97
B	28	SER	PHE	variant	UNP Q93K97
B	209	GLU	-	cloning artifact	UNP Q93K97
B	210	GLY	-	cloning artifact	UNP Q93K97
B	211	GLY	-	cloning artifact	UNP Q93K97
B	212	SER	-	cloning artifact	UNP Q93K97
B	213	HIS	-	cloning artifact	UNP Q93K97
B	214	HIS	-	cloning artifact	UNP Q93K97
B	215	HIS	-	cloning artifact	UNP Q93K97

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	216	HIS	-	cloning artifact	UNP Q93K97
B	217	HIS	-	cloning artifact	UNP Q93K97
B	218	HIS	-	cloning artifact	UNP Q93K97
C	-1	MET	-	cloning artifact	UNP Q93K97
C	0	SER	-	cloning artifact	UNP Q93K97
C	28	SER	PHE	variant	UNP Q93K97
C	209	GLU	-	cloning artifact	UNP Q93K97
C	210	GLY	-	cloning artifact	UNP Q93K97
C	211	GLY	-	cloning artifact	UNP Q93K97
C	212	SER	-	cloning artifact	UNP Q93K97
C	213	HIS	-	cloning artifact	UNP Q93K97
C	214	HIS	-	cloning artifact	UNP Q93K97
C	215	HIS	-	cloning artifact	UNP Q93K97
C	216	HIS	-	cloning artifact	UNP Q93K97
C	217	HIS	-	cloning artifact	UNP Q93K97
C	218	HIS	-	cloning artifact	UNP Q93K97

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	80	Total O 80 80	0	0
2	B	91	Total O 91 91	0	0
2	C	80	Total O 80 80	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ADP-ribose pyrophosphatase

Chain A: 




• Molecule 1: ADP-ribose pyrophosphatase

Chain B: 



• Molecule 1: ADP-ribose pyrophosphatase

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	98.18Å 98.18Å 155.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	51.98 – 2.40 51.98 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (51.98-2.40) 99.8 (51.98-2.40)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.39Å)	Xtriage
Refinement program	REFMAC 4.0	Depositor
R, R_{free}	0.244 , 0.314 0.222 , 0.286	Depositor DCC
R_{free} test set	1764 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.830	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.047 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4976	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/1628 (0.1%)	1.03	2/2209 (0.1%)
1	B	0.84	1/1629 (0.1%)	1.04	2/2207 (0.1%)
1	C	0.79	0/1562	1.01	2/2117 (0.1%)
All	All	0.82	2/4819 (0.0%)	1.03	6/6533 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	179	VAL	CA-CB	5.81	1.60	1.54
1	A	143	MET	SD-CE	5.69	1.93	1.79

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	83	ASP	N-CA-C	5.76	119.57	112.54
1	A	35	PHE	N-CA-C	5.59	117.76	108.99
1	A	29	SER	N-CA-C	5.48	118.33	109.40
1	C	52	ILE	N-CA-C	5.37	115.58	107.80
1	B	139	ARG	N-CA-C	-5.24	99.05	108.48
1	B	175	ALA	N-CA-C	-5.06	105.46	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1597	0	1552	21	1
1	B	1595	0	1565	44	0
1	C	1533	0	1496	35	0
2	A	80	0	0	4	0
2	B	91	0	0	6	0
2	C	80	0	0	1	0
All	All	4976	0	4613	87	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:GLN:HG2	2:B:229:HOH:O	1.66	0.96
1:B:177:GLN:O	1:B:181:GLU:HG3	1.78	0.84
1:A:114:GLU:HG2	1:B:68:ARG:NH2	1.96	0.80
1:A:72:VAL:HG21	1:A:118:LEU:HD11	1.72	0.71
1:B:139:ARG:NH2	1:C:185:ASP:OD2	2.17	0.71
1:A:61:LEU:HD12	1:A:108:ALA:HB1	1.74	0.69
1:B:84:THR:CG2	1:C:139:ARG:HH11	2.07	0.68
1:A:56:GLY:HA3	1:A:97:MET:HE3	1.76	0.67
1:B:176:TYR:HB2	2:B:240:HOH:O	1.95	0.67
1:B:185:ASP:OD2	1:C:139:ARG:NH2	2.20	0.67
1:B:131:ALA:HB2	2:B:247:HOH:O	1.95	0.65
1:B:22:THR:HG21	1:B:25[B]:ARG:CZ	2.27	0.65
1:B:54:GLU:OE2	1:B:55:ARG:N	2.31	0.64
1:B:195:TRP:CD1	1:B:199:HIS:CE1	2.87	0.63
1:B:84:THR:HG23	1:C:139:ARG:HD2	1.84	0.60
1:B:84:THR:HG23	1:C:139:ARG:HH11	1.67	0.59
1:C:48:VAL:HG11	1:C:50:ARG:HH11	1.67	0.59
1:B:105:GLU:HA	1:B:143:MET:HE1	1.84	0.59
1:B:104:VAL:HG22	1:B:143:MET:HE2	1.83	0.59
1:C:152:ALA:HA	2:C:230:HOH:O	2.04	0.58
1:B:164:ASP:N	2:B:285:HOH:O	2.39	0.56
1:C:68:ARG:HD3	1:C:70:GLU:OE1	2.07	0.55
1:A:68:ARG:HD3	1:A:70:GLU:OE1	2.09	0.53
1:C:68:ARG:NH1	1:C:70:GLU:OE2	2.41	0.53
1:A:99:GLU:O	1:A:102:GLU:HB2	2.09	0.53
1:B:23:LEU:HB3	1:C:24:TYR:CE1	2.43	0.53
1:B:69:ASP:OD2	1:B:172:ARG:NH2	2.40	0.53
1:C:121:LYS:HB2	1:C:146:GLU:HB3	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:GLN:OE1	1:B:181:GLU:OE2	2.28	0.52
1:B:51:GLU:C	1:B:52:ILE:HD13	2.33	0.52
1:C:95:ALA:C	1:C:111:GLU:HG3	2.34	0.52
1:B:6:LEU:HG	1:C:91:LEU:HD22	1.92	0.52
1:B:111:GLU:O	1:B:115:GLU:HB2	2.10	0.51
1:B:200:HIS:HD2	1:B:204:LYS:NZ	2.09	0.51
1:B:51:GLU:O	1:B:52:ILE:HD13	2.11	0.50
1:C:22:THR:HA	1:C:31:ASP:OD1	2.12	0.50
1:C:163:GLU:HG2	1:C:165:ILE:HD11	1.94	0.50
1:C:104:VAL:HG11	1:C:141:SER:HB2	1.94	0.50
1:B:68:ARG:HD3	1:B:70:GLU:OE1	2.11	0.49
1:B:22:THR:HG22	1:B:31:ASP:OD1	2.12	0.49
1:B:102:GLU:OE2	1:B:110:ARG:NH1	2.35	0.49
1:B:172:ARG:HG2	1:B:172:ARG:HH11	1.78	0.49
1:B:54:GLU:OE2	1:B:56:GLY:N	2.25	0.48
1:A:107:VAL:HG22	1:A:110:ARG:HH12	1.79	0.48
1:B:80:ALA:HB3	1:C:133:PRO:O	2.14	0.48
1:B:195:TRP:NE1	1:B:199:HIS:CE1	2.83	0.47
1:B:109:ARG:NH1	1:B:120:VAL:O	2.48	0.47
1:B:75:GLU:HA	1:B:90:LEU:O	2.14	0.47
1:B:80:ALA:HB2	1:C:52:ILE:HD12	1.97	0.47
1:A:86:GLU:HB3	2:A:295:HOH:O	2.15	0.46
1:A:61:LEU:CD1	1:A:108:ALA:HB1	2.42	0.46
1:B:36:ARG:HD3	2:B:290:HOH:O	2.15	0.45
1:C:65:ASP:HA	1:C:66:PRO:HD2	1.79	0.45
1:C:48:VAL:HG11	1:C:50:ARG:NH1	2.31	0.45
1:C:12:LYS:HA	1:C:15:VAL:HG23	1.98	0.45
1:C:136:THR:HG23	1:C:138:GLU:H	1.82	0.45
1:B:84:THR:CG2	1:C:139:ARG:HD2	2.45	0.44
1:A:102:GLU:OE1	1:A:110:ARG:NH1	2.51	0.44
1:B:53:PHE:CE1	1:C:30:LEU:HD11	2.53	0.44
1:A:126:VAL:HG13	1:A:195:TRP:CG	2.53	0.44
1:B:33:TYR:O	1:B:49:ARG:HA	2.18	0.43
1:B:121:LYS:NZ	2:B:235:HOH:O	2.51	0.43
1:C:6:LEU:CB	1:C:7:PRO:CD	2.96	0.43
1:C:74:ILE:O	1:C:74:ILE:HG13	2.18	0.43
1:A:130:LEU:HD21	1:A:139:ARG:NH1	2.33	0.43
1:A:174:GLN:HA	1:A:177:GLN:HE21	1.84	0.43
1:A:18:ILE:N	1:A:34:ARG:O	2.48	0.43
1:C:176:TYR:O	1:C:179:VAL:HB	2.19	0.43
1:B:121:LYS:HB2	1:B:146:GLU:HB3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:HIS:CD2	1:B:139:ARG:HB3	2.54	0.43
1:C:126:VAL:HG12	1:C:127:LEU:N	2.33	0.43
1:B:165:ILE:CG2	1:B:166:ARG:N	2.82	0.42
1:A:39:LEU:HA	2:A:220:HOH:O	2.18	0.42
1:A:120:VAL:HG23	2:A:222:HOH:O	2.19	0.42
1:C:48:VAL:CG1	1:C:50:ARG:HH11	2.32	0.42
1:C:176:TYR:CZ	1:C:180:GLU:HG2	2.55	0.42
1:C:34:ARG:HE	1:C:47:GLU:CD	2.27	0.42
1:C:105:GLU:OE2	1:C:109:ARG:NE	2.50	0.42
1:C:38:ARG:HA	1:C:38:ARG:HD3	1.79	0.42
1:A:55:ARG:O	1:A:55:ARG:HG3	2.19	0.41
1:A:170:VAL:HG11	1:A:178:TRP:CZ3	2.55	0.41
1:C:195:TRP:O	1:C:196:LEU:C	2.63	0.41
1:A:190:VAL:O	1:A:194:GLN:HG3	2.21	0.41
1:A:121:LYS:HB3	1:A:122:ARG:H	1.68	0.41
1:B:23:LEU:HB3	1:C:24:TYR:CD1	2.56	0.41
1:A:13:ASN:HB2	2:A:229:HOH:O	2.21	0.41
1:B:24:TYR:CZ	1:B:26:GLY:HA3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ARG:NH2	1:A:185:ASP:OD2[6_556]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/220 (91%)	191 (96%)	9 (4%)	0	100	100
1	B	198/220 (90%)	181 (91%)	17 (9%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	190/220 (86%)	182 (96%)	6 (3%)	2 (1%)	11	18
All	All	588/660 (89%)	554 (94%)	32 (5%)	2 (0%)	36	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	121	LYS
1	C	100	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/186 (89%)	147 (89%)	19 (11%)	5	8
1	B	168/186 (90%)	138 (82%)	30 (18%)	2	2
1	C	160/186 (86%)	142 (89%)	18 (11%)	5	8
All	All	494/558 (88%)	427 (86%)	67 (14%)	4	5

All (67) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	15	VAL
1	A	16	GLU
1	A	17	ILE
1	A	21	GLU
1	A	28	SER
1	A	29	SER
1	A	48	VAL
1	A	61	LEU
1	A	68	ARG
1	A	85	SER
1	A	91	LEU
1	A	94	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	104	VAL
1	A	121	LYS
1	A	147	VAL
1	A	183	LYS
1	A	201	GLN
1	A	206	GLU
1	B	6	LEU
1	B	15	VAL
1	B	25[A]	ARG
1	B	25[B]	ARG
1	B	29	SER
1	B	49	ARG
1	B	50	ARG
1	B	54	GLU
1	B	71	VAL
1	B	74	ILE
1	B	77	ILE
1	B	84	THR
1	B	85	SER
1	B	91	LEU
1	B	94	VAL
1	B	97	MET
1	B	103	SER
1	B	104	VAL
1	B	105	GLU
1	B	113	ILE
1	B	115	GLU
1	B	127	LEU
1	B	139	ARG
1	B	147	VAL
1	B	153	SER
1	B	166	ARG
1	B	173	GLU
1	B	177	GLN
1	B	181	GLU
1	B	201	GLN
1	C	13	ASN
1	C	15	VAL
1	C	18	ILE
1	C	22	THR
1	C	29	SER
1	C	38	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	50	ARG
1	C	62	LEU
1	C	86	GLU
1	C	91	LEU
1	C	99	GLU
1	C	105	GLU
1	C	119	ILE
1	C	147	VAL
1	C	163	GLU
1	C	174	GLN
1	C	184	ILE
1	C	204	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	177	GLN
1	A	200	HIS
1	A	201	GLN
1	B	57	HIS
1	B	168	HIS
1	B	174	GLN
1	B	177	GLN
1	B	200	HIS
1	C	186	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/220 (92%)	-0.10	0 100 100	21, 36, 53, 59	0
1	B	201/220 (91%)	-0.15	2 (0%) 79 76	19, 32, 46, 55	1 (0%)
1	C	194/220 (88%)	0.08	4 (2%) 63 59	22, 39, 54, 59	0
All	All	599/660 (90%)	-0.06	6 (1%) 79 76	19, 35, 53, 59	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	162	ASN	3.3
1	C	6	LEU	2.4
1	B	0	SER	2.3
1	C	152	ALA	2.2
1	B	9	THR	2.1
1	C	46	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.