



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 08:19 AM EDT

PDB ID : 3VIQ / pdb_00003viq
Title : Crystal structure of Swi5-Sfr1 complex from fission yeast
Authors : Kuwabara, N.; Murayama, Y.; Hashimoto, H.; Kokabu, Y.; Ikeguchi, M.; Sato, M.; Mayanagi, K.; Tsutsui, Y.; Iwasaki, H.; Shimizu, T.
Deposited on : 2011-10-06
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	16% 85% 14% .
1	C	122	19% 83% 11% 5%
2	B	85	% 92% 8%
2	D	85	8% 75% 7% 18%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Swi5-dependent recombination DNA repair protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	122	Total	C	N	O	S	0	2	0
			1005	628	185	187	5			
1	C	116	Total	C	N	O	S	0	1	0
			940	587	169	180	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	GLY	-	expression tag	UNP Q9USV1
A	179	HIS	-	expression tag	UNP Q9USV1
A	180	MET	-	expression tag	UNP Q9USV1
C	178	GLY	-	expression tag	UNP Q9USV1
C	179	HIS	-	expression tag	UNP Q9USV1
C	180	MET	-	expression tag	UNP Q9USV1

- Molecule 2 is a protein called Mating-type switching protein swi5.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	85	Total	C	N	O	S	0	2	0
			698	431	128	136	3			
2	D	70	Total	C	N	O	S	0	1	0
			573	356	103	112	2			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	62	Total 62	O 62	0	0
6	B	42	Total 42	O 42	0	0
6	C	31	Total 31	O 31	0	0
6	D	29	Total 29	O 29	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Swi5-dependent recombination DNA repair protein 1





- Molecule 1: Swi5-dependent recombination DNA repair protein 1



- Molecule 2: Mating-type switching protein swi5



- Molecule 2: Mating-type switching protein swi5



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.16Å 128.68Å 59.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 20.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-2.20) 98.9 (20.00-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.6.0113	Depositor
R, R_{free}	0.230 , 0.266 0.238 , 0.276	Depositor DCC
R_{free} test set	1751 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3431	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1024	0.85	0/1369
1	C	0.78	0/955	0.89	0/1280
2	B	0.82	0/711	0.89	0/948
2	D	0.81	0/580	0.84	0/772
All	All	0.82	0/3270	0.87	0/4369

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1005	0	1016	16	0
1	C	940	0	932	15	0
2	B	698	0	708	10	0
2	D	573	0	580	7	0
3	A	12	0	16	0	0
3	C	24	0	32	0	0
3	D	6	0	8	1	0
4	A	4	0	0	1	0
5	C	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	62	0	0	0	0
6	B	42	0	0	1	0
6	C	31	0	0	0	0
6	D	29	0	0	2	0
All	All	3431	0	3292	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:70:CYS:SG	2:B:75:LEU:HD11	2.02	0.99
1:C:189:LEU:HD21	2:D:13:LEU:HB3	1.52	0.89
1:C:250:VAL:HG12	1:C:270:THR:O	1.72	0.88
1:C:262:LYS:O	1:C:264:GLN:N	2.20	0.73
1:A:230:GLN:HG3	1:A:288:MET:HE3	1.74	0.69
2:B:1:MET:HE1	2:B:9:ARG:HG3	1.79	0.64
2:D:81:VAL:HG12	2:D:81:VAL:O	1.97	0.63
1:C:250:VAL:HG12	1:C:270:THR:C	2.24	0.61
2:B:1:MET:HE2	2:B:5:GLN:C	2.28	0.58
1:C:253:SER:HB2	1:C:265:ILE:CG2	2.35	0.57
1:A:270:THR:HG22	1:A:271:GLU:N	2.19	0.56
1:C:270:THR:HG22	1:C:271:GLU:N	2.20	0.56
1:A:229:ALA:C	1:A:288:MET:HE1	2.31	0.56
1:C:189:LEU:HD22	2:D:13:LEU:HD13	1.88	0.55
2:B:70:CYS:SG	2:B:75:LEU:CD1	2.89	0.54
1:A:207[B]:ARG:CZ	2:B:30:LYS:HD3	2.38	0.54
2:B:1:MET:HE1	2:B:9:ARG:CG	2.38	0.54
1:C:250:VAL:CG1	1:C:270:THR:O	2.51	0.54
1:A:276:MET:HE2	1:C:252:GLN:NE2	2.25	0.52
1:C:270:THR:CG2	1:C:271:GLU:N	2.73	0.52
2:B:75:LEU:HD12	2:B:75:LEU:N	2.26	0.51
1:C:262:LYS:C	1:C:264:GLN:H	2.19	0.50
2:D:73:VAL:CG2	6:D:163:HOH:O	2.59	0.50
2:D:60:MET:HG3	3:D:86:GOL:H11	1.95	0.49
2:B:67:HIS:HB3	6:B:136:HOH:O	2.13	0.48
2:B:70:CYS:HG	2:B:75:LEU:HD11	1.74	0.48
1:A:230:GLN:CG	1:A:288:MET:HE3	2.41	0.47
1:A:265:ILE:N	1:A:265:ILE:HD12	2.29	0.47
1:C:187:LEU:HD22	1:C:191:LYS:NZ	2.28	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:VAL:HG11	1:A:288:MET:HB3	1.96	0.47
1:C:270:THR:HG22	1:C:271:GLU:O	2.14	0.47
1:C:291:ASP:OD1	1:C:292:GLU:N	2.48	0.46
2:D:42:LYS:HG2	6:D:137:HOH:O	2.15	0.46
1:A:244:ILE:HG22	1:A:249:GLY:HA2	1.98	0.45
1:A:267:GLU:O	1:A:268:VAL:HG23	2.17	0.45
1:C:189:LEU:CD2	2:D:13:LEU:HD13	2.45	0.45
1:A:270:THR:CG2	1:A:271:GLU:N	2.81	0.43
1:A:191:LYS:HG2	1:A:194:ARG:HH22	1.83	0.43
2:B:1:MET:HE3	2:B:5:GLN:HB3	2.00	0.43
1:A:191:LYS:HG2	1:A:194:ARG:NH2	2.35	0.41
1:A:230:GLN:N	1:A:288:MET:CE	2.84	0.41
1:A:179:HIS:HA	4:A:13:NO3:O1	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/122 (100%)	118 (97%)	4 (3%)	0	100	100
1	C	115/122 (94%)	108 (94%)	6 (5%)	1 (1%)	14	14
2	B	85/85 (100%)	85 (100%)	0	0	100	100
2	D	68/85 (80%)	68 (100%)	0	0	100	100
All	All	390/414 (94%)	379 (97%)	10 (3%)	1 (0%)	36	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	263	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/106 (101%)	104 (97%)	3 (3%)	38	52
1	C	99/106 (93%)	97 (98%)	2 (2%)	48	64
2	B	78/76 (103%)	78 (100%)	0	100	100
2	D	63/76 (83%)	62 (98%)	1 (2%)	55	71
All	All	347/364 (95%)	341 (98%)	6 (2%)	51	69

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	246	LEU
1	A	260	GLU
1	A	268	VAL
1	C	250	VAL
1	C	261	ASN
2	D	77	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	HIS
1	A	230	GLN
1	A	261	ASN
2	B	51	ASN
1	C	199	GLN
1	C	230	GLN
1	C	280	GLN
2	D	41	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	5	-	5,5,5	0.39	0	5,5,5	0.29	0
4	NO3	A	13	-	1,3,3	3.36	1 (100%)	0,3,3	-	-
3	GOL	C	7	-	5,5,5	0.37	0	5,5,5	0.15	0
3	GOL	A	4	-	5,5,5	0.40	0	5,5,5	0.49	0
3	GOL	A	3	-	5,5,5	0.45	0	5,5,5	0.74	0
3	GOL	D	86	-	5,5,5	0.32	0	5,5,5	0.84	0
3	GOL	C	6	-	5,5,5	0.37	0	5,5,5	0.41	0
5	SO4	C	1	-	4,4,4	0.25	0	6,6,6	0.12	0
3	GOL	C	8	-	5,5,5	0.64	0	5,5,5	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	5	-	-	0/4/4/4	-
3	GOL	C	7	-	-	0/4/4/4	-
3	GOL	A	4	-	-	3/4/4/4	-
3	GOL	A	3	-	-	2/4/4/4	-
3	GOL	D	86	-	-	3/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	6	-	-	2/4/4/4	-
3	GOL	C	8	-	-	4/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	13	NO3	O1-N	3.36	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3	GOL	C1-C2-C3-O3
3	A	3	GOL	O2-C2-C3-O3
3	C	6	GOL	C1-C2-C3-O3
3	C	8	GOL	O1-C1-C2-C3
3	C	8	GOL	C1-C2-C3-O3
3	C	8	GOL	O2-C2-C3-O3
3	D	86	GOL	C1-C2-C3-O3
3	A	4	GOL	O1-C1-C2-C3
3	C	6	GOL	O2-C2-C3-O3
3	C	8	GOL	O1-C1-C2-O2
3	D	86	GOL	O2-C2-C3-O3
3	A	4	GOL	O1-C1-C2-O2
3	A	4	GOL	C1-C2-C3-O3
3	D	86	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	13	NO3	1	0
3	D	86	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	122/122 (100%)	0.93	19 (15%) 5 4	31, 57, 82, 102	2 (1%)
1	C	116/122 (95%)	1.31	23 (19%) 3 2	45, 69, 100, 119	1 (0%)
2	B	85/85 (100%)	0.42	1 (1%) 76 74	28, 50, 70, 76	2 (2%)
2	D	70/85 (82%)	0.68	7 (10%) 12 10	34, 57, 93, 113	1 (1%)
All	All	393/414 (94%)	0.89	50 (12%) 8 6	28, 59, 96, 119	6 (1%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	187	LEU	5.4
1	C	299	SER	5.1
1	C	256	ILE	4.9
2	D	13	LEU	4.2
1	C	265	ILE	4.1
1	C	295	GLY	3.9
1	A	249	GLY	3.7
1	A	299	SER	3.7
1	A	270	THR	3.4
1	A	285	VAL	3.4
2	D	81	VAL	3.4
2	D	34[A]	ARG	3.4
1	C	291	ASP	3.3
1	A	296	ASP	3.2
1	C	255	ARG	3.2
1	C	184	SER	3.2
1	A	269[A]	ARG	3.1
2	D	85	GLU	3.1
1	C	250	VAL	3.1
1	C	254	PHE	3.1
1	A	246	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	264	GLN	2.9
1	A	250	VAL	2.9
1	C	186	ARG	2.9
1	C	270	THR	2.7
1	C	245[A]	ARG	2.6
1	A	286	HIS	2.6
2	D	19	GLN	2.5
2	D	15	GLN	2.5
1	C	185	ARG	2.4
1	C	260	GLU	2.4
1	A	195	ASN	2.4
1	A	262	LYS	2.4
1	C	294	ASN	2.3
1	A	256	ILE	2.3
1	C	189	LEU	2.3
1	C	262	LYS	2.3
1	A	281	PHE	2.3
1	C	188	LYS	2.3
1	C	298	LYS	2.3
2	D	20	LEU	2.2
1	A	257	GLU	2.2
2	B	69	LYS	2.2
1	A	268	VAL	2.1
1	A	251	THR	2.1
1	A	244	ILE	2.1
1	C	258	GLU	2.1
1	C	268	VAL	2.1
1	A	261	ASN	2.1
1	A	241	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	C	5	6/6	0.71	0.21	91,95,98,100	0
3	GOL	C	7	6/6	0.74	0.27	97,100,104,107	0
3	GOL	C	6	6/6	0.76	0.25	74,85,88,91	0
3	GOL	A	3	6/6	0.80	0.18	58,67,71,78	0
5	SO4	C	1	5/5	0.81	0.12	94,95,102,104	0
3	GOL	A	4	6/6	0.82	0.18	65,76,81,83	0
3	GOL	C	8	6/6	0.84	0.20	58,78,83,89	0
3	GOL	D	86	6/6	0.89	0.14	61,66,69,72	0
4	NO3	A	13	4/4	0.95	0.12	52,54,55,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.