



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:49 PM UTC

PDB ID : 5VIL / pdb\_00005vil  
Title : Crystal structure of ASK1 kinase domain with a potent inhibitor (analog 6)  
Authors : Jasti, J.; Chang, J.; Kurumbail, R.  
Deposited on : 2017-04-17  
Resolution : 2.64 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2053 (2.66-2.62)
Clashscore	190562	2097 (2.66-2.62)
Ramachandran outliers	187476	2066 (2.66-2.62)
Sidechain outliers	187428	2066 (2.66-2.62)
RSRZ outliers	180081	2052 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	 3% 72% 17% • 8%
1	B	293	 3% 75% 15% • 8%
1	C	293	 3% 76% 14% • 8%
1	D	293	 7% 71% 18% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	B	1003	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8958 atoms, of which 104 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase kinase kinase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	271	Total 2146	C 1376	N 358	O 402	S 10	0	0	0
1	B	271	Total 2146	C 1376	N 358	O 402	S 10	0	0	0
1	C	271	Total 2146	C 1376	N 358	O 402	S 10	0	0	0
1	D	271	Total 2146	C 1376	N 358	O 402	S 10	0	0	0

- Molecule 2 is 2-methoxy-N-{6-[4-(propan-2-yl)-4H-1,2,4-triazol-3-yl]pyridin-2-yl}-5-sulfamoylbenzamide (CCD ID: 9E1) (formula: C<sub>18</sub>H<sub>20</sub>N<sub>6</sub>O<sub>4</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
2	A	1	Total 49	C 18	H 20	N 6	O 4	S 1	20	0
2	B	1	Total 49	C 18	H 20	N 6	O 4	S 1	20	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	C	1	Total	C	H	N	O	S	20	0
			49	18	20	6	4	1		
2	D	1	Total	C	H	N	O	S	20	0
			49	18	20	6	4	1		

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
3	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
3	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		
3	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	63	Total	O	0	0
			63	63		
4	C	9	Total	O	0	0
			9	9		

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	D	14	Total	O	0	0
			14	14		





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.54Å 132.09Å 135.22Å 90.00° 92.82° 90.00°	Depositor
Resolution (Å)	135.00 – 2.64 135.00 – 2.64	Depositor EDS
% Data completeness (in resolution range)	82.7 (135.00-2.64) 82.7 (135.00-2.64)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.65Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, $R_{free}$	0.180 , 0.213 0.189 , 0.218	Depositor DCC
$R_{free}$ test set	1773 reflections (4.23%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	71.4	Xtriage
Anisotropy	0.468	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 116.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k 0.012 for -h,-l,-k 0.038 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8958	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	122.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9E1, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.92	2/2195 (0.1%)	1.41	10/2963 (0.3%)
1	B	0.91	0/2195	1.37	9/2963 (0.3%)
1	C	0.78	0/2195	1.37	9/2963 (0.3%)
1	D	0.79	1/2195 (0.0%)	1.38	15/2963 (0.5%)
All	All	0.85	3/8780 (0.0%)	1.38	43/11852 (0.4%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	832	ILE	CA-C	5.67	1.57	1.53
1	D	832	ILE	CA-C	5.46	1.57	1.53
1	A	737	ILE	CA-CB	5.27	1.61	1.54

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	897	PHE	N-CA-C	-9.71	102.05	112.93
1	D	722	LEU	N-CA-C	-7.92	103.42	113.23
1	C	893	LYS	CA-C-N	7.72	130.88	120.77
1	C	893	LYS	C-N-CA	7.72	130.88	120.77
1	B	717	ARG	N-CA-C	-7.08	102.77	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2146	0	2136	24	0
1	B	2146	0	2136	24	0
1	C	2146	0	2136	19	0
1	D	2146	0	2136	25	0
2	A	29	20	0	0	0
2	B	29	20	0	0	0
2	C	29	20	0	1	0
2	D	29	20	0	0	0
3	A	4	6	6	0	0
3	B	12	18	18	8	0
4	A	52	0	0	0	0
4	B	63	0	0	0	0
4	C	9	0	0	0	0
4	D	14	0	0	0	0
All	All	8854	104	8568	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:715:ASP:O	1:B:716:SER:HB3	1.51	1.11
1:A:711:ILE:HD13	1:A:712:PRO:HD2	1.34	1.03
1:B:711:ILE:HD13	1:B:712:PRO:HD2	1.42	1.02
1:B:770:TRP:HE1	3:B:1002:DMS:H12	1.24	1.00
1:D:848:PRO:HB3	1:D:895:GLY:HA3	1.55	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/293 (92%)	249 (93%)	18 (7%)	2 (1%)	18	27
1	B	269/293 (92%)	254 (94%)	11 (4%)	4 (2%)	8	11
1	C	269/293 (92%)	237 (88%)	27 (10%)	5 (2%)	6	8
1	D	269/293 (92%)	241 (90%)	22 (8%)	6 (2%)	5	7
All	All	1076/1172 (92%)	981 (91%)	78 (7%)	17 (2%)	7	10

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	939	LYS
1	B	715	ASP
1	B	939	LYS
1	C	713	GLU
1	D	894	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/252 (92%)	211 (91%)	20 (9%)	9	15
1	B	231/252 (92%)	213 (92%)	18 (8%)	11	18
1	C	231/252 (92%)	214 (93%)	17 (7%)	13	20
1	D	231/252 (92%)	213 (92%)	18 (8%)	11	18
All	All	924/1008 (92%)	851 (92%)	73 (8%)	11	18

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	706	ILE
1	D	888	GLN
1	D	711	ILE
1	D	827	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	713	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	703	GLN
1	D	888	GLN
1	C	703	GLN
1	C	720	GLN
1	C	756	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DMS	A	1002	-	3,3,3	0.30	0	3,3,3	0.71	0
2	9E1	C	4000	-	31,31,31	1.77	5 (16%)	43,45,45	1.26	5 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DMS	B	1004	-	3,3,3	0.28	0	3,3,3	0.76	0
3	DMS	B	1002	-	3,3,3	0.23	0	3,3,3	0.60	0
2	9E1	D	4000	-	31,31,31	1.78	5 (16%)	43,45,45	1.29	5 (11%)
2	9E1	B	1001	-	31,31,31	1.75	5 (16%)	43,45,45	1.34	5 (11%)
2	9E1	A	1001	-	31,31,31	1.77	5 (16%)	43,45,45	1.28	5 (11%)
3	DMS	B	1003	-	3,3,3	0.92	0	3,3,3	1.48	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9E1	B	1001	-	-	4/24/24/24	0/3/3/3
2	9E1	A	1001	-	-	4/24/24/24	0/3/3/3
2	9E1	C	4000	-	-	4/24/24/24	0/3/3/3
2	9E1	D	4000	-	-	6/24/24/24	0/3/3/3

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	9E1	C14-S1	-6.88	1.66	1.77
2	B	1001	9E1	C14-S1	-6.80	1.66	1.77
2	D	4000	9E1	C14-S1	-6.77	1.66	1.77
2	C	4000	9E1	C14-S1	-6.74	1.66	1.77
2	D	4000	9E1	C5-N3	3.47	1.37	1.32

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4000	9E1	N1-C5-N3	-4.33	108.29	110.31
2	B	1001	9E1	N1-C5-N3	-4.29	108.31	110.31
2	A	1001	9E1	N1-C5-N3	-4.17	108.36	110.31
2	C	4000	9E1	N1-C5-N3	-4.10	108.39	110.31
2	B	1001	9E1	C14-S1-N6	-3.47	103.57	108.40

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1001	9E1	C15-C14-S1-O3
2	C	4000	9E1	C15-C14-S1-O3
2	A	1001	9E1	C13-C14-S1-N6
2	B	1001	9E1	C13-C14-S1-N6
2	C	4000	9E1	C13-C14-S1-O3


There are no ring outliers.


3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4000	9E1	1	0
3	B	1002	DMS	3	0
3	B	1003	DMS	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/293 (92%)	-0.21	9 (3%) 49 43	46, 84, 143, 197	0
1	B	271/293 (92%)	-0.35	9 (3%) 49 43	46, 78, 136, 173	0
1	C	271/293 (92%)	0.34	10 (3%) 45 39	93, 165, 230, 243	0
1	D	271/293 (92%)	0.42	20 (7%) 20 17	96, 161, 208, 229	0
All	All	1084/1172 (92%)	0.05	48 (4%) 39 32	46, 117, 207, 243	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	894	VAL	4.6
1	B	836	THR	4.1
1	A	940	VAL	3.8
1	C	670	LEU	3.7
1	C	889	ALA	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	DMS	B	1002	4/4	0.92	0.12	98,100,103,104	0
3	DMS	B	1003	4/4	0.92	0.14	63,75,77,84	0
2	9E1	C	4000	29/29	0.95	0.08	102,123,139,143	20
3	DMS	A	1002	4/4	0.95	0.19	78,86,88,90	0
2	9E1	D	4000	29/29	0.96	0.07	104,114,131,135	20
3	DMS	B	1004	4/4	0.96	0.15	80,86,89,90	0
2	9E1	A	1001	29/29	0.98	0.05	50,58,63,66	20
2	9E1	B	1001	29/29	0.98	0.06	35,47,55,59	20

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.