



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:28 AM UTC

PDB ID : 5VIU / pdb_00005viu
Title : Crystal Structure of Acetylornithine Aminotransferase from Elizabethkingia anophelis
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2017-04-17
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	5% 90% 5% .
1	B	419	3% 86% 5% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CIT	A	509	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylornithine aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	15	0
			3113	1997	521	585	10			
1	B	383	Total	C	N	O	S	0	14	0
			2953	1897	491	554	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A0A077E919
A	-6	ALA	-	expression tag	UNP A0A077E919
A	-5	HIS	-	expression tag	UNP A0A077E919
A	-4	HIS	-	expression tag	UNP A0A077E919
A	-3	HIS	-	expression tag	UNP A0A077E919
A	-2	HIS	-	expression tag	UNP A0A077E919
A	-1	HIS	-	expression tag	UNP A0A077E919
A	0	HIS	-	expression tag	UNP A0A077E919
B	-7	MET	-	expression tag	UNP A0A077E919
B	-6	ALA	-	expression tag	UNP A0A077E919
B	-5	HIS	-	expression tag	UNP A0A077E919
B	-4	HIS	-	expression tag	UNP A0A077E919
B	-3	HIS	-	expression tag	UNP A0A077E919
B	-2	HIS	-	expression tag	UNP A0A077E919
B	-1	HIS	-	expression tag	UNP A0A077E919
B	0	HIS	-	expression tag	UNP A0A077E919

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	399	Total	O	0	8
			404	404		
4	B	314	Total	O	0	9
			319	319		

- Molecule 1: Acetylornithine aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	136.60Å 77.47Å 104.96Å 90.00° 129.46° 90.00°	Depositor
Resolution (Å)	42.35 – 1.65 42.35 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (42.35-1.65) 99.8 (42.35-1.65)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 1.65Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.155 , 0.180 0.157 , 0.180	Depositor DCC
R_{free} test set	1923 reflections (1.89%)	wwPDB-VP
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6856	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3215	0.48	0/4355
1	B	0.23	0/3048	0.43	0/4135
All	All	0.24	0/6263	0.46	0/8490

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3113	0	3150	19	0
1	B	2953	0	2981	18	4
2	A	24	0	36	2	0
2	B	4	0	6	0	0
3	A	39	0	15	3	0
4	A	404	0	0	5	0
4	B	319	0	0	4	0
All	All	6856	0	6188	34	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:GLN:NE2	4:A:602[A]:HOH:O	2.23	0.70
1:A:179:ILE:HG22	2:A:506:EDO:H11	1.74	0.68
1:B:68:LEU:HD12	1:B:297:LEU:HD11	1.73	0.68
1:A:73[B]:SER:OG	4:A:601:HOH:O	2.12	0.66
1:A:68:LEU:HD12	1:A:297:LEU:HD11	1.78	0.66
1:B:345:LYS:NZ	4:B:604:HOH:O	2.29	0.63
1:B:106[B]:MET:HE1	1:B:273:VAL:HG23	1.81	0.63
1:A:24[B]:PRO:HG2	1:B:83:TYR:CE1	2.40	0.57
2:A:502:EDO:H21	1:B:76:ALA:HB3	1.89	0.55
1:A:73[B]:SER:HB3	4:A:618[B]:HOH:O	2.06	0.54
1:B:138:ILE:HD11	1:B:193:ILE:HD13	1.89	0.53
1:A:291:THR:HG21	4:B:612:HOH:O	2.08	0.53
1:B:49:LEU:HG	1:B:52:TYR:H	1.74	0.53
1:A:155:SER:O	1:B:121[A]:ARG:NH1	2.43	0.52
1:B:106[A]:MET:HE3	1:B:111[A]:GLU:HB3	1.93	0.51
1:A:223:LYS:NZ	4:A:611:HOH:O	2.43	0.50
1:A:138:ILE:HD11	1:A:193[B]:ILE:HD13	1.93	0.50
1:A:269:PRO:HD3	1:B:268:TYR:CZ	2.48	0.49
3:A:509:CIT:H42	4:A:726:HOH:O	2.13	0.47
1:B:223:LYS:HD3	4:B:615:HOH:O	2.14	0.46
1:B:235:GLN:HG3	1:B:261:LYS:HE3	1.98	0.45
1:A:69:VAL:HG22	1:B:69[B]:VAL:HG12	1.99	0.45
1:B:261:LYS:NZ	4:B:612:HOH:O	2.48	0.45
1:A:124:SER:HB2	1:A:130:ILE:HD12	1.99	0.44
1:A:69:VAL:HG22	1:B:69[A]:VAL:HG22	1.99	0.44
1:A:95:THR:HB	1:A:100:PHE:O	2.18	0.44
1:A:136:LYS:HB2	1:A:193[A]:ILE:HD13	1.99	0.44
3:A:509:CIT:O2	3:A:509:CIT:O7	2.36	0.43
1:B:106[B]:MET:HE1	1:B:273:VAL:CG2	2.49	0.43
1:A:292:PHE:HD1	3:A:507:CIT:H41	1.85	0.42
1:B:324:LEU:HD23	1:B:397:LEU:HD23	2.00	0.42
1:B:238:ILE:HG22	1:B:309:VAL:HG21	2.03	0.41
1:A:49:LEU:HD23	1:A:52:TYR:H	1.85	0.41
1:A:36:PHE:CD1	1:A:44:LYS:HE2	2.57	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:GLU:CD	1:B:66:GLU:OE2[2_555]	1.35	0.85
1:B:66:GLU:OE2	1:B:66:GLU:OE2[2_555]	1.55	0.65
1:B:66:GLU:OE1	1:B:66:GLU:OE2[2_555]	1.80	0.40
1:B:66:GLU:CD	1:B:66:GLU:CD[2_555]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/419 (98%)	399 (97%)	9 (2%)	2 (0%)	24	10
1	B	393/419 (94%)	381 (97%)	11 (3%)	1 (0%)	36	21
All	All	803/838 (96%)	780 (97%)	20 (2%)	3 (0%)	36	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261[A]	LYS
1	A	261[B]	LYS
1	B	261	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/350 (95%)	332 (100%)	1 (0%)	86	80
1	B	313/350 (89%)	311 (99%)	2 (1%)	78	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	646/700 (92%)	643 (100%)	3 (0%)	81	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	LEU
1	B	89	GLU
1	B	235	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	B	107	ASN
1	B	225	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	501	-	3,3,3	0.41	0	2,2,2	0.48	0
2	EDO	A	502	-	3,3,3	0.55	0	2,2,2	0.35	0
2	EDO	A	503	-	3,3,3	0.44	0	2,2,2	0.43	0
2	EDO	A	506	-	3,3,3	0.42	0	2,2,2	0.39	0
2	EDO	B	501	-	3,3,3	0.46	0	2,2,2	0.40	0
3	CIT	A	507	-	12,12,12	1.06	0	17,17,17	1.53	2 (11%)
2	EDO	A	505	-	3,3,3	0.37	0	2,2,2	0.44	0
3	CIT	A	508	-	12,12,12	1.17	0	17,17,17	1.73	5 (29%)
3	CIT	A	509	-	12,12,12	1.10	0	17,17,17	1.94	6 (35%)
2	EDO	A	504	-	3,3,3	0.41	0	2,2,2	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	501	-	-	0/1/1/1	-
2	EDO	A	502	-	-	0/1/1/1	-
2	EDO	A	503	-	-	0/1/1/1	-
2	EDO	A	506	-	-	1/1/1/1	-
2	EDO	B	501	-	-	0/1/1/1	-
3	CIT	A	507	-	-	6/16/16/16	-
2	EDO	A	505	-	-	0/1/1/1	-
3	CIT	A	508	-	-	3/16/16/16	-
3	CIT	A	509	-	-	12/16/16/16	-
2	EDO	A	504	-	-	0/1/1/1	-

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	509	CIT	O6-C6-C3	3.52	119.90	113.14
3	A	507	CIT	O6-C6-C3	3.51	119.88	113.14
3	A	508	CIT	O6-C6-C3	3.38	119.63	113.14
3	A	509	CIT	O4-C5-C4	3.22	124.55	114.35
3	A	508	CIT	C2-C3-C6	-2.74	103.98	110.03
3	A	509	CIT	C4-C3-C6	2.55	115.68	110.03
3	A	509	CIT	C3-C4-C5	2.46	120.64	113.92
3	A	508	CIT	O2-C1-C2	2.41	121.97	114.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	509	CIT	O4-C5-O3	-2.40	117.16	123.33
3	A	507	CIT	O2-C1-O1	-2.34	117.31	123.33
3	A	509	CIT	O7-C3-C6	-2.21	105.83	108.96
3	A	508	CIT	O4-C5-C4	2.17	121.24	114.35
3	A	508	CIT	O4-C5-O3	-2.12	117.87	123.33

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	509	CIT	C1-C2-C3-C4
3	A	509	CIT	C1-C2-C3-C6
3	A	509	CIT	C2-C3-C4-C5
3	A	509	CIT	O7-C3-C4-C5
3	A	509	CIT	C2-C3-C6-O5
3	A	509	CIT	O7-C3-C6-O5
3	A	509	CIT	O7-C3-C6-O6
3	A	509	CIT	C1-C2-C3-O7
3	A	509	CIT	C2-C3-C6-O6
3	A	509	CIT	C6-C3-C4-C5
2	A	506	EDO	O1-C1-C2-O2
3	A	507	CIT	O7-C3-C6-O5
3	A	507	CIT	C2-C3-C6-O5
3	A	507	CIT	C2-C3-C6-O6
3	A	507	CIT	C4-C3-C6-O5
3	A	507	CIT	C4-C3-C6-O6
3	A	509	CIT	C3-C4-C5-O4
3	A	508	CIT	C2-C3-C6-O6
3	A	508	CIT	O2-C1-C2-C3
3	A	507	CIT	O7-C3-C6-O6
3	A	509	CIT	C3-C4-C5-O3
3	A	508	CIT	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	EDO	1	0
2	A	506	EDO	1	0
3	A	507	CIT	1	0
3	A	509	CIT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/419 (95%)	-0.14	20 (4%) 34 38	11, 22, 56, 90	15 (3%)
1	B	383/419 (91%)	-0.04	12 (3%) 51 55	13, 27, 53, 86	14 (3%)
All	All	784/838 (93%)	-0.09	32 (4%) 41 45	11, 24, 55, 90	29 (3%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	205	ALA	6.0
1	A	151	ILE	5.4
1	A	154	PHE	5.1
1	A	4	THR	4.1
1	A	149	THR	4.1
1	B	133	ASN	3.9
1	A	148	THR	3.8
1	A	150	THR	3.7
1	A	144	PHE	3.5
1	B	206	GLY	3.5
1	A	125	TYR	3.2
1	A	146	GLY	3.2
1	A	133	ASN	3.1
1	B	168	THR	3.1
1	A	287	GLN	3.1
1	A	145	HIS	3.0
1	B	142	ASN	3.0
1	A	167	PHE	3.0
1	B	289	GLY	3.0
1	A	147	ARG	2.9
1	A	155	SER	2.9
1	A	288	HIS	2.9
1	B	4	THR	2.7
1	B	290	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	287	GLN	2.6
1	B	131	ALA	2.6
1	A	165	GLY	2.6
1	A	166	PRO	2.4
1	B	380	HIS	2.1
1	B	204	GLU	2.0
1	A	159	ASP	2.0
1	A	131	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CIT	A	509	13/13	0.82	0.17	30,66,81,84	0
2	EDO	A	505	4/4	0.85	0.19	48,49,50,54	0
2	EDO	A	506	4/4	0.85	0.17	53,62,62,67	0
2	EDO	A	503	4/4	0.85	0.16	39,51,56,63	0
3	CIT	A	508	13/13	0.89	0.12	19,33,39,42	13
2	EDO	B	501	4/4	0.90	0.12	33,40,42,52	0
2	EDO	A	501	4/4	0.91	0.13	37,40,42,49	0
3	CIT	A	507	13/13	0.91	0.11	23,27,36,38	13
2	EDO	A	504	4/4	0.92	0.10	40,48,48,50	0
2	EDO	A	502	4/4	0.94	0.13	19,20,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.