



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:16 AM UTC

PDB ID : 2VJL / pdb_00002vjL
Title : Formyl-CoA transferase with aspartyl-CoA thioester intermediate derived from formyl-CoA
Authors : Berthold, C.L.; Toyota, C.G.; Richards, N.G.J.; Lindqvist, Y.
Deposited on : 2007-12-11
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	7% 90% 10%
1	B	428	12% 94% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FORMYL-COENZYME A TRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	6	0
			3339	2114	571	631	23			
1	B	427	Total	C	N	O	S	259	2	0
			3323	2103	570	627	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	ILE	MET	conflict	UNP O06644
B	186	ILE	MET	conflict	UNP O06644

- Molecule 2 is COENZYME A (CCD ID: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	2	Total	Cl	0	0
			2	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	369	Total	O	0	0
			369	369		
5	B	325	Total	O	0	0
			325	325		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FORMYL-COENZYME A TRANSFERASE



• Molecule 1: FORMYL-COENZYME A TRANSFERASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	151.76Å 151.76Å 100.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (30.00-2.00) 98.2 (30.00-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.197 , 0.242 0.199 , 0.242	Depositor DCC
R_{free} test set	3765 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7457	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/3431	0.75	0/4638
1	B	0.62	2/3403 (0.1%)	0.73	0/4602
All	All	0.57	2/6834 (0.0%)	0.74	0/9240

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	316	LYS	C-N	-15.45	1.12	1.33
1	B	285	ALA	C-N	12.72	1.50	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	285	ALA	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3339	0	3291	31	0
1	B	3323	0	3265	12	0
2	A	48	0	31	1	0
2	B	48	0	31	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	2	0	0	0	0
5	A	369	0	0	3	0
5	B	325	0	0	2	0
All	All	7457	0	6618	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:ILE:HG21	1:A:408:VAL:HG21	1.45	0.96
1:B:126:GLY:HA3	1:B:138:VAL:HG13	1.70	0.73
1:A:11:LEU:HD21	1:A:36:ILE:HD11	1.72	0.71
1:A:319:ASP:OD1	5:A:2270:HOH:O	2.09	0.69
1:B:102:LEU:HD23	1:B:105:MET:HE3	1.78	0.63
1:A:422:LEU:HD22	1:A:427:VAL:HG11	1.83	0.61
1:A:219:GLU:OE2	1:B:358:HIS:HE1	1.83	0.61
1:B:234:PHE:CZ	1:B:318:MET:HE2	2.35	0.61
1:A:208:VAL:HG12	1:A:208:VAL:O	2.02	0.59
1:A:124:VAL:HG23	1:A:176:LEU:HD22	1.87	0.56
1:B:17:GLN:HG3	1:B:63:PHE:CE2	2.42	0.54
1:A:70:ILE:HG21	1:A:408:VAL:CG2	2.30	0.54
1:A:422:LEU:CD2	1:A:427:VAL:HG11	2.39	0.52
1:B:16:VAL:HA	1:B:45:THR:HG23	1.92	0.51
1:B:136:LEU:O	5:B:2138:HOH:O	2.19	0.51
1:A:209:ARG:HD2	1:B:63:PHE:CZ	2.47	0.50
1:A:74:MET:HE3	1:A:105:MET:HE1	1.93	0.49
1:A:83:LEU:HD23	1:A:86:MET:HE3	1.96	0.48
1:A:70:ILE:HG22	1:A:405:THR:HA	1.96	0.48
1:A:17:GLN:NE2	1:A:59:TYR:OH	2.47	0.48
1:A:11:LEU:HD21	1:A:36:ILE:CD1	2.44	0.47
1:A:186:ILE:HD13	1:A:194:GLN:NE2	2.30	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ASN:HB3	1:A:136:LEU:HD12	1.97	0.46
1:A:95:GLU:HG2	1:A:102:LEU:HD22	1.97	0.46
1:B:392:GLN:NE2	5:B:2303:HOH:O	2.48	0.46
1:A:288:MET:HE1	1:A:344:GLY:C	2.42	0.45
1:B:102:LEU:HD23	1:B:105:MET:CE	2.44	0.45
1:B:133:ASN:HB3	1:B:136:LEU:HD12	1.98	0.44
1:A:298:LYS:NZ	5:A:2247:HOH:O	2.51	0.43
1:A:262:GLN:NE2	5:A:2219:HOH:O	2.51	0.43
1:A:14:THR:HB	1:A:19:GLY:HA3	2.01	0.43
1:A:70:ILE:CG2	1:A:408:VAL:HG21	2.33	0.43
1:B:186:ILE:HG21	1:B:194:GLN:NE2	2.34	0.43
1:A:404:HIS:O	1:A:408:VAL:HG13	2.19	0.42
1:A:36:ILE:HG12	1:A:70:ILE:HD11	2.02	0.42
1:A:208:VAL:HG12	1:A:211:LYS:HB2	2.01	0.42
1:A:137:LYS:HB3	2:A:1169:COA:H142	2.02	0.41
1:A:16:VAL:HG23	1:A:17:GLN:H	1.86	0.41
1:A:179:GLY:HA3	1:A:196:VAL:HG21	2.02	0.41
1:A:15:HIS:O	1:A:16:VAL:HG22	2.21	0.41
1:A:288:MET:HE2	1:A:288:MET:HB3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/428 (101%)	420 (97%)	10 (2%)	1 (0%)	43	42
1	B	427/428 (100%)	411 (96%)	12 (3%)	4 (1%)	14	9
All	All	858/856 (100%)	831 (97%)	22 (3%)	5 (1%)	21	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	140	GLU
1	B	134	GLU
1	B	233	ASN
1	A	16	VAL
1	B	16	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/351 (101%)	350 (98%)	6 (2%)	53	60
1	B	352/351 (100%)	349 (99%)	3 (1%)	70	78
All	All	708/702 (101%)	699 (99%)	9 (1%)	61	68

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LEU
1	A	35	LYS
1	A	38	ARG
1	A	44	MET
1	A	78	GLU
1	A	386	PHE
1	B	138	VAL
1	B	288	MET
1	B	386	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	342	GLN
1	B	113	GLN
1	B	233	ASN
1	B	358	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COA	A	1169	1	47,50,50	1.22	4 (8%)	69,75,75	1.50	9 (13%)
2	COA	B	1169	1	47,50,50	1.23	4 (8%)	69,75,75	1.46	7 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	1169	1	-	10/48/64/64	0/3/3/3
2	COA	B	1169	1	-	4/48/64/64	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1169	COA	O9P-C9P	5.33	1.33	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1169	COA	O9P-C9P	5.26	1.33	1.23
2	B	1169	COA	P2A-O3A	2.85	1.62	1.59
2	B	1169	COA	P1A-O3A	2.67	1.62	1.59
2	A	1169	COA	P2A-O3A	2.43	1.62	1.59
2	A	1169	COA	P1A-O3A	2.41	1.62	1.59
2	B	1169	COA	C5A-N7A	-2.33	1.34	1.39
2	A	1169	COA	C8A-N7A	2.23	1.36	1.31

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1169	COA	C5A-C4A-N3A	-5.45	119.21	126.72
2	A	1169	COA	C5A-C4A-N3A	-5.24	119.50	126.72
2	A	1169	COA	N3A-C2A-N1A	-4.93	121.12	128.58
2	B	1169	COA	N3A-C2A-N1A	-4.71	121.45	128.58
2	A	1169	COA	N9A-C8A-N7A	-3.57	108.88	113.94
2	A	1169	COA	C2A-N3A-C4A	3.52	120.44	111.83
2	B	1169	COA	C2A-N3A-C4A	3.47	120.30	111.83
2	A	1169	COA	N3A-C4A-N9A	3.33	132.84	127.17
2	B	1169	COA	N3A-C4A-N9A	3.31	132.80	127.17
2	B	1169	COA	N9A-C8A-N7A	-2.99	109.69	113.94
2	A	1169	COA	C4A-C5A-N7A	-2.93	107.23	110.58
2	B	1169	COA	C4A-C5A-N7A	-2.86	107.31	110.58
2	A	1169	COA	C5A-N7A-C8A	2.71	107.71	103.45
2	A	1169	COA	C4A-N9A-C8A	2.66	108.53	105.74
2	B	1169	COA	C5A-N7A-C8A	2.61	107.55	103.45
2	A	1169	COA	O4B-C1B-N9A	2.31	112.53	108.09

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1169	COA	CCP-O6A-P2A-O3A
2	A	1169	COA	CCP-O6A-P2A-O4A
2	A	1169	COA	CCP-O6A-P2A-O5A
2	B	1169	COA	C5B-O5B-P1A-O1A
2	B	1169	COA	C5B-O5B-P1A-O3A
2	A	1169	COA	O5P-C5P-N4P-C3P
2	A	1169	COA	C6P-C5P-N4P-C3P
2	B	1169	COA	C5B-O5B-P1A-O2A
2	A	1169	COA	C3B-O3B-P3B-O8A
2	A	1169	COA	C4B-C3B-O3B-P3B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1169	COA	P1A-O3A-P2A-O4A
2	A	1169	COA	P1A-O3A-P2A-O5A
2	A	1169	COA	C2B-C3B-O3B-P3B
2	B	1169	COA	C3B-O3B-P3B-O8A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1169	COA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	316:LYS	C	317:LEU	N	1.12

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	427/428 (99%)	0.31	31 (7%) 21 19	13, 27, 61, 71	6 (1%)
1	B	396/428 (92%)	0.57	53 (13%) 7 6	15, 30, 64, 73	2 (0%)
All	All	823/856 (96%)	0.43	84 (10%) 12 11	13, 28, 62, 73	8 (0%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	321	PHE	5.8
1	B	320	ILE	5.6
1	B	323	PHE	5.0
1	B	284	ILE	4.4
1	B	236	PHE	4.1
1	B	343	TYR	4.0
1	B	420	LYS	4.0
1	B	285	ALA	3.9
1	A	427	VAL	3.8
1	B	240	GLY	3.8
1	B	421	GLU	3.8
1	B	413	GLY	3.7
1	B	424	ALA	3.7
1	B	247	ASN	3.6
1	B	322	SER	3.5
1	B	261	GLY	3.5
1	A	408	VAL	3.4
1	A	54[A]	ASN	3.3
1	A	419	ILE	3.3
1	A	99	PRO	3.3
1	A	417	ALA	3.2
1	B	233	ASN	3.2
1	B	422	LEU	3.2
1	A	247	ASN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	238	ARG	3.1
1	B	234	PHE	3.0
1	B	6	ASP	3.0
1	A	109	TRP	3.0
1	B	326	THR	3.0
1	B	417	ALA	3.0
1	B	344	GLY	3.0
1	B	317	LEU	3.0
1	B	239	ASP	3.0
1	A	139	TYR	3.0
1	B	339	TRP	3.0
1	B	235	ALA	2.9
1	B	260	GLY	2.9
1	B	410	LYS	2.9
1	B	282	PHE	2.8
1	B	414	LEU	2.8
1	A	415	ASP	2.8
1	A	40	GLY	2.8
1	B	2	THR	2.8
1	B	345	ILE	2.8
1	A	418	LYS	2.7
1	B	346	PRO	2.7
1	A	81	GLU	2.6
1	B	342	GLN	2.6
1	A	2	THR	2.6
1	B	324	ILE	2.6
1	A	424	ALA	2.6
1	A	414	LEU	2.5
1	B	419	ILE	2.5
1	A	78	GLU	2.5
1	A	48	TRP	2.5
1	A	106	GLY	2.5
1	A	275	ASP	2.4
1	B	105	MET	2.3
1	B	106	GLY	2.3
1	B	412	LEU	2.3
1	A	83	LEU	2.3
1	B	119	VAL	2.3
1	B	111	TYR	2.3
1	B	76	THR	2.2
1	B	241	ASN	2.2
1	A	44	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	GLY	2.2
1	A	111	TYR	2.2
1	B	243	LEU	2.2
1	B	135	HIS	2.2
1	A	101	ALA	2.2
1	B	340	ALA	2.2
1	B	318	MET	2.2
1	A	421	GLU	2.1
1	A	77	PRO	2.1
1	A	97	PHE	2.1
1	B	409	LEU	2.1
1	B	392	GLN	2.1
1	A	186	ILE	2.1
1	A	409	LEU	2.1
1	A	105	MET	2.1
1	B	418	LYS	2.0
1	B	110	GLU	2.0
1	B	338	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	COA	A	1169	48/48	0.86	0.13	41,58,65,65	0
4	MG	A	4000	1/1	0.92	0.06	40,40,40,40	0
2	COA	B	1169	48/48	0.93	0.10	23,39,45,46	0
3	CL	A	3000	1/1	0.97	0.10	35,35,35,35	0
3	CL	B	3001	1/1	0.99	0.11	28,28,28,28	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	B	3000	1/1	0.99	0.07	26,26,26,26	0
4	MG	A	4001	1/1	0.99	0.08	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around COA A 1169:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around COA B 1169:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.