



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:24 AM UTC

PDB ID : 5VLP / pdb_00005vlp
Title : PCSK9 complex with LDLR antagonist peptide and Fab7G7
Authors : Eigenbrot, C.; Ultsch, M.
Deposited on : 2017-04-25
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	700	
2	H	223	
3	L	214	
4	Z	17	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	0	0
			4332	2688	802	808	34			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	474	ILE	VAL	variant	UNP Q8NBP7
A	670	GLU	GLY	variant	UNP Q8NBP7
A	693	HIS	-	expression tag	UNP Q8NBP7
A	694	HIS	-	expression tag	UNP Q8NBP7
A	695	HIS	-	expression tag	UNP Q8NBP7
A	696	HIS	-	expression tag	UNP Q8NBP7
A	697	HIS	-	expression tag	UNP Q8NBP7
A	698	HIS	-	expression tag	UNP Q8NBP7
A	699	HIS	-	expression tag	UNP Q8NBP7
A	700	HIS	-	expression tag	UNP Q8NBP7

- Molecule 2 is a protein called Fab7G7 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	0	0
			1677	1066	269	335	7			

- Molecule 3 is a protein called Fab7G7 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	214	Total	C	N	O	S	0	0	0
			1661	1035	278	340	8			

- Molecule 4 is a protein called LDLR antagonist peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	Z	17	Total	C	N	O	S	0	0	0
			136	90	24	21	1			

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	O	0	0
			3	3		
6	H	1	Total	O	0	0
			1	1		
6	L	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proprotein convertase subtilisin/kexin type 9




- Molecule 2: Fab7G7 heavy chain




- Molecule 3: Fab7G7 light chain





● Molecule 4: LDLR antagonist peptide



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	110.86Å 142.17Å 239.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.60 – 2.90 70.60 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.60-2.90) 99.9 (70.60-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.171 , 0.209 0.180 , 0.216	Depositor DCC
R_{free} test set	2164 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7814	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, HRG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/4415	1.32	19/5988 (0.3%)
2	H	0.76	0/1723	1.19	7/2353 (0.3%)
3	L	0.73	0/1698	1.20	7/2301 (0.3%)
4	Z	0.61	0/124	0.99	0/166
All	All	0.78	0/7960	1.26	33/10808 (0.3%)

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	356	GLY	CA-C-N	8.64	133.78	120.82
1	A	356	GLY	C-N-CA	8.64	133.78	120.82
1	A	224	ASP	CA-CB-CG	8.34	120.94	112.60
1	A	547	GLY	CA-C-N	8.29	132.06	120.29
1	A	547	GLY	C-N-CA	8.29	132.06	120.29
1	A	252	VAL	N-CA-CB	-6.29	104.85	112.33
1	A	129	ASP	CA-CB-CG	6.17	118.77	112.60
1	A	223	CYS	CA-C-N	6.00	132.98	122.79
1	A	223	CYS	C-N-CA	6.00	132.98	122.79
2	H	140	GLY	N-CA-C	5.81	118.82	111.09
2	H	138	GLY	CA-C-N	5.71	129.82	121.31
2	H	138	GLY	C-N-CA	5.71	129.82	121.31
3	L	25	THR	CA-C-N	5.57	127.75	120.28
3	L	25	THR	C-N-CA	5.57	127.75	120.28
1	A	437	THR	CB-CA-C	5.49	116.11	109.85
3	L	151	ASP	CA-CB-CG	5.47	118.07	112.60
1	A	643	HIS	CA-C-N	5.39	129.32	122.37
1	A	643	HIS	C-N-CA	5.39	129.32	122.37
1	A	517	GLY	CA-C-N	5.37	127.74	120.38
1	A	517	GLY	C-N-CA	5.37	127.74	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	158	PRO	CA-C-N	5.37	131.80	121.54
2	H	158	PRO	C-N-CA	5.37	131.80	121.54
1	A	252	VAL	CB-CA-C	5.29	116.33	110.62
1	A	418	PHE	CA-CB-CG	-5.28	108.52	113.80
1	A	131	LEU	CA-C-N	5.20	127.51	120.38
1	A	131	LEU	C-N-CA	5.20	127.51	120.38
2	H	167	SER	N-CA-C	-5.16	108.73	114.62
1	A	589	VAL	N-CA-CB	-5.15	105.11	110.72
3	L	137	ASN	N-CA-C	5.10	118.12	109.76
3	L	31	THR	CB-CA-C	5.10	118.52	109.75
3	L	211	ARG	CA-C-N	5.04	129.13	120.72
3	L	211	ARG	C-N-CA	5.04	129.13	120.72
2	H	131	SER	N-CA-C	-5.03	101.04	109.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4332	0	4263	35	0
2	H	1677	0	1612	17	0
3	L	1661	0	1580	13	0
4	Z	136	0	140	3	0
5	A	2	0	0	0	0
6	A	3	0	0	0	0
6	H	1	0	0	0	0
6	L	2	0	0	0	0
All	All	7814	0	7595	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:586:ASN:H	1:A:586:ASN:HD22	1.32	0.78
1:A:272:ARG:HE	1:A:275:GLN:HE21	1.38	0.71
1:A:560:THR:HB	1:A:633:THR:HG21	1.72	0.71
2:H:23:LYS:HA	2:H:78:ASN:HB3	1.74	0.68
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.76	0.67
2:H:33:TYR:CE2	2:H:52:TYR:HB2	2.36	0.60
1:A:490:SER:HB2	1:A:520:VAL:HG12	1.83	0.60
1:A:346:VAL:HG22	1:A:366:GLU:HB2	1.84	0.60
1:A:399:MET:HE1	1:A:415:LEU:HG	1.84	0.60
2:H:174:VAL:HG22	2:H:192:VAL:HG23	1.84	0.59
1:A:66:ARG:HH11	1:A:73:ARG:HD2	1.67	0.59
3:L:3:VAL:H	3:L:26:SER:HB3	1.67	0.58
2:H:35:ASN:HD22	2:H:47:TRP:HE1	1.50	0.58
1:A:66:ARG:HD3	1:A:73:ARG:CZ	2.34	0.57
3:L:2:ILE:HG12	3:L:27:GLN:HG2	1.87	0.56
2:H:101:PHE:CD1	2:H:109:TRP:HB3	2.40	0.56
1:A:615:ILE:HD12	1:A:622:VAL:HG13	1.87	0.55
3:L:146:VAL:HG21	3:L:175:MET:HE1	1.88	0.55
1:A:560:THR:HB	1:A:633:THR:CG2	2.35	0.55
1:A:586:ASN:HD22	1:A:586:ASN:N	2.02	0.55
1:A:99:GLN:HG3	1:A:109:THR:HG22	1.89	0.55
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.88	0.54
2:H:35:ASN:HD22	2:H:47:TRP:NE1	2.06	0.53
3:L:195:GLU:HG2	3:L:206:VAL:HG22	1.90	0.53
2:H:18:VAL:HG22	2:H:86:LEU:HD11	1.92	0.52
1:A:88:LEU:HD13	1:A:116:HIS:HB3	1.92	0.51
3:L:149:LYS:HB2	3:L:193:THR:HB	1.92	0.51
1:A:476:ARG:HH21	1:A:503:GLN:HE22	1.58	0.50
1:A:364:PRO:HB3	4:Z:1:TRP:CH2	2.46	0.50
1:A:407:THR:HG23	1:A:410:GLU:H	1.77	0.49
1:A:369:ILE:HG13	4:Z:-1:PRO:HB3	1.95	0.49
1:A:332:GLU:H	1:A:332:GLU:CD	2.21	0.48
1:A:69:LYS:HG3	1:A:71:PRO:HD2	1.95	0.48
1:A:484:LEU:HD22	1:A:527:CYS:HB2	1.95	0.48
2:H:30:THR:HA	2:H:53:PRO:HB2	1.95	0.48
3:L:117:ILE:HD12	3:L:194:CYS:HB3	1.95	0.48
3:L:163:TRP:CD2	3:L:175:MET:HG3	2.49	0.47
3:L:3:VAL:H	3:L:26:SER:CB	2.27	0.47
2:H:130:PRO:HB3	2:H:156:TYR:HB3	1.96	0.47
1:A:548:THR:HG21	1:A:581:PRO:HG2	1.96	0.47
3:L:122:SER:O	3:L:126:THR:HG23	2.15	0.46
4:Z:4:VAL:HG12	4:Z:9:LEU:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:ALA:HB1	1:A:365:GLY:HA3	1.97	0.46
2:H:33:TYR:CZ	2:H:52:TYR:HB2	2.50	0.46
1:A:529:LEU:HD21	1:A:606:LEU:HD22	1.97	0.46
1:A:586:ASN:H	1:A:586:ASN:ND2	2.05	0.46
1:A:426:GLU:HB3	1:A:434:ARG:HG2	1.97	0.45
1:A:180:VAL:HG11	1:A:397:ALA:HA	1.99	0.44
1:A:186:ASP:OD2	1:A:288:PRO:HG2	2.16	0.44
2:H:27:TYR:OH	2:H:34:ILE:HD11	2.18	0.44
2:H:114:TRP:HB2	3:L:43:SER:HB2	1.99	0.44
2:H:35:ASN:ND2	2:H:47:TRP:HE1	2.14	0.44
3:L:35:TRP:CE2	3:L:73:PHE:HB2	2.54	0.43
1:A:407:THR:HG22	1:A:410:GLU:CG	2.48	0.43
1:A:453:TRP:HH2	1:A:604:PRO:HG2	1.83	0.42
1:A:82:LEU:HD13	1:A:86:THR:HG21	2.02	0.42
3:L:124:GLN:HE22	3:L:131:SER:H	1.69	0.41
1:A:560:THR:O	1:A:633:THR:HG22	2.20	0.41
2:H:163:VAL:HG22	2:H:208:VAL:HG22	2.03	0.41
1:A:276:LEU:HD12	1:A:276:LEU:HA	1.87	0.41
2:H:149:LEU:HB3	2:H:221:LEU:HD22	2.01	0.41
1:A:576:PRO:HA	1:A:577:PRO:HD3	1.99	0.40
2:H:133:TYR:HA	2:H:134:PRO:HD3	2.01	0.40
1:A:343:ASP:O	1:A:424:ILE:HA	2.21	0.40
1:A:554:GLN:HB2	1:A:557:HIS:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	563/700 (80%)	531 (94%)	27 (5%)	5 (1%)	14	41
2	H	216/223 (97%)	199 (92%)	13 (6%)	4 (2%)	6	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	L	212/214 (99%)	206 (97%)	6 (3%)	0	100	100
4	Z	14/17 (82%)	14 (100%)	0	0	100	100
All	All	1005/1154 (87%)	950 (94%)	46 (5%)	9 (1%)	14	41

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	SER
1	A	530	PRO
1	A	546	MET
2	H	159	GLU
2	H	166	ASN
2	H	168	GLY
1	A	604	PRO
2	H	160	SER
1	A	616	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/566 (82%)	423 (91%)	41 (9%)	9	29
2	H	190/193 (98%)	175 (92%)	15 (8%)	11	34
3	L	189/189 (100%)	174 (92%)	15 (8%)	11	34
4	Z	13/13 (100%)	13 (100%)	0	100	100
All	All	856/961 (89%)	785 (92%)	71 (8%)	10	32

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	THR
1	A	110	LYS
1	A	132	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	133	LEU
1	A	181	GLU
1	A	189	ILE
1	A	196	ILE
1	A	237	ARG
1	A	252	VAL
1	A	272	ARG
1	A	274	SER
1	A	276	LEU
1	A	289	LEU
1	A	344	GLN
1	A	361	LEU
1	A	369	ILE
1	A	375	CYS
1	A	381	SER
1	A	398	MET
1	A	405	GLU
1	A	411	LEU
1	A	435	VAL
1	A	444	LEU
1	A	470	MET
1	A	474	ILE
1	A	484	LEU
1	A	485	SER
1	A	491	ARG
1	A	501	GLU
1	A	533	ASN
1	A	546	MET
1	A	548	THR
1	A	558	VAL
1	A	575	LYS
1	A	586	ASN
1	A	589	VAL
1	A	596	ILE
1	A	612	GLU
1	A	623	THR
1	A	638	LEU
1	A	682	ARG
2	H	10	GLU
2	H	12	VAL
2	H	18	VAL
2	H	23	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	50	ARG
2	H	62	GLU
2	H	67	LYS
2	H	71	THR
2	H	82	GLN
2	H	113	VAL
2	H	128	THR
2	H	139	CYS
2	H	159	GLU
2	H	191	SER
2	H	221	LEU
3	L	12	SER
3	L	31	THR
3	L	50	SER
3	L	56	THR
3	L	60	ASP
3	L	80	SER
3	L	116	SER
3	L	125	LEU
3	L	134	CYS
3	L	153	SER
3	L	160	LEU
3	L	181	LEU
3	L	183	LYS
3	L	194	CYS
3	L	214	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	GLN
1	A	417	HIS
1	A	425	ASN
1	A	433	GLN
1	A	464	HIS
1	A	513	ASN
1	A	586	ASN
1	A	643	HIS
2	H	3	GLN
2	H	35	ASN
2	H	182	GLN
3	L	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	53	ASN
3	L	89	HIS
3	L	124	GLN
3	L	189	HIS
3	L	190	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HRG	Z	5	4	10,11,12	0.78	1 (10%)	6,12,14	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HRG	Z	5	4	-	1/9/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	Z	5	HRG	CZ-NH2	-2.08	1.26	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Z	5	HRG	NH2-CZ-NE-CD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	577/700 (82%)	-0.13	17 (2%) 53 45	29, 46, 75, 108	0
2	H	220/223 (98%)	0.11	7 (3%) 50 41	32, 59, 87, 109	0
3	L	214/214 (100%)	-0.15	6 (2%) 55 46	32, 53, 81, 101	0
4	Z	15/17 (88%)	0.74	1 (6%) 24 19	42, 51, 70, 82	1 (6%)
All	All	1026/1154 (88%)	-0.07	31 (3%) 52 43	29, 50, 80, 109	1 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	Z	10	ARG	11.9
2	H	169	SER	5.5
1	A	659	ARG	5.1
1	A	639	PRO	4.4
1	A	279	PRO	4.1
1	A	643	HIS	3.8
1	A	453	TRP	3.4
2	H	160	SER	3.3
2	H	222	GLU	3.3
1	A	671	ALA	3.1
2	H	172	SER	3.1
1	A	618	PRO	3.1
3	L	214	CYS	2.9
1	A	642	SER	2.9
2	H	166	ASN	2.7
1	A	117	GLY	2.6
3	L	199	LYS	2.6
3	L	156	GLN	2.6
1	A	375	CYS	2.5
1	A	116	HIS	2.5
1	A	61	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	11	MET	2.4
1	A	617	ALA	2.4
1	A	446	PRO	2.3
1	A	470	MET	2.3
3	L	14	SER	2.2
1	A	179	LEU	2.2
3	L	16	GLY	2.2
2	H	167	SER	2.0
2	H	202	GLN	2.0
1	A	177	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	HRG	Z	5	12/13	0.93	0.12	44,48,53,55	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NA	A	802	1/1	0.90	0.26	59,59,59,59	0
5	NA	A	801	1/1	0.91	0.12	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.