



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:27 PM UTC

PDB ID : 1VNG / pdb_00001vng
Title : CHLOROPEROXIDASE FROM THE FUNGUS CURVULARIA INAE-QUALIS: MUTANT H404A
Authors : Macedo-Ribeiro, S.; Messerschmidt, A.
Deposited on : 1999-01-20
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	609	 82% 12% 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called VANADIUM CHLOROPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	574	4477	2850	769	846	12	73	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	404	ALA	HIS	engineered mutation	UNP P49053
A	544	GLU	PRO	SEE REMARK 999	UNP P49053

- Molecule 2 is VANADATE ION (CCD ID: VO4) (formula: O₄V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	V		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	216	Total 216	O 216	0	0

i

Note EDS was not executed.

Chain A: 82% 12% 6%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	129.60 Å 129.60 Å 104.45 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.20)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.180 , 0.210	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4698	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/4610	0.95	19/6303 (0.3%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	381	LEU	N-CA-C	-9.62	100.88	111.36
1	A	123	PRO	N-CA-CB	7.37	110.19	103.19
1	A	322	ALA	N-CA-C	7.22	119.04	111.03
1	A	222	HIS	N-CA-C	6.87	119.61	108.63
1	A	271	GLN	N-CA-C	6.85	119.62	111.33
1	A	47	PRO	N-CA-C	6.79	118.98	110.70
1	A	125	PRO	N-CA-CB	6.55	110.13	103.25
1	A	399	ALA	N-CA-C	6.52	119.36	111.40
1	A	121	PRO	N-CA-CB	6.37	109.94	103.25
1	A	533	ILE	N-CA-C	5.96	116.98	108.93
1	A	173	LEU	N-CA-C	5.83	120.56	113.38
1	A	449	ARG	N-CA-C	5.78	116.18	108.38
1	A	192	LYS	N-CA-C	5.63	118.42	109.24
1	A	323	ASN	N-CA-C	5.46	119.25	112.59
1	A	27	ASN	N-CA-C	-5.45	105.42	111.36
1	A	210	ASN	CA-C-N	5.35	125.01	119.56
1	A	210	ASN	C-N-CA	5.35	125.01	119.56
1	A	451	LEU	N-CA-C	5.13	117.73	110.50
1	A	518	ALA	N-CA-C	5.06	117.64	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4477	0	4256	31	0
2	A	5	0	0	0	0
3	A	216	0	0	1	0
All	All	4698	0	4256	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:437:ASP:HB3	1:A:475:HIS:HD2	1.51	0.75
1:A:220:GLN:OE1	1:A:395:PRO:HG3	2.01	0.61
1:A:437:ASP:HB3	1:A:475:HIS:CD2	2.35	0.60
1:A:177:ARG:HH21	1:A:180:SER:HB3	1.67	0.59
1:A:479:ALA:HA	1:A:482:LEU:HD12	1.85	0.58
1:A:444:LEU:HD13	1:A:472:ILE:HD12	1.85	0.57
1:A:449:ARG:HD3	1:A:461:ILE:HG13	1.88	0.56
1:A:572:PRO:HG2	1:A:575:ILE:HG12	1.90	0.53
1:A:553:VAL:HB	1:A:554:PRO:HD3	1.91	0.52
1:A:202:VAL:HG23	1:A:218:PHE:O	2.11	0.52
1:A:449:ARG:HB2	1:A:467:ILE:O	2.10	0.51
1:A:11:LYS:HE3	3:A:1195:HOH:O	2.09	0.51
1:A:228:LYS:HG3	1:A:229:THR:HG23	1.92	0.51
1:A:241:LEU:HD23	1:A:549:PRO:HG2	1.94	0.49
1:A:218:PHE:CE2	1:A:220:GLN:HG2	2.48	0.48
1:A:40:VAL:HG11	1:A:136:GLN:HG3	1.95	0.48
1:A:360:ARG:HD3	1:A:399:ALA:HB2	1.98	0.46
1:A:345:ALA:HB1	1:A:406:THR:HG22	1.97	0.45
1:A:490:ARG:HB3	1:A:496:HIS:HB2	1.99	0.44
1:A:177:ARG:HH22	1:A:225:PHE:HB2	1.82	0.44
1:A:292:ASP:CG	1:A:383:ALA:HB1	2.43	0.44
1:A:389:ASN:HB3	1:A:453:GLN:O	2.19	0.43
1:A:549:PRO:HB2	1:A:554:PRO:HB2	2.01	0.43
1:A:291:TYR:O	1:A:300:PRO:HD3	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:ASP:O	1:A:168:VAL:HG23	2.20	0.41
1:A:136:GLN:HA	1:A:139:LEU:HD12	2.02	0.41
1:A:375:ASP:HB3	1:A:378:TRP:HB2	2.03	0.41
1:A:385:ALA:O	1:A:388:THR:HG22	2.21	0.41
1:A:336:ALA:O	1:A:340:VAL:HG23	2.21	0.40
1:A:286:GLY:HA2	1:A:489:SER:HB2	2.04	0.40
1:A:300:PRO:HB2	1:A:301:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	572/609 (94%)	547 (96%)	19 (3%)	6 (1%)	12 11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLN
1	A	124	ASN
1	A	128	ASN
1	A	290	ALA
1	A	127	ALA
1	A	178	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	466/505 (92%)	461 (99%)	5 (1%)	65 79

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	VAL
1	A	262	ARG
1	A	292	ASP
1	A	449	ARG
1	A	512	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	ASN
1	A	136	GLN
1	A	271	GLN
1	A	475	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	VO4	A	610	1	0,4,4	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.