



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:48 PM UTC

PDB ID : 6VNE / pdb_00006vne
Title : JAK2 JH1 in complex with Fedratinib
Authors : Davis, R.R.; Schonbrunn, E.
Deposited on : 2020-01-29
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	 6% 79% 15% • 5%
1	B	308	 6% 79% 14% • 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	289	Total	C	N	O	P	S	0	0	0
			2395	1521	417	442	1	14			
1	A	292	Total	C	N	O	P	S	0	0	0
			2417	1534	420	448	1	14			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	825	HIS	-	expression tag	UNP O60674
B	826	HIS	-	expression tag	UNP O60674
B	827	HIS	-	expression tag	UNP O60674
B	828	HIS	-	expression tag	UNP O60674
B	829	HIS	-	expression tag	UNP O60674
B	830	HIS	-	expression tag	UNP O60674
B	831	HIS	-	expression tag	UNP O60674
B	832	HIS	-	expression tag	UNP O60674
B	833	GLU	-	expression tag	UNP O60674
B	834	ASN	-	expression tag	UNP O60674
B	835	LEU	-	expression tag	UNP O60674
B	836	TYR	-	expression tag	UNP O60674
B	837	PHE	-	expression tag	UNP O60674
B	838	GLN	-	expression tag	UNP O60674
B	839	GLY	-	expression tag	UNP O60674
A	825	HIS	-	expression tag	UNP O60674
A	826	HIS	-	expression tag	UNP O60674
A	827	HIS	-	expression tag	UNP O60674
A	828	HIS	-	expression tag	UNP O60674
A	829	HIS	-	expression tag	UNP O60674
A	830	HIS	-	expression tag	UNP O60674
A	831	HIS	-	expression tag	UNP O60674
A	832	HIS	-	expression tag	UNP O60674
A	833	GLU	-	expression tag	UNP O60674
A	834	ASN	-	expression tag	UNP O60674

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	835	LEU	-	expression tag	UNP O60674
A	836	TYR	-	expression tag	UNP O60674
A	837	PHE	-	expression tag	UNP O60674
A	838	GLN	-	expression tag	UNP O60674
A	839	GLY	-	expression tag	UNP O60674


- ## 2TA

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	20	Total O 20 20	0	0
3	A	14	Total O 14 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK2



- Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	114.24Å 114.24Å 70.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.48 – 2.32 44.48 – 2.32	Depositor EDS
% Data completeness (in resolution range)	96.3 (44.48-2.32) 99.7 (44.48-2.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.32Å)	Xtriage
Refinement program	PHENIX 1.14-3260_3260	Depositor
R, R_{free}	0.217 , 0.256 (Not available) , 0.284	Depositor DCC
R_{free} test set	1087 reflections (2.74%)	wwPDB-VP
Wilson B-factor (Å ²)	57.6	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4920	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2TA, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2452	0.70	4/3300 (0.1%)
1	B	0.35	0/2429	0.58	1/3267 (0.0%)
All	All	0.37	0/4881	0.65	5/6567 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	922	ARG	CG-CD-NE	7.38	128.24	112.00
1	A	1011	LYS	CA-C-N	6.92	138.68	121.80
1	A	1011	LYS	C-N-CA	6.92	138.68	121.80
1	A	864	GLU	CA-CB-CG	5.89	125.88	114.10
1	B	1125	GLN	CA-CB-CG	-5.50	103.11	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2417	0	2396	35	0
1	B	2395	0	2377	27	2
2	A	37	0	36	2	0
2	B	37	0	36	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	14	0	0	1	0
3	B	20	0	0	1	0
All	All	4920	0	4845	64	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:896:GLU:OE1	1:A:922:ARG:NH1	1.93	1.00
1:A:896:GLU:CD	1:A:922:ARG:HH12	1.75	0.95
1:A:864:GLU:OE1	1:A:883:LYS:NZ	2.18	0.76
1:A:857:LYS:HG3	1:A:862:SER:HB2	1.68	0.74
1:B:853:GLN:HG2	1:B:865:MET:HE2	1.74	0.69
1:A:857:LYS:HG3	1:A:862:SER:CB	2.26	0.65
1:B:1018:ILE:HD13	1:B:1075:VAL:HG22	1.78	0.65
1:A:922:ARG:O	1:A:923:ARG:HB2	1.99	0.62
1:B:951:ILE:H	1:B:951:ILE:HD12	1.64	0.61
1:B:845:GLU:OE1	1:B:847:ARG:NH1	2.35	0.60
1:A:925:LEU:HD12	1:A:926:LYS:H	1.67	0.59
1:B:989:ARG:NH2	1:B:991:LYS:NZ	2.53	0.57
1:B:881:VAL:HG22	1:B:928:ILE:HD12	1.86	0.57
1:B:977:LEU:HD11	1:B:982:ILE:HD11	1.89	0.55
1:B:989:ARG:HH21	1:B:991:LYS:NZ	2.05	0.54
1:B:989:ARG:NH2	1:B:991:LYS:HZ1	2.06	0.53
1:A:952:LYS:NZ	1:A:988:ASN:OD1	2.42	0.53
1:A:1098:ILE:HG22	1:A:1102:MET:HE3	1.89	0.52
1:A:860:PHE:CD1	1:A:860:PHE:N	2.77	0.52
1:B:848:HIS:CG	1:B:870:PRO:HA	2.44	0.52
1:B:960:ILE:HG12	1:B:982:ILE:HD12	1.93	0.51
1:A:1012:GLU:O	1:A:1012:GLU:OE1	2.28	0.51
1:A:859:ASN:HB3	1:A:860:PHE:CD1	2.45	0.50
1:A:919:SER:OG	1:A:924:ASN:ND2	2.44	0.50
1:B:1009:LYS:HA	1:B:1029:SER:O	2.12	0.49
1:A:975:ARG:HG3	1:A:1031:PHE:CD2	2.47	0.49
1:B:1125:GLN:O	1:B:1125:GLN:HG3	2.13	0.49
1:A:859:ASN:C	1:A:860:PHE:HD1	2.21	0.48
1:A:924:ASN:N	1:A:924:ASN:OD1	2.44	0.48
1:A:954:LEU:HD12	1:A:1130:MET:HE1	1.95	0.48
1:B:971:ARG:NH2	1:B:1002:PRO:O	2.45	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:930:GLU:O	2:A:1201:2TA:H19	2.14	0.47
1:A:999:LYS:HD3	1:A:999:LYS:HA	1.64	0.47
1:A:1035:SER:HB3	3:A:1303:HOH:O	2.15	0.47
1:A:1083:LYS:HB3	1:A:1083:LYS:HE2	1.66	0.47
1:A:911:VAL:HG22	1:A:992:ILE:O	2.15	0.46
1:A:1110:VAL:HG12	1:A:1113:ARG:NH1	2.31	0.46
1:B:1035:SER:OG	3:B:1301:HOH:O	2.21	0.45
1:A:848:HIS:HA	1:A:850:LYS:HE2	1.98	0.45
1:A:918:TYR:HD2	1:A:925:LEU:HD13	1.80	0.45
1:A:1090:ARG:NH1	1:A:1094:CYS:O	2.45	0.45
1:B:902:LEU:HG	1:B:913:TYR:HB2	1.98	0.44
1:A:892:LEU:HD23	1:A:892:LEU:HA	1.72	0.44
1:A:860:PHE:N	1:A:860:PHE:HD1	2.16	0.44
1:B:1027:THR:HG22	1:B:1079:ILE:HD13	2.00	0.43
1:A:978:ALA:HA	1:A:1043:VAL:HG22	1.99	0.43
1:B:989:ARG:HH21	1:B:991:LYS:HZ1	1.62	0.43
1:A:893:ARG:HD3	1:A:893:ARG:HA	1.75	0.43
1:B:857:LYS:HE3	1:B:885:GLN:HE22	1.84	0.43
1:B:952:LYS:NZ	1:B:988:ASN:OD1	2.51	0.43
1:B:1090:ARG:HD2	1:B:1094:CYS:O	2.19	0.43
1:B:965:GLU:O	1:B:969:THR:HG23	2.18	0.43
1:A:945:LYS:HE2	1:A:945:LYS:HB2	1.75	0.43
1:B:1045:TYR:O	1:B:1049:THR:HG23	2.19	0.42
1:A:921:GLY:O	1:A:922:ARG:HB2	2.19	0.42
2:A:1201:2TA:N3	2:A:1201:2TA:H17	2.34	0.42
1:B:890:GLU:H	1:B:890:GLU:CD	2.28	0.42
1:A:925:LEU:HD12	1:A:926:LYS:N	2.32	0.42
1:B:999:LYS:HG2	1:B:1008:TYR:CZ	2.54	0.42
1:B:1069:LYS:HE3	1:B:1077:HIS:CE1	2.55	0.41
1:A:976:ASP:HB2	1:A:997:LEU:HD12	2.02	0.41
1:B:1038:TRP:CE3	1:B:1106:TRP:HA	2.56	0.40
2:B:1201:2TA:OAM	2:B:1201:2TA:H7	2.20	0.40
1:A:1044:LEU:HD21	1:A:1098:ILE:HG23	2.02	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:883:LYS:NZ	1:B:1007:PTR:O3P[2_654]	1.30	0.90
1:B:883:LYS:NZ	1:B:1007:PTR:P[2_654]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/308 (94%)	275 (95%)	12 (4%)	2 (1%)	18	22
1	B	286/308 (93%)	278 (97%)	7 (2%)	1 (0%)	36	45
All	All	575/616 (93%)	553 (96%)	19 (3%)	3 (0%)	24	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	923	ARG
1	A	920	ALA
1	B	920	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/280 (95%)	265 (100%)	1 (0%)	84	91
1	B	263/280 (94%)	263 (100%)	0	100	100
All	All	529/560 (94%)	528 (100%)	1 (0%)	87	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	840	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	874	ASN
1	B	885	GLN
1	B	891	HIS
1	A	843	GLN
1	A	942	GLN
1	A	955	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	1007	1	15,16,17	1.27	1 (6%)	17,22,24	0.50	0
1	PTR	B	1007	1	15,16,17	1.34	1 (6%)	17,22,24	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1007	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1007	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1007	PTR	OH-CZ	-4.40	1.30	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1007	PTR	OH-CZ	-3.90	1.31	1.40

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	1007	PTR	0	2

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	2TA	A	1201	-	40,40,40	3.27	12 (30%)	56,57,57	2.79	12 (21%)
2	2TA	B	1201	-	40,40,40	3.18	13 (32%)	56,57,57	2.69	12 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2TA	A	1201	-	-	5/26/33/33	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2TA	B	1201	-	-	0/26/33/33	0/4/4/4

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	2TA	CAO-NAP	-11.50	1.21	1.47
2	B	1201	2TA	CAO-NAP	-11.42	1.21	1.47
2	A	1201	2TA	CAQ-NAP	-9.56	1.28	1.47
2	B	1201	2TA	CAQ-NAP	-9.44	1.28	1.47
2	A	1201	2TA	SBD-NBE	8.97	1.75	1.61
2	B	1201	2TA	SBD-NBE	8.68	1.74	1.61
2	A	1201	2TA	C4-NAW	4.92	1.45	1.36
2	A	1201	2TA	CBB-SBD	4.24	1.83	1.76
2	A	1201	2TA	C2-NAF	4.17	1.45	1.36
2	B	1201	2TA	C2-NAF	4.10	1.45	1.36
2	B	1201	2TA	C4-NAW	3.79	1.43	1.36
2	B	1201	2TA	CAS-CAR	-3.49	1.26	1.48
2	A	1201	2TA	CAS-CAR	-3.38	1.26	1.48
2	B	1201	2TA	CBB-SBD	3.30	1.81	1.76
2	B	1201	2TA	CAR-CAQ	3.16	1.62	1.51
2	A	1201	2TA	CAR-CAQ	3.05	1.62	1.51
2	A	1201	2TA	OBK-SBD	2.86	1.46	1.43
2	A	1201	2TA	OBJ-SBD	2.76	1.46	1.43
2	B	1201	2TA	OBJ-SBD	2.73	1.46	1.43
2	B	1201	2TA	OBK-SBD	2.61	1.46	1.43
2	B	1201	2TA	CAG-NAF	2.59	1.46	1.40
2	A	1201	2TA	CAG-NAF	2.59	1.46	1.40
2	A	1201	2TA	CAT-NAP	2.26	1.52	1.47
2	B	1201	2TA	CAT-NAP	2.18	1.52	1.47
2	B	1201	2TA	C5-C4	-2.05	1.38	1.41

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	2TA	OBK-SBD-OBJ	-16.79	99.13	119.52
2	B	1201	2TA	OBK-SBD-OBJ	-15.45	100.75	119.52
2	A	1201	2TA	C5-C6-N1	-5.89	119.18	125.09
2	B	1201	2TA	C5-C6-N1	-5.22	119.86	125.09
2	B	1201	2TA	N1-C2-N3	-4.99	121.60	126.42
2	A	1201	2TA	N1-C2-N3	-4.19	122.37	126.42
2	B	1201	2TA	CAA-C5-C4	-3.67	118.87	122.10
2	A	1201	2TA	C6-N1-C2	3.48	120.58	115.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1201	2TA	C6-N1-C2	3.45	120.55	115.81
2	B	1201	2TA	OBJ-SBD-NBE	3.40	115.14	107.36
2	A	1201	2TA	CBB-SBD-NBE	3.02	113.48	108.11
2	B	1201	2TA	C5-C4-NAW	-2.92	115.25	119.44
2	A	1201	2TA	C5-C4-N3	-2.90	120.08	123.46
2	A	1201	2TA	CAT-NAP-CAQ	2.80	107.34	104.04
2	B	1201	2TA	CAX-NAW-C4	-2.74	122.71	129.43
2	A	1201	2TA	OBK-SBD-CBB	2.74	111.43	107.98
2	B	1201	2TA	NAW-C4-N3	2.68	123.05	119.08
2	B	1201	2TA	CAT-NAP-CAQ	2.63	107.14	104.04
2	B	1201	2TA	CAA-C5-C6	2.58	124.51	120.01
2	A	1201	2TA	OBK-SBD-NBE	2.50	113.09	107.36
2	A	1201	2TA	CBI-CBF-CBH	-2.42	104.61	109.95
2	A	1201	2TA	CAA-C5-C4	-2.23	120.13	122.10
2	B	1201	2TA	CBB-SBD-NBE	2.23	112.09	108.11
2	A	1201	2TA	CAX-NAW-C4	-2.04	124.42	129.43

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	2TA	CAN-CAO-NAP-CAT
2	A	1201	2TA	OAM-CAN-CAO-NAP
2	A	1201	2TA	CBF-NBE-SBD-OBK
2	A	1201	2TA	CAK-CAJ-OAM-CAN
2	A	1201	2TA	CAI-CAJ-OAM-CAN



There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1201	2TA	2	0
2	B	1201	2TA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	291/308 (94%)	0.72	20 (6%) 23 25	49, 72, 113, 150	0
1	B	288/308 (93%)	0.70	17 (5%) 28 31	47, 69, 116, 142	0
All	All	579/616 (93%)	0.71	37 (6%) 25 28	47, 71, 116, 150	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	920	ALA	4.4
1	B	886	HIS	3.9
1	A	841	PRO	3.6
1	A	860	PHE	3.4
1	A	842	THR	3.2
1	B	1013	PRO	3.0
1	B	860	PHE	2.8
1	A	864	GLU	2.7
1	A	1011	LYS	2.6
1	B	887	SER	2.6
1	A	890	GLU	2.6
1	A	1010	VAL	2.6
1	A	1053	LYS	2.6
1	B	924	ASN	2.5
1	A	1131	ALA	2.5
1	A	923	ARG	2.5
1	A	1100	MET	2.5
1	B	843	GLN	2.5
1	B	1076	PHE	2.4
1	A	1101	ILE	2.4
1	B	1066	GLY	2.4
1	A	946	GLU	2.3
1	B	1053	LYS	2.3
1	A	922	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1014	GLY	2.2
1	B	1121	LEU	2.2
1	A	1014	GLY	2.2
1	A	877	GLU	2.2
1	B	1008	TYR	2.2
1	B	923	ARG	2.2
1	A	1016	SER	2.2
1	B	1064	MET	2.1
1	A	902	LEU	2.1
1	B	918	TYR	2.1
1	A	840	ASP	2.1
1	B	994	ASP	2.0
1	A	976	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	PTR	B	1007	16/17	0.91	0.12	51,81,137,140	0
1	PTR	A	1007	16/17	0.93	0.10	76,98,149,149	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	2TA	A	1201	37/37	0.94	0.11	49,64,98,107	0
2	2TA	B	1201	37/37	0.95	0.10	33,53,93,95	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.