

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	145	Total 145	O 145	0	1
7	B	146	Total 146	O 146	0	0

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	12	0	16	0	0
5	B	6	0	8	0	0
5	C	12	0	16	0	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
7	A	145	0	0	1	0
7	B	146	0	0	0	0
7	C	49	0	0	0	0
7	D	57	0	0	0	0
All	All	3867	0	2946	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:417:DA:H62	2:B:263:GLN:HE22	1.38	0.72
1:D:513[B]:DA:O3'	1:D:514[B]:DG:P	2.58	0.62
2:A:91:GLY:HA2	7:A:810:HOH:O	2.14	0.47
1:C:417:DA:H62	2:B:263:GLN:NE2	2.09	0.47
1:D:513[B]:DA:O3'	1:D:514[B]:DG:OP2	2.33	0.46
1:C:407:DC:H2'	1:C:408:DT:C6	2.53	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	160/162 (99%)	159 (99%)	1 (1%)	0	100	100
2	B	160/162 (99%)	160 (100%)	0	0	100	100
All	All	320/324 (99%)	319 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	133/133 (100%)	133 (100%)	0	100	100
2	B	133/133 (100%)	132 (99%)	1 (1%)	73	51
All	All	266/266 (100%)	265 (100%)	1 (0%)	84	70

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	271	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	36	GLN
2	A	69	ASN
2	A	129	ASN
2	B	205	ASN
2	B	263	GLN
2	B	269	ASN
2	B	273	HIS
2	B	288	ASN
2	B	360	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 8 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	C	503	-	5,5,5	0.09	0	5,5,5	0.35	0
5	GOL	A	603	-	5,5,5	0.09	0	5,5,5	0.54	0
5	GOL	A	604	-	5,5,5	0.07	0	5,5,5	0.39	0
5	GOL	B	403	-	5,5,5	0.06	0	5,5,5	0.29	0
5	GOL	C	504	-	5,5,5	0.08	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	503	-	-	0/4/4/4	-
5	GOL	A	603	-	-	3/4/4/4	-
5	GOL	A	604	-	-	2/4/4/4	-
5	GOL	B	403	-	-	0/4/4/4	-
5	GOL	C	504	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	603	GOL	C1-C2-C3-O3
5	A	604	GOL	C1-C2-C3-O3
5	A	603	GOL	O2-C2-C3-O3
5	A	604	GOL	O2-C2-C3-O3
5	A	603	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	413[B]:DA	O3'	414[B]:DG	P	2.78
1	D	513[B]:DA	O3'	514[B]:DG	P	2.58

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	C	21/21 (100%)	0.05	0	100	100	9, 25, 30, 33	2 (9%)
1	D	21/21 (100%)	0.01	0	100	100	9, 26, 30, 32	2 (9%)
2	A	162/162 (100%)	0.25	9 (5%)	30	37	15, 20, 32, 42	0
2	B	162/162 (100%)	0.28	8 (4%)	35	43	15, 20, 37, 49	0
All	All	366/366 (100%)	0.24	17 (4%)	37	45	9, 21, 33, 49	4 (1%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	203	LEU	7.0
2	B	202	ALA	6.9
2	A	91	GLY	6.2
2	A	90	ASN	5.3
2	B	271	VAL	4.0
2	A	89	ILE	4.0
2	B	290	ASN	3.6
2	A	69	ASN	3.0
2	B	291	GLY	2.9
2	B	204	THR	2.9
2	A	2	ALA	2.8
2	A	70	GLN	2.8
2	B	270	GLN	2.5
2	B	269	ASN	2.5
2	A	68	ILE	2.4
2	A	35	LEU	2.1
2	A	129	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	C	503	6/6	0.74	0.21	26,30,34,50	0
5	GOL	A	604	6/6	0.78	0.18	37,40,43,44	0
5	GOL	C	504	6/6	0.83	0.16	26,30,36,45	0
5	GOL	A	603	6/6	0.91	0.12	21,26,35,37	0
5	GOL	B	403	6/6	0.95	0.08	18,20,21,25	0
4	NA	D	602	1/1	0.97	0.04	16,16,16,16	1
4	NA	C	502	1/1	0.98	0.06	16,16,16,16	1
3	MG	D	601	1/1	0.99	0.03	16,16,16,16	1
6	ZN	A	601	1/1	0.99	0.02	18,18,18,18	0
6	ZN	A	602	1/1	0.99	0.02	16,16,16,16	0
3	MG	C	501	1/1	1.00	0.05	16,16,16,16	0
6	ZN	B	401	1/1	1.00	0.02	17,17,17,17	0
6	ZN	B	402	1/1	1.00	0.01	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NA D 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NA C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG D 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.