



Full wwPDB EM Validation Report ⓘ

Mar 28, 2026 – 10:01 PM UTC

PDB ID : 2VOY / pdb_00002voy
EMDB ID : EMD-5004
Title : CryoEM model of CopA, the copper transporting ATPase from *Archaeoglobus fulgidus*
Authors : Wu, C.-C.; Rice, W.J.; Stokes, D.L.
Deposited on : 2008-02-25
Resolution : 18.00 Å (reported)
Based on initial models : 2EAR, 2B8E, 2HC8

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : **FAILED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 18.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Q-score	25397	45 (17.00 - 18.00)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 5270 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POTENTIAL COPPER-TRANSPORTING ATPASE.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	80	Total	C	N	O	S	0	0
			615	382	104	126	3		

- Molecule 2 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	42	Total	C	N	O	S	0	0
			341	227	52	61	1		

- Molecule 3 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	22	Total	C	N	O	S	0	0
			176	123	25	27	1		

- Molecule 4 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	23	Total	C	N	O	S	0	0
			176	121	28	26	1		

- Molecule 5 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
5	E	30	Total	C	N	O	0	0
			234	155	38	41		

- Molecule 6 is a protein called CATION-TRANSPORTING ATPASE, P-TYPE.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	113	Total	C	N	O	S	0	0
			829	525	138	164	2		

- Molecule 7 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	36	Total	C	N	O	S	0	0
			287	185	47	54	1		

- Molecule 8 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	48	Total	C	N	O	S	0	0
			356	233	64	57	2		

- Molecule 9 is a protein called CATION-TRANSPORTING ATPASE.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	128	Total	C	N	O	Se	0	0
			959	604	165	188	2		

- Molecule 10 is a protein called CATION-TRANSPORTING ATPASE.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	118	Total	C	N	O	Se	0	0
			887	553	157	176	1		

- Molecule 11 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	32	Total	C	N	O	S	0	0
			253	163	43	45	2		

- Molecule 12 is a protein called SARCOPLASMIC/ENDOPLASMIC RETICULUM CALCIUM ATPASE 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
12	L	21	Total	C	N	O	0	0
			157	106	24	27		

MolProbity failed to run properly - this section is therefore empty.

3 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	Not provided	
Resolution determination method	Not provided	
CTF correction method	INDIVIDUAL TUBES	Depositor
Microscope	FEI/PHILIPS CM200FEG/ST	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	10	Depositor
Minimum defocus (nm)	900.00	Depositor
Maximum defocus (nm)	2500.00	Depositor
Magnification	51300	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	75.388	Depositor
Minimum map value	-48.614	Depositor
Average map value	0.529	Depositor
Map value standard deviation	4.943	Depositor
Recommended contour level	10	Depositor
Map size (Å)	202, 202, 202	wwPDB
Map dimensions	101, 101, 101	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	2, 2, 2	Depositor

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
9	I	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	I	636:GLY	C	645:SER	N	12.26
1	I	431:LYS	C	550:LEU	N	7.12


5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-5004. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

5.1 Orthogonal projections [i](#)


5.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

5.2 Central slices [i](#)

5.2.1 Primary map



X Index: 50



Y Index: 50



Z Index: 50

The images above show central slices of the map in three orthogonal directions.

5.3 Largest variance slices [i](#)

5.3.1 Primary map



X Index: 50



Y Index: 47



Z Index: 57


The images above show the largest variance slices of the map in three orthogonal directions.

5.4 Orthogonal standard-deviation projections (False-color) [i](#)

5.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

5.5 Orthogonal surface views [i](#)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 10.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


6.2 Volume estimate [i](#)



The volume at the recommended contour level is 203 nm^3 ; this corresponds to an approximate mass of 183 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

6.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.056 Å⁻¹


7 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-5004 and PDB model 2VOY. Per-residue inclusion information can be found in section ?? on page ??.


8.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 10.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (10).

8.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (10) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7870	0.0410
A	0.7380	0.0140
B	0.7590	0.0670
C	0.8060	0.0640
D	0.9350	0.1190
E	0.7620	-0.0070
F	0.7290	0.0520
G	0.4930	0.0500
H	0.8240	-0.0010
I	0.8130	0.0450
J	0.9100	0.0490
K	0.7450	0.0430
L	0.8850	0.0080

