



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 08:56 PM UTC

PDB ID : 5VOP / pdb_00005vop
Title : Methionine synthase folate-binding domain from *Thermus thermophilus* HB8 native
Authors : Koutmos, M.; Yamada, K.
Deposited on : 2017-05-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	90% 5% 5%
1	B	296	2% 90% 5% 5%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 9262 atoms, of which 4601 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5-methyltetrahydrofolate homocysteine S-methyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	282	Total	C	H	N	O	S	0	0	0
			4480	1412	2263	390	408	7			
1	B	282	Total	C	H	N	O	S	0	0	0
			4480	1412	2263	390	408	7			

- Molecule 2 is 5-METHYL-5,6,7,8-TETRAHYDROFOLIC ACID (CCD ID: C2F) (formula: $C_{20}H_{25}N_7O_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			56	20	23	7	6		
2	B	1	Total	C	H	N	O	0	0
			56	20	23	7	6		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		
3	A	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			18	6	5	7		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Na 1	0	0
5	B	1	Total 1	Na 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	81	Total 81	O 81	0	0
6	B	47	Total 47	O 47	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5-methyltetrahydrofolate homocysteine S-methyltransferase

Chain A: 



- Molecule 1: 5-methyltetrahydrofolate homocysteine S-methyltransferase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	111.48Å 111.48Å 242.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.44 – 2.10 29.44 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.44-2.10) 99.9 (29.44-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.194 , 0.222 (Not available) , 0.224	Depositor DCC
R_{free} test set	2667 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9262	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, C2F, NA, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2255	0.90	3/3048 (0.1%)
1	B	0.65	0/2255	0.86	1/3048 (0.0%)
All	All	0.67	0/4510	0.88	4/6096 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	616	PRO	CA-C-N	11.31	135.80	120.54
1	A	616	PRO	C-N-CA	11.31	135.80	120.54
1	A	519	GLU	N-CA-C	-5.72	102.43	110.50
1	B	574	VAL	CB-CA-C	-5.52	103.61	112.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2217	2263	2263	9	0
1	B	2217	2263	2263	9	0
2	A	33	23	23	0	0
2	B	33	23	23	0	0
3	A	18	24	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	13	5	5	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	81	0	0	1	0
6	B	47	0	0	0	0
All	All	4661	4601	4601	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:581:ARG:NH2	1:A:648:HIS:O	2.18	0.77
1:A:442:VAL:HG21	1:A:453:ALA:CB	2.35	0.56
1:A:426:ARG:NH1	6:A:3101:HOH:O	2.39	0.54
1:A:442:VAL:HG21	1:A:453:ALA:HB1	1.90	0.53
1:A:368:LEU:HD13	1:A:591:LEU:HD11	1.93	0.50
1:B:555:MET:HE1	1:B:569:LEU:HD11	1.94	0.48
1:B:534:THR:HG23	1:B:574:VAL:HG13	1.95	0.48
1:B:592:ASP:OD1	1:B:595:ARG:NH1	2.47	0.48
1:A:491:VAL:HG11	1:A:517:LEU:HD11	1.96	0.47
1:B:571:VAL:O	1:B:574:VAL:HG22	2.15	0.47
1:B:414:VAL:HG23	1:B:421:GLU:HG3	1.97	0.46
3:A:3003:GOL:H12	1:B:519:GLU:HB2	1.97	0.46
1:A:516:ARG:O	1:A:519:GLU:O	2.34	0.45
1:A:368:LEU:CD1	1:A:591:LEU:HD11	2.47	0.45
1:B:503:ARG:NH1	1:B:553:LEU:HD13	2.33	0.44
1:A:473:GLU:HB3	4:A:3004:FLC:HA2	2.00	0.43
1:B:534:THR:CG2	1:B:574:VAL:HG13	2.50	0.42
1:B:639:LEU:HD23	1:B:642:MET:CE	2.50	0.42

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/296 (95%)	274 (98%)	5 (2%)	1 (0%)	30	28
1	B	280/296 (95%)	274 (98%)	5 (2%)	1 (0%)	30	28
All	All	560/592 (95%)	548 (98%)	10 (2%)	2 (0%)	30	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	533	LEU
1	B	533	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/242 (96%)	230 (100%)	1 (0%)	84	89
1	B	231/242 (96%)	230 (100%)	1 (0%)	84	89
All	All	462/484 (96%)	460 (100%)	2 (0%)	84	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	633	GLU
1	B	574	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	587	ASN
1	A	614	GLN
1	A	648	HIS
1	B	587	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	C2F	B	3001	-	32,35,35	1.15	3 (9%)	36,49,49	1.43	5 (13%)
3	GOL	A	3002	-	5,5,5	0.39	0	5,5,5	0.45	0
4	FLC	A	3004	-	12,12,12	1.37	1 (8%)	17,17,17	1.91	7 (41%)
2	C2F	A	3001	-	32,35,35	1.13	3 (9%)	36,49,49	1.53	6 (16%)
3	GOL	A	3003	-	5,5,5	0.34	0	5,5,5	0.41	0
3	GOL	A	3006	-	5,5,5	0.80	0	5,5,5	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2F	B	3001	-	-	3/22/35/35	0/3/3/3
3	GOL	A	3002	-	-	0/4/4/4	-
4	FLC	A	3004	-	-	6/16/16/16	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2F	A	3001	-	-	2/22/35/35	0/3/3/3
3	GOL	A	3003	-	-	2/4/4/4	-
3	GOL	A	3006	-	-	2/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3001	C2F	C4-N3	-3.12	1.33	1.38
4	A	3004	FLC	CB-CBC	-2.99	1.50	1.53
2	A	3001	C2F	C7-C6	2.63	1.55	1.52
2	B	3001	C2F	C7-C6	2.60	1.55	1.52
2	B	3001	C2F	C4A-C4	2.14	1.48	1.43
2	A	3001	C2F	C4-N3	-2.12	1.34	1.38
2	A	3001	C2F	C4A-C4	2.07	1.48	1.43

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3001	C2F	C2-N1-C8A	4.18	120.74	113.36
2	A	3001	C2F	C2-N1-C8A	4.07	120.55	113.36
2	B	3001	C2F	C4A-C4-N3	3.50	117.10	110.94
4	A	3004	FLC	OHB-CB-CBC	3.37	113.74	108.96
2	A	3001	C2F	CB-CA-CT	-3.12	102.95	110.35
2	A	3001	C2F	C4A-C4-N3	3.04	116.28	110.94
4	A	3004	FLC	OHB-CB-CG	-2.90	102.76	109.38
2	A	3001	C2F	C12-C-N	2.79	122.22	117.04
4	A	3004	FLC	CB-CG-CGC	-2.75	106.39	113.92
4	A	3004	FLC	OB2-CBC-CB	2.46	117.86	113.14
2	A	3001	C2F	O-C-C12	-2.39	116.16	120.90
4	A	3004	FLC	CG-CB-CBC	-2.35	104.83	110.03
2	B	3001	C2F	CG-CB-CA	2.32	117.44	113.16
4	A	3004	FLC	CB-CA-CAC	2.30	120.22	113.92
2	B	3001	C2F	C12-C-N	2.20	121.12	117.04
2	A	3001	C2F	CB-CA-N	2.14	115.14	110.91
2	B	3001	C2F	CB-CG-CD	-2.12	106.85	112.49
4	A	3004	FLC	OA2-CAC-CA	2.03	120.77	114.35

There are no chirality outliers.

All (15) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	3003	GOL	C1-C2-C3-O3
3	A	3003	GOL	O2-C2-C3-O3
3	A	3006	GOL	C1-C2-C3-O3
3	A	3006	GOL	O2-C2-C3-O3
4	A	3004	FLC	CAC-CA-CB-OHB
4	A	3004	FLC	OHB-CB-CG-CGC
4	A	3004	FLC	CAC-CA-CB-CBC
4	A	3004	FLC	CBC-CB-CG-CGC
4	A	3004	FLC	CA-CB-CG-CGC
4	A	3004	FLC	CAC-CA-CB-CG
2	B	3001	C2F	OE1-CD-CG-CB
2	B	3001	C2F	OE2-CD-CG-CB
2	A	3001	C2F	OE1-CD-CG-CB
2	B	3001	C2F	N-CA-CT-O1
2	A	3001	C2F	OE2-CD-CG-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3004	FLC	1	0
3	A	3003	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/296 (95%)	-0.32	3 (1%) 78 80	44, 52, 74, 94	0
1	B	282/296 (95%)	0.07	7 (2%) 58 61	48, 61, 102, 134	0
All	All	564/592 (95%)	-0.13	10 (1%) 67 70	44, 56, 93, 134	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	635	PHE	4.1
1	B	648	HIS	3.7
1	A	417	THR	3.5
1	A	459	GLY	2.9
1	B	574	VAL	2.7
1	A	458	PRO	2.7
1	B	540	GLY	2.5
1	B	460	ARG	2.4
1	B	647	ALA	2.4
1	B	641	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NA	B	3002	1/1	0.69	0.23	82,82,82,82	0
3	GOL	A	3002	6/6	0.82	0.15	78,90,95,95	0
5	NA	A	3005	1/1	0.85	0.17	68,68,68,68	0
3	GOL	A	3006	6/6	0.88	0.16	55,67,69,69	0
2	C2F	A	3001	33/33	0.92	0.11	42,59,69,72	56
2	C2F	B	3001	33/33	0.92	0.12	55,64,78,82	56
4	FLC	A	3004	13/13	0.94	0.09	40,48,56,63	18
3	GOL	A	3003	6/6	0.95	0.10	54,60,65,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.