



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:06 PM UTC

PDB ID : 6VO8 / pdb_00006vo8
Title : X-ray structure of the Cj1427 in the presence of NADH and GDP-D-glycero-D-mannoheptose, an essential NAD-dependent dehydrogenase from *Campylobacter jejuni*
Authors : Spencer, K.D.; Anderson, T.K.; Thoden, J.B.; Huddleston, J.P.; Raushel, F.M.; Holden, H.M.
Deposited on : 2020-01-30
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	 82% 10% • 6%
1	B	321	 2% 76% 15% • 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5165 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative sugar-nucleotide epimerase/dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2416	1549	406	446	15			
1	B	299	Total	C	N	O	S	0	0	0
			2385	1531	401	438	15			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	314	LEU	-	expression tag	UNP Q0P8I7
A	315	GLU	-	expression tag	UNP Q0P8I7
A	316	HIS	-	expression tag	UNP Q0P8I7
A	317	HIS	-	expression tag	UNP Q0P8I7
A	318	HIS	-	expression tag	UNP Q0P8I7
A	319	HIS	-	expression tag	UNP Q0P8I7
A	320	HIS	-	expression tag	UNP Q0P8I7
A	321	HIS	-	expression tag	UNP Q0P8I7
B	314	LEU	-	expression tag	UNP Q0P8I7
B	315	GLU	-	expression tag	UNP Q0P8I7
B	316	HIS	-	expression tag	UNP Q0P8I7
B	317	HIS	-	expression tag	UNP Q0P8I7
B	318	HIS	-	expression tag	UNP Q0P8I7
B	319	HIS	-	expression tag	UNP Q0P8I7
B	320	HIS	-	expression tag	UNP Q0P8I7
B	321	HIS	-	expression tag	UNP Q0P8I7

- Molecule 2 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (CCD ID: NAI) (formula: C₂₁H₂₉N₇O₁₄P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is [(2 {R},3 {S},4 {R},5 {R})-5-(2-azanylidene-3 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl [(2 {R},3 {S},4 {S},5 {S},6 {S})-6-[(1 {S})-1,2-bis(oxidanyl)ethyl]-3,4,5-tris(oxidanyl)oxan-2-yl] hydrogen phosphate (CCD ID: GZ0) (formula: C₁₇H₂₇N₅O₁₇P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			41	17	5	17	2		
3	B	1	Total	C	N	O	P	0	0
			41	17	5	17	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	128	Total	O	0	0
			128	128		
4	B	66	Total	O	0	0
			66	66		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative sugar-nucleotide epimerase/dehydratase



- Molecule 1: Putative sugar-nucleotide epimerase/dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	98.72Å 98.85Å 72.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.01 – 2.40 30.01 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.9 (30.01-2.40) 98.9 (30.01-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.206 , 0.286 0.211 , 0.285	Depositor DCC
R_{free} test set	1590 reflections (5.56%)	wwPDB-VP
Wilson B-factor (Å ²)	20.9	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.032 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5165	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, GZ0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.09	0/2465	1.49	5/3325 (0.2%)
1	B	1.05	0/2433	1.50	6/3281 (0.2%)
All	All	1.07	0/4898	1.49	11/6606 (0.2%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ASP	CA-CB-CG	5.66	118.26	112.60
1	A	244	GLN	N-CA-C	-5.35	105.44	111.28
1	B	292	ASP	CA-C-N	5.35	126.03	120.03
1	B	292	ASP	C-N-CA	5.35	126.03	120.03
1	B	104	ILE	CA-C-N	5.26	127.65	120.54
1	B	104	ILE	C-N-CA	5.26	127.65	120.54
1	A	205	ASN	CA-C-O	-5.15	115.58	121.40
1	A	112	GLN	CA-C-N	5.08	128.92	122.37
1	A	112	GLN	C-N-CA	5.08	128.92	122.37
1	B	234	GLY	CA-C-O	-5.05	116.31	121.61
1	B	276	ASN	N-CA-C	-5.01	106.86	113.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2433	20	0
1	B	2385	0	2404	39	0
2	A	44	0	27	3	0
2	B	44	0	27	2	0
3	A	41	0	0	1	0
3	B	41	0	0	2	0
4	A	128	0	0	4	0
4	B	66	0	0	2	0
All	All	5165	0	4891	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:MET:HE2	1:B:95:MET:HB3	1.53	0.89
1:A:130:ALA:O	4:A:501:HOH:O	1.98	0.81
1:B:59:MET:CE	1:B:95:MET:HB3	2.24	0.64
1:A:55:ASN:C	1:A:55:ASN:HD22	2.06	0.62
1:B:55:ASN:C	1:B:55:ASN:HD22	2.07	0.62
1:B:304:MET:O	4:B:501:HOH:O	2.17	0.57
1:A:166:LEU:HB2	2:A:401:NAI:H5N	1.87	0.57
1:B:90:PRO:O	1:B:93:ALA:HB3	2.05	0.57
1:A:120:ASN:ND2	3:A:402:GZ0:O71	2.39	0.56
1:B:191:ARG:HD2	1:B:192:ASP:OD1	2.06	0.56
1:A:71:ALA:O	1:A:112:GLN:NE2	2.35	0.55
1:A:134:GLU:HG2	4:A:603:HOH:O	2.06	0.55
1:B:96:ILE:HD12	2:B:401:NAI:H61A	1.71	0.55
1:B:262:ASN:H	1:B:262:ASN:HD22	1.56	0.53
1:B:291:GLU:HG3	4:B:559:HOH:O	2.09	0.53
1:B:13:TYR:HB2	1:B:177:ARG:HG3	1.91	0.52
1:B:262:ASN:HD22	1:B:262:ASN:N	2.08	0.51
1:B:262:ASN:N	1:B:262:ASN:ND2	2.58	0.51
1:A:268:ASP:O	1:A:269:LYS:C	2.53	0.50
1:B:59:MET:CE	1:B:95:MET:CB	2.89	0.50
1:B:59:MET:HE3	1:B:96:ILE:HD13	1.93	0.50
1:A:229:GLN:HB2	1:A:231:TYR:CE2	2.47	0.50
1:A:145:GLY:O	1:A:149:VAL:HG23	2.12	0.50
1:A:169:VAL:HA	1:A:207:ILE:O	2.12	0.50
1:B:191:ARG:NH2	1:B:192:ASP:OD2	2.43	0.50
2:A:401:NAI:O1N	2:A:401:NAI:N7N	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:LEU:HD12	2:B:401:NAI:C5N	2.43	0.48
1:A:166:LEU:HB2	2:A:401:NAI:C5N	2.43	0.48
1:B:191:ARG:CD	1:B:192:ASP:OD1	2.62	0.48
1:B:229:GLN:HB2	1:B:231:TYR:CE2	2.49	0.47
1:A:128:LYS:HA	4:A:528:HOH:O	2.13	0.47
1:B:68:VAL:O	1:B:112:GLN:NE2	2.48	0.46
1:A:98:TYR:CZ	1:A:150:HIS:NE2	2.84	0.45
1:B:40:ILE:HB	1:B:43:LEU:HD22	1.98	0.45
1:B:120:ASN:C	1:B:120:ASN:HD22	2.23	0.45
1:B:120:ASN:ND2	1:B:272:TYR:OH	2.46	0.45
1:B:43:LEU:HD12	1:B:43:LEU:HA	1.87	0.45
1:B:131:MET:O	1:B:131:MET:HG3	2.14	0.44
1:B:118:ASN:OD1	1:B:148:LYS:HB3	2.17	0.44
1:B:198:PHE:CE2	3:B:402:GZ0:C5	3.00	0.44
1:B:102:LYS:HG3	1:B:154:TYR:CE1	2.52	0.44
1:A:200:GLU:OE2	1:A:243:ARG:NH2	2.51	0.43
1:B:134:GLU:HG2	1:B:230:ALA:O	2.18	0.43
1:A:210:ARG:HD2	1:B:47:HIS:CD2	2.53	0.43
1:B:242:LYS:NZ	3:B:402:GZ0:N7	2.67	0.43
1:B:4:LYS:HD2	1:B:70:LYS:O	2.19	0.43
1:B:177:ARG:NH1	1:B:180:LEU:HD21	2.34	0.43
1:A:120:ASN:HD22	1:A:120:ASN:C	2.26	0.42
1:B:120:ASN:C	1:B:120:ASN:ND2	2.76	0.42
1:B:96:ILE:O	1:B:100:ALA:HB3	2.20	0.42
1:A:99:GLU:O	1:A:102:LYS:HB3	2.20	0.42
1:A:266:ASP:HB2	4:A:573:HOH:O	2.20	0.41
1:A:231:TYR:HB3	1:A:279:LEU:HB2	2.01	0.41
1:B:55:ASN:C	1:B:55:ASN:ND2	2.75	0.41
1:B:191:ARG:HD2	1:B:192:ASP:CG	2.46	0.41
1:B:167:ALA:HB1	1:B:205:ASN:C	2.45	0.41
1:B:83:ALA:N	1:B:84:PRO:CD	2.84	0.41
1:A:242:LYS:O	1:A:245:LEU:HB3	2.21	0.40
1:B:196:VAL:HA	1:B:259:HIS:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/321 (94%)	285 (95%)	16 (5%)	0	100	100
1	B	295/321 (92%)	275 (93%)	18 (6%)	2 (1%)	18	28
All	All	596/642 (93%)	560 (94%)	34 (6%)	2 (0%)	36	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	60	ASP
1	B	181	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/280 (94%)	246 (94%)	17 (6%)	15	27
1	B	260/280 (93%)	243 (94%)	17 (6%)	15	27
All	All	523/560 (93%)	489 (94%)	34 (6%)	15	27

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	4	LYS
1	A	54	ILE
1	A	55	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	61	GLU
1	A	91	LYS
1	A	92	LEU
1	A	120	ASN
1	A	128	LYS
1	A	129	ASP
1	A	134	GLU
1	A	175	ARG
1	A	203	ARG
1	A	205	ASN
1	A	237	SER
1	A	243	ARG
1	A	272	TYR
1	B	4	LYS
1	B	43	LEU
1	B	55	ASN
1	B	61	GLU
1	B	87	LYS
1	B	96	ILE
1	B	105	SER
1	B	120	ASN
1	B	131	MET
1	B	150	HIS
1	B	175	ARG
1	B	205	ASN
1	B	243	ARG
1	B	260	SER
1	B	262	ASN
1	B	272	TYR
1	B	275	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	120	ASN
1	A	205	ASN
1	B	55	ASN
1	B	66	GLN
1	B	120	ASN
1	B	205	ASN
1	B	262	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAI	B	401	-	47,48,48	1.24	7 (14%)	64,73,73	1.76	16 (25%)
2	NAI	A	401	-	47,48,48	1.31	5 (10%)	64,73,73	1.80	14 (21%)
3	GZ0	A	402	-	43,44,44	1.21	3 (6%)	58,68,68	2.04	9 (15%)
3	GZ0	B	402	-	43,44,44	0.99	3 (6%)	58,68,68	2.34	14 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAI	B	401	-	-	11/29/72/72	0/5/5/5
2	NAI	A	401	-	-	10/29/72/72	0/5/5/5
3	GZ0	A	402	-	-	12/27/63/63	0/4/4/4
3	GZ0	B	402	-	-	11/27/63/63	0/4/4/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAI	C5A-C4A	4.15	1.46	1.39
3	A	402	GZ0	O51-C51	-4.02	1.37	1.44
2	B	401	NAI	C5A-C4A	3.65	1.45	1.39
3	A	402	GZ0	C4-N9	-3.60	1.33	1.37
3	B	402	GZ0	C4-N9	-3.05	1.34	1.37
3	B	402	GZ0	C6-N1	-2.81	1.33	1.38
2	B	401	NAI	C2N-C3N	2.76	1.42	1.35
2	A	401	NAI	C4A-N9A	-2.75	1.32	1.37
2	A	401	NAI	C5A-C6A	2.74	1.48	1.41
2	B	401	NAI	C4A-N9A	-2.58	1.32	1.37
2	A	401	NAI	PN-O3	2.58	1.62	1.59
3	A	402	GZ0	C5-N7	-2.40	1.34	1.39
2	A	401	NAI	C4N-C5N	-2.29	1.43	1.49
3	B	402	GZ0	C5-N7	-2.28	1.34	1.39
2	B	401	NAI	C5A-C6A	2.23	1.47	1.41
2	B	401	NAI	C6N-C5N	2.22	1.39	1.33
2	B	401	NAI	C8A-N7A	2.21	1.35	1.31
2	B	401	NAI	C5A-N7A	-2.03	1.35	1.39

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	GZ0	O51-C51-C41	9.19	120.93	107.94
3	B	402	GZ0	C11-O51-C51	8.93	127.39	113.07
3	B	402	GZ0	O51-C11-O1B	-8.84	99.81	111.36
3	A	402	GZ0	C5-C4-N9	5.63	108.85	106.25
2	A	401	NAI	N3A-C2A-N1A	-5.53	120.22	128.58
2	B	401	NAI	N3A-C2A-N1A	-5.41	120.39	128.58
3	A	402	GZ0	C11-O51-C51	5.33	121.61	113.07
2	A	401	NAI	C5A-C4A-N3A	-5.17	119.60	126.72
3	B	402	GZ0	O51-C51-C41	5.03	115.05	107.94
2	A	401	NAI	C2A-N3A-C4A	4.66	123.22	111.83
2	A	401	NAI	N3A-C4A-N9A	4.49	134.81	127.17
2	B	401	NAI	C2A-N3A-C4A	4.38	122.52	111.83
2	B	401	NAI	C5A-C4A-N3A	-4.13	121.03	126.72
2	B	401	NAI	N3A-C4A-N9A	3.50	133.12	127.17
2	B	401	NAI	C4A-N9A-C8A	3.47	109.38	105.74
2	A	401	NAI	C4A-N9A-C8A	3.32	109.22	105.74
3	B	402	GZ0	O31-C31-C41	-3.20	102.82	110.38
2	A	401	NAI	C2A-N1A-C6A	3.17	123.94	118.73
2	B	401	NAI	C6A-C5A-N7A	3.16	138.17	132.09
3	B	402	GZ0	C8-N9-C4	3.12	109.30	106.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	GZ0	O51-C51-C61	3.11	112.65	106.60
3	A	402	GZ0	O2A-PA-O3A	3.08	115.59	107.27
3	B	402	GZ0	O3B-PB-O3A	2.92	115.17	107.27
2	A	401	NAI	O7N-C7N-C3N	-2.85	115.52	120.90
3	A	402	GZ0	O4'-C4'-C5'	2.84	118.42	109.33
3	A	402	GZ0	O71-C71-C61	-2.81	105.26	111.16
3	B	402	GZ0	O4'-C1'-C2'	-2.76	100.71	106.62
2	B	401	NAI	O3-PN-O2N	2.70	118.84	110.70
2	B	401	NAI	C4A-N9A-C1B	-2.70	120.32	126.63
3	B	402	GZ0	N9-C8-N7	-2.63	108.52	113.40
2	A	401	NAI	C4A-C5A-N7A	-2.58	107.63	110.58
3	A	402	GZ0	O4'-C1'-C2'	-2.55	101.16	106.62
2	A	401	NAI	C6A-C5A-N7A	2.54	136.98	132.09
3	B	402	GZ0	C31-C41-C51	2.52	115.40	109.68
2	B	401	NAI	C4A-C5A-N7A	-2.49	107.74	110.58
2	B	401	NAI	N9A-C8A-N7A	-2.44	110.48	113.94
3	A	402	GZ0	O5'-PA-O1A	2.40	118.45	108.94
2	B	401	NAI	C3N-C7N-N7N	2.37	121.88	117.67
2	A	401	NAI	C3B-C2B-C1B	-2.35	97.01	101.46
2	B	401	NAI	C1D-N1N-C6N	-2.31	115.89	120.77
3	B	402	GZ0	O51-C11-C21	2.29	115.07	110.37
2	B	401	NAI	C6A-C5A-C4A	-2.28	114.07	117.18
3	A	402	GZ0	O51-C11-C21	2.22	114.94	110.37
2	B	401	NAI	O5B-PA-O1A	2.21	117.71	108.94
2	A	401	NAI	C3N-C7N-N7N	2.18	121.54	117.67
2	B	401	NAI	O3-PA-O1A	-2.12	104.32	110.70
3	B	402	GZ0	O3A-PA-O1A	-2.11	104.37	110.70
3	B	402	GZ0	C71-C61-C51	2.09	116.21	112.05
2	B	401	NAI	C2A-N1A-C6A	2.08	122.16	118.73
3	B	402	GZ0	O21-C21-C31	2.07	115.26	110.38
2	A	401	NAI	O4D-C1D-C2D	-2.05	102.24	106.62
2	A	401	NAI	N9A-C8A-N7A	-2.03	111.06	113.94
2	A	401	NAI	O4D-C1D-N1N	-2.01	104.26	108.08

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAI	PN-O3-PA-O5B
2	A	401	NAI	C5D-O5D-PN-O3
2	A	401	NAI	C5D-O5D-PN-O2N
2	B	401	NAI	C5B-O5B-PA-O1A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	401	NAI	C5B-O5B-PA-O3
2	B	401	NAI	C5D-O5D-PN-O3
2	B	401	NAI	C5D-O5D-PN-O1N
2	B	401	NAI	C5D-O5D-PN-O2N
3	A	402	GZ0	C11-O1B-PB-O2B
3	A	402	GZ0	C11-O1B-PB-O3A
3	A	402	GZ0	O51-C51-C61-C71
3	A	402	GZ0	O51-C51-C61-O6A
3	A	402	GZ0	C5'-O5'-PA-O3A
3	A	402	GZ0	C5'-O5'-PA-O1A
3	B	402	GZ0	O51-C51-C61-C71
3	B	402	GZ0	O51-C51-C61-O6A
3	B	402	GZ0	O6A-C61-C71-O71
3	B	402	GZ0	C5'-O5'-PA-O3A
3	B	402	GZ0	C5'-O5'-PA-O1A
3	B	402	GZ0	C5'-O5'-PA-O2A
3	A	402	GZ0	O6A-C61-C71-O71
3	B	402	GZ0	C51-C61-C71-O71
3	A	402	GZ0	C51-C61-C71-O71
2	A	401	NAI	O4D-C1D-N1N-C6N
2	B	401	NAI	O4B-C4B-C5B-O5B
2	B	401	NAI	C3B-C4B-C5B-O5B
3	A	402	GZ0	PA-O3A-PB-O1B
2	A	401	NAI	PA-O3-PN-O2N
2	A	401	NAI	C5B-O5B-PA-O2A
2	A	401	NAI	C5B-O5B-PA-O3
2	A	401	NAI	C5D-O5D-PN-O1N
2	B	401	NAI	C5B-O5B-PA-O2A
3	A	402	GZ0	C5'-O5'-PA-O2A
3	B	402	GZ0	O4'-C4'-C5'-O5'
3	B	402	GZ0	O51-C11-O1B-PB
2	B	401	NAI	O4D-C1D-N1N-C6N
3	B	402	GZ0	C3'-C4'-C5'-O5'
2	A	401	NAI	O4B-C4B-C5B-O5B
2	A	401	NAI	PA-O3-PN-O1N
3	B	402	GZ0	PA-O3A-PB-O2B
2	B	401	NAI	O4D-C4D-C5D-O5D
2	B	401	NAI	PA-O3-PN-O1N
3	A	402	GZ0	PB-O3A-PA-O1A
3	A	402	GZ0	PB-O3A-PA-O2A

There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	NAI	2	0
2	A	401	NAI	3	0
3	A	402	GZ0	1	0
3	B	402	GZ0	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand NAI A 401



Ligand GZ0 A 402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/321 (94%)	-0.27	1 (0%) 90 88	10, 21, 39, 58	0
1	B	299/321 (93%)	0.15	6 (2%) 65 60	14, 32, 54, 76	0
All	All	602/642 (93%)	-0.06	7 (1%) 76 73	10, 25, 50, 76	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	129	ASP	3.1
1	B	159	GLY	3.1
1	B	225	LYS	2.8
1	B	131	MET	2.8
1	B	160	ASN	2.4
1	B	126	GLY	2.3
1	B	224	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAI	B	401	44/44	0.96	0.07	21,25,31,32	0
3	GZ0	B	402	41/41	0.96	0.07	23,26,30,34	0
3	GZ0	A	402	41/41	0.98	0.06	10,11,18,18	0
2	NAI	A	401	44/44	0.98	0.06	8,11,15,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAI B 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GZ0 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GZ0 A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAI A 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.