



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 09:02 AM EDT

PDB ID : 7VOE / pdb_00007voe
Title : Crystal structure of 5-HT2AR in complex with aripiprazole
Authors : Chen, Z.; Fan, L.; Wang, H.; Yu, J.; Lu, D.; Qi, J.; Nie, F.; Luo, Z.; Liu, Z.;
Cheng, J.; Wang, S.
Deposited on : 2021-10-13
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 5-hydroxytryptamine receptor 2A, Soluble cytochrome b562.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	356	2718	1785	429	486	18	0	0	0

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	67	GLY	-	expression tag	UNP P28223
A	68	GLY	-	expression tag	UNP P28223
A	69	THR	-	expression tag	UNP P28223
A	162	LYS	SER	engineered mutation	UNP P28223
A	164	TRP	MET	engineered mutation	UNP P28223
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1041	GLY	-	linker	UNP P0ABE7
A	1062	SER	-	linker	UNP P0ABE7
A	1063	GLY	-	linker	UNP P0ABE7
A	1064	SER	-	linker	UNP P0ABE7
A	1065	GLY	-	linker	UNP P0ABE7
A	1098	ILE	ARG	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	GLY	ARG	engineered mutation	UNP P0ABE7
A	372	ASN	SER	engineered mutation	UNP P28223

- Molecule 2 is CHOLESTEROL (CCD ID: CLR) (formula: C₂₇H₄₆O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	19	1		
2	A	1	Total	C	O	0	0
			28	27	1		
2	A	1	Total	C	O	0	0
			28	27	1		

- Molecule 3 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (CCD ID: OLC) (formula: $C_{21}H_{40}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			25	21	4		
4	A	1	Total	C	O	0	0
			15	13	2		
4	A	1	Total	C	O	0	0
			18	14	4		
4	A	1	Total	C	O	0	0
			20	16	4		

- Molecule 5 is 7-[4-[4-[2,3-bis(chloranyl)phenyl]piperazin-1-yl]butoxy]-3,4-dihydro-1H-quinolin-2-one (CCD ID: 9SC) (formula: $C_{23}H_{27}Cl_2N_3O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0	0
			30	23	2	3	2		


- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5-hydroxytryptamine receptor 2A, Soluble cytochrome b562



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	50.27Å 54.28Å 179.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.41 – 2.90 48.41 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.8 (48.41-2.90) 98.8 (48.41-2.90)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.223 , 0.244 0.227 , 0.246	Depositor DCC
R_{free} test set	589 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	72.8	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 70.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2919	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, OLC, CLR, MG, 9SC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	1/2769 (0.0%)	1.58	13/3772 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	265	ALA	C-N	8.01	1.44	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	265	ALA	CA-C-N	-7.87	110.21	120.44
1	A	265	ALA	C-N-CA	-7.87	110.21	120.44
1	A	381	THR	CA-CB-OG1	-6.71	99.53	109.60
1	A	174	TYR	CA-C-O	-5.89	114.63	120.70
1	A	313	GLN	N-CA-C	-5.62	105.24	111.36
1	A	1106	GLY	CA-C-N	-5.46	112.53	120.29
1	A	1106	GLY	C-N-CA	-5.46	112.53	120.29
1	A	265	ALA	O-C-N	5.44	127.97	122.09
1	A	1066	ASP	CA-C-N	5.37	127.33	120.56
1	A	1066	ASP	C-N-CA	5.37	127.33	120.56
1	A	313	GLN	O-C-N	5.28	128.17	122.15
1	A	368	ILE	N-CA-C	-5.22	105.45	110.72
1	A	217	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2718	0	2753	26	0
2	A	76	0	120	10	0
3	A	16	0	22	0	0
4	A	78	0	110	2	0
5	A	30	0	0	4	0
6	A	1	0	0	0	0
All	All	2919	0	3005	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:PHE:CE2	5:A:1709:9SC:CL2	2.61	0.90
1:A:342:THR:HG23	2:A:1708:CLR:H191	1.61	0.82
1:A:339:PHE:CZ	5:A:1709:9SC:CL2	2.71	0.81
2:A:1708:CLR:H183	2:A:1708:CLR:H212	1.67	0.75
1:A:342:THR:HG23	2:A:1708:CLR:C19	2.18	0.72
1:A:122:LEU:HD21	2:A:1706:CLR:H213	1.80	0.64
1:A:339:PHE:HE2	5:A:1709:9SC:CL2	2.16	0.63
1:A:349:CYS:HB2	1:A:352:SER:HB2	1.90	0.53
1:A:103:LYS:HA	1:A:106:GLN:HG2	1.92	0.52
1:A:173:ARG:NH2	1:A:318:GLU:O	2.45	0.50
1:A:336:TRP:HB3	5:A:1709:9SC:C20	2.43	0.48
1:A:354:ASN:H	2:A:1708:CLR:H231	1.77	0.48
1:A:81:THR:O	1:A:85:ILE:HG13	2.14	0.47
1:A:361:LEU:CD1	2:A:1708:CLR:H193	2.45	0.47
1:A:146:LYS:HB2	4:A:1704:OLC:H3A	1.98	0.45
1:A:223:LYS:HB3	1:A:228:LEU:HD21	1.98	0.45
1:A:240:PHE:HA	1:A:244:PHE:HB3	1.98	0.45
1:A:361:LEU:HD13	2:A:1708:CLR:H193	1.99	0.45
2:A:1706:CLR:H232	2:A:1706:CLR:H211	1.85	0.44
1:A:349:CYS:SG	1:A:349:CYS:O	2.76	0.44
1:A:202:ILE:O	1:A:206:ILE:HG13	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:206:ILE:HG21	1:A:242:SER:HB3	2.01	0.43
2:A:1706:CLR:H222	2:A:1706:CLR:H162	1.74	0.43
1:A:342:THR:CG2	2:A:1708:CLR:C19	2.92	0.42
1:A:362:LEU:HD12	1:A:362:LEU:HA	1.79	0.42
1:A:122:LEU:HD23	1:A:122:LEU:HA	1.87	0.41
1:A:163:ILE:HG21	1:A:243:PHE:HD1	1.86	0.41
1:A:1091:ALA:HA	1:A:1094:LEU:HD12	2.03	0.41
1:A:220:LYS:HE2	4:A:1703:OLC:H15	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/376 (93%)	344 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/322 (89%)	279 (97%)	9 (3%)	35	69

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	LYS
1	A	228	LEU
1	A	244	PHE
1	A	245	ILE
1	A	260	SER
1	A	1008	GLU
1	A	1019	LYS
1	A	322	CYS
1	A	362	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	178	GLN
1	A	216	GLN
1	A	1099	ASN
1	A	1103	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1PE	A	1702	-	15,15,15	0.30	0	14,14,14	0.24	0
4	OLC	A	1703	-	24,24,24	0.22	0	25,25,25	0.32	0
4	OLC	A	1705	-	17,17,24	0.42	0	18,18,25	0.48	0
4	OLC	A	1707	-	19,19,24	0.31	0	20,20,25	0.44	0
2	CLR	A	1706	-	31,31,31	0.43	0	48,48,48	0.79	0
2	CLR	A	1701	-	23,23,31	0.43	0	37,37,48	0.68	0
2	CLR	A	1708	-	31,31,31	0.47	0	48,48,48	0.91	3 (6%)
5	9SC	A	1709	-	33,33,33	0.43	0	45,45,45	0.99	3 (6%)
4	OLC	A	1704	-	14,14,24	0.46	0	14,14,25	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1PE	A	1702	-	-	6/13/13/13	-
4	OLC	A	1703	-	-	5/24/24/24	-
4	OLC	A	1705	-	-	9/17/17/24	-
4	OLC	A	1707	-	-	8/19/19/24	-
2	CLR	A	1706	-	-	3/10/68/68	0/4/4/4
2	CLR	A	1701	-	-	-	0/4/4/4
2	CLR	A	1708	-	-	0/10/68/68	0/4/4/4
5	9SC	A	1709	-	-	6/12/31/31	0/4/4/4
4	OLC	A	1704	-	-	2/12/12/24	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1709	9SC	C2-C1-CL1	-3.16	117.61	120.51
2	A	1708	CLR	C17-C13-C14	2.78	103.29	100.10
2	A	1708	CLR	C1-C2-C3	2.64	113.97	110.48
5	A	1709	9SC	C3-C2-CL2	2.12	122.07	120.07
5	A	1709	9SC	C21-C18-C17	2.05	125.32	121.05
2	A	1708	CLR	C1-C10-C9	2.01	111.40	108.74

There are no chirality outliers.

All (39) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1705	OLC	C21-C22-C24-O25
4	A	1707	OLC	C10-C11-C12-C13
4	A	1707	OLC	C2-C1-O20-C21
4	A	1707	OLC	O19-C1-O20-C21
5	A	1709	9SC	C12-C11-N2-C8
4	A	1707	OLC	O23-C22-C24-O25
5	A	1709	9SC	C12-C13-C14-O1
4	A	1703	OLC	O20-C21-C22-O23
4	A	1705	OLC	O20-C21-C22-O23
3	A	1702	1PE	C13-C23-OH3-C22
3	A	1702	1PE	C25-C15-OH6-C26
4	A	1703	OLC	O20-C21-C22-C24
4	A	1705	OLC	O20-C21-C22-C24
4	A	1704	OLC	C10-C11-C12-C13
4	A	1707	OLC	C21-C22-C24-O25
4	A	1705	OLC	O23-C22-C24-O25
4	A	1705	OLC	C2-C3-C4-C5
2	A	1706	CLR	C20-C22-C23-C24
3	A	1702	1PE	OH5-C14-C24-OH4
4	A	1707	OLC	C4-C5-C6-C7
4	A	1707	OLC	O20-C21-C22-C24
2	A	1706	CLR	C22-C23-C24-C25
4	A	1705	OLC	C4-C5-C6-C7
5	A	1709	9SC	N2-C11-C12-C13
5	A	1709	9SC	C13-C14-O1-C15
4	A	1707	OLC	O20-C21-C22-O23
5	A	1709	9SC	C12-C11-N2-C9
5	A	1709	9SC	C11-C12-C13-C14
4	A	1705	OLC	C5-C6-C7-C8
4	A	1703	OLC	C3-C4-C5-C6
4	A	1704	OLC	C7-C8-C9-C10
2	A	1706	CLR	C16-C17-C20-C22
4	A	1703	OLC	C5-C6-C7-C8
3	A	1702	1PE	OH6-C15-C25-OH5
3	A	1702	1PE	OH7-C16-C26-OH6
4	A	1705	OLC	C3-C4-C5-C6
4	A	1705	OLC	C7-C8-C9-C10
4	A	1703	OLC	C6-C7-C8-C9
3	A	1702	1PE	OH4-C13-C23-OH3




There are no ring outliers.


5 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1703	OLC	1	0
2	A	1706	CLR	3	0
2	A	1708	CLR	7	0
5	A	1709	9SC	4	0
4	A	1704	OLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/376 (94%)	-0.03	10 (2%) 55 46	43, 71, 107, 138	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	74	LYS	3.9
1	A	242	SER	3.8
1	A	1102	ILE	3.5
1	A	1006	ASN	3.4
1	A	243	PHE	3.0
1	A	84	VAL	2.8
1	A	1066	ASP	2.5
1	A	244	PHE	2.4
1	A	1037	ALA	2.3
1	A	193	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CLR	A	1708	28/28	0.66	0.21	83,120,135,153	0
5	9SC	A	1709	30/30	0.81	0.27	51,96,159,212	30
4	OLC	A	1707	20/25	0.85	0.17	72,97,149,154	0
4	OLC	A	1704	15/25	0.85	0.17	48,68,116,138	0
4	OLC	A	1703	25/25	0.86	0.23	76,94,117,122	0
3	1PE	A	1702	16/16	0.86	0.12	89,105,114,118	0
2	CLR	A	1701	20/28	0.91	0.12	82,98,114,120	0
4	OLC	A	1705	18/25	0.91	0.18	79,89,106,116	0
2	CLR	A	1706	28/28	0.93	0.16	81,104,120,127	0
6	MG	A	1710	1/1	1.00	0.01	9,9,9,9	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 9SC A 1709:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 1707:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 1704:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 1703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 1705:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CLR A 1706:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.