



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:52 AM UTC

PDB ID : 1VPE / pdb_00001vpe
Title : CRYSTALLOGRAPHIC ANALYSIS OF PHOSPHOGLYCERATE KINASE
FROM THE HYPERTHERMOPHILIC BACTERIUM THERMOTOGA
MARITIMA
Authors : Auerbach, G.; Huber, R.; Graettinger, M.; Zaiss, K.; Schurig, H.; Jaenicke, R.;
Jacob, U.
Deposited on : 1997-05-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	398	 82% 17% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOGLYCERATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	398	Total	C	N	O	S	59	0	0
			3026	1939	509	569	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	250	VAL	LEU	conflict	UNP P36204
A	394	MET	ILE	conflict	UNP P36204
A	395	ARG	ALA	conflict	UNP P36204
A	396	ILE	ASP	conflict	UNP P36204

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 4 is 3-PHOSPHOGLYCERIC ACID (CCD ID: 3PG) (formula: $\text{C}_3\text{H}_7\text{O}_7\text{P}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			11	3	7	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	226	Total 226	O 226	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PHOSPHOGLYCERATE KINASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.01Å 76.86Å 87.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.00)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.198 , 0.288	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3295	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 3PG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/3075	0.99	8/4143 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	VAL	N-CA-C	7.27	118.35	108.17
1	A	119	PHE	N-CA-C	-6.36	105.49	113.18
1	A	297	GLY	CA-C-N	6.28	126.02	119.05
1	A	297	GLY	C-N-CA	6.28	126.02	119.05
1	A	391	ILE	N-CA-C	-5.72	106.89	111.81
1	A	236	SER	N-CA-C	5.58	117.50	110.24
1	A	138	ALA	N-CA-C	5.29	116.84	108.96
1	A	223	PHE	N-CA-C	5.01	117.63	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3026	0	3141	46	0
2	A	1	0	0	0	0
3	A	31	0	13	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	11	0	4	0	0
5	A	226	0	0	9	0
All	All	3295	0	3158	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:374:THR:HG23	5:A:573:HOH:O	1.67	0.94
1:A:179:SER:HB2	5:A:504:HOH:O	1.86	0.75
1:A:319:PRO:HD2	3:A:400:ANP:H5'2	1.70	0.73
1:A:201:LYS:O	1:A:204:VAL:HG22	1.95	0.66
1:A:386:LYS:H	1:A:386:LYS:HD2	1.63	0.63
1:A:240:GLU:HA	1:A:243:ILE:HD11	1.82	0.62
1:A:204:VAL:HG23	1:A:208:LEU:HD12	1.84	0.60
1:A:48:LEU:HD21	1:A:111:VAL:HG22	1.85	0.57
1:A:322:VAL:HA	3:A:400:ANP:O2'	2.07	0.55
1:A:208:LEU:HD23	5:A:587:HOH:O	2.07	0.54
1:A:201:LYS:HB2	1:A:221:MET:HE1	1.91	0.53
1:A:141:HIS:HD2	1:A:165:SER:OG	1.91	0.53
1:A:150:HIS:H	1:A:150:HIS:CD2	2.26	0.53
1:A:381:GLU:HG3	5:A:500:HOH:O	2.07	0.52
1:A:216:LEU:HD23	1:A:216:LEU:N	2.25	0.52
1:A:302:GLU:O	1:A:306:GLN:HG3	2.09	0.52
1:A:108:GLU:HG3	5:A:530:HOH:O	2.09	0.51
1:A:320:MET:HG3	1:A:335:VAL:HG21	1.92	0.51
1:A:268:ILE:HG22	1:A:293:GLY:HA2	1.93	0.51
1:A:391:ILE:HA	1:A:394:MET:HE2	1.92	0.51
1:A:345:LYS:HG3	5:A:579:HOH:O	2.11	0.50
1:A:301:ILE:O	1:A:305:LYS:HG3	2.13	0.49
1:A:240:GLU:HA	1:A:243:ILE:CD1	2.42	0.49
1:A:101:LYS:HE3	1:A:105:GLU:OE2	2.13	0.49
1:A:224:THR:HA	1:A:238:VAL:HG13	1.95	0.49
1:A:172:GLU:O	1:A:176:LYS:HG2	2.13	0.48
1:A:78:LYS:HB3	5:A:577:HOH:O	2.14	0.48
1:A:45:LYS:HE2	5:A:557:HOH:O	2.14	0.47
1:A:158:GLY:O	1:A:161:GLN:HG2	2.15	0.47
1:A:331:GLY:O	1:A:335:VAL:HG23	2.15	0.46
1:A:319:PRO:CD	3:A:400:ANP:H5'2	2.41	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:VAL:HG21	1:A:307:LYS:HG2	1.99	0.44
1:A:68:SER:HB2	1:A:71:PHE:HD2	1.83	0.44
1:A:15:ARG:HH11	1:A:15:ARG:HG3	1.83	0.43
1:A:233:VAL:HG21	1:A:238:VAL:HG21	2.00	0.43
1:A:374:THR:O	1:A:376:GLY:N	2.49	0.43
1:A:23:ASN:HB3	1:A:36:ARG:HH11	1.84	0.43
1:A:77:ALA:HB2	1:A:113:LEU:HD23	2.01	0.42
1:A:268:ILE:HA	1:A:292:MET:O	2.19	0.42
1:A:223:PHE:CZ	1:A:293:GLY:HA3	2.55	0.41
1:A:271:LYS:HA	5:A:537:HOH:O	2.19	0.41
1:A:227:LYS:HD2	1:A:238:VAL:HG11	2.02	0.41
1:A:268:ILE:HD11	1:A:281:VAL:HG11	2.01	0.41
1:A:386:LYS:HD2	1:A:386:LYS:N	2.33	0.41
1:A:264:VAL:O	1:A:283:ILE:HG22	2.21	0.40
1:A:140:ILE:HG22	1:A:141:HIS:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/398 (100%)	379 (96%)	16 (4%)	1 (0%)	36	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	231	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	322/322 (100%)	316 (98%)	6 (2%)	50 56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	LYS
1	A	216	LEU
1	A	226	LEU
1	A	241	ASP
1	A	278	LYS
1	A	326	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	141	HIS
1	A	150	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	A	400	2	33,33,33	2.77	8 (24%)	45,52,52	2.07	11 (24%)
4	3PG	A	401	-	9,10,10	3.39	5 (55%)	11,14,14	1.54	3 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	400	2	-	6/18/38/38	0/3/3/3
4	3PG	A	401	-	-	1/10/10/10	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	400	ANP	PA-O3A	-8.68	1.50	1.59
3	A	400	ANP	PG-N3B	8.32	1.85	1.63
3	A	400	ANP	PB-O3A	-6.31	1.51	1.59
4	A	401	3PG	P-O2P	6.06	1.69	1.50
4	A	401	3PG	O1-C1	4.35	1.35	1.22
4	A	401	3PG	P-O3P	4.04	1.69	1.54
4	A	401	3PG	P-O4P	3.84	1.69	1.54
4	A	401	3PG	C2-C1	-3.31	1.47	1.52
3	A	400	ANP	PB-N3B	-3.24	1.54	1.63
3	A	400	ANP	PG-O3G	-2.68	1.49	1.56
3	A	400	ANP	PG-O1G	2.45	1.49	1.46
3	A	400	ANP	C5-N7	2.16	1.43	1.39
3	A	400	ANP	O4'-C1'	2.12	1.46	1.42

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	400	ANP	O1B-PB-N3B	-7.35	100.94	111.77
3	A	400	ANP	O4'-C1'-N9	5.99	119.59	108.09
3	A	400	ANP	O4'-C4'-C3'	4.20	113.49	105.15
3	A	400	ANP	O2A-PA-O3A	4.14	118.47	107.27
4	A	401	3PG	O2-C1-C2	3.44	120.02	112.74
3	A	400	ANP	O2G-PG-O3G	3.31	116.49	107.59
3	A	400	ANP	O2B-PB-O3A	2.74	113.78	104.64
3	A	400	ANP	O5'-C5'-C4'	-2.62	100.09	108.99
3	A	400	ANP	O1G-PG-N3B	-2.52	108.06	111.77
4	A	401	3PG	O2-C1-O1	-2.42	118.59	124.08
3	A	400	ANP	C4'-O4'-C1'	-2.37	104.23	109.47
3	A	400	ANP	O2A-PA-O5'	-2.28	97.21	107.57
4	A	401	3PG	O4P-P-O1P	2.14	112.24	106.67
3	A	400	ANP	C2'-C1'-N9	-2.04	108.24	113.30

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	400	ANP	PB-N3B-PG-O1G
3	A	400	ANP	C5'-O5'-PA-O1A
3	A	400	ANP	C5'-O5'-PA-O2A
4	A	401	3PG	O3-C2-C3-O1P
3	A	400	ANP	C5'-O5'-PA-O3A
3	A	400	ANP	C3'-C4'-C5'-O5'
3	A	400	ANP	PB-O3A-PA-O1A


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	ANP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.