



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:03 PM UTC

PDB ID : 6VP5 / pdb_00006vp5
Title : Ethylene forming enzyme (EFE) D191E variant in complex with Fe(II), L-arginine, and 2OG
Authors : Davis, K.M.; Copeland, R.A.; Boal, A.K.
Deposited on : 2020-02-01
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	 3% 91% 5% .
1	B	351	 4% 92% 5% .
1	C	351	 3% 91% 5% .
1	D	351	 3% 89% 5% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11986 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-oxoglutarate-dependent ethylene/succinate-forming enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	3	0
			2651	1689	455	492	15			
1	B	342	Total	C	N	O	S	0	5	0
			2708	1722	460	509	17			
1	C	337	Total	C	N	O	S	0	3	0
			2664	1699	456	493	16			
1	D	331	Total	C	N	O	S	0	3	0
			2601	1661	444	481	15			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P32021
A	191	GLU	ASP	engineered mutation	UNP P32021
B	0	SER	-	expression tag	UNP P32021
B	191	GLU	ASP	engineered mutation	UNP P32021
C	0	SER	-	expression tag	UNP P32021
C	191	GLU	ASP	engineered mutation	UNP P32021
D	0	SER	-	expression tag	UNP P32021
D	191	GLU	ASP	engineered mutation	UNP P32021

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		

- Molecule 3 is 2-OXOGLUTARIC ACID (CCD ID: AKG) (formula: $C_5H_6O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		
3	C	1	Total	C	O	0	0
			10	5	5		
3	D	1	Total	C	O	0	0
			10	5	5		

- Molecule 4 is ARGinine (CCD ID: ARG) (formula: $C_6H_{15}N_4O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	6	4	2		
4	B	1	Total	C	N	O	0	0
			12	6	4	2		
4	C	1	Total	C	N	O	0	0
			12	6	4	2		
4	D	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 5 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: $C_{11}H_{26}N_2O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			19	11	2	6		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	294	Total	O	0	0
			294	294		
6	B	338	Total	O	0	0
			338	338		
6	C	310	Total	O	0	0
			310	310		
6	D	309	Total	O	0	0
			309	309		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-oxoglutarate-dependent ethylene/succinate-forming enzyme

Chain A: 




- Molecule 1: 2-oxoglutarate-dependent ethylene/succinate-forming enzyme

Chain B: 




- Molecule 1: 2-oxoglutarate-dependent ethylene/succinate-forming enzyme

Chain C: 



- Molecule 1: 2-oxoglutarate-dependent ethylene/succinate-forming enzyme

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.54Å 78.50Å 195.49Å 90.00° 92.34° 90.00°	Depositor
Resolution (Å)	49.52 – 1.97 49.52 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.52-1.97) 93.8 (49.52-1.97)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 1.97Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.173 , 0.207 0.173 , 0.206	Depositor DCC
R_{free} test set	1592 reflections (1.50%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11986	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, AKG, B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/2725	0.35	0/3707
1	B	0.30	2/2786 (0.1%)	0.40	1/3791 (0.0%)
1	C	0.13	0/2741	0.34	0/3726
1	D	0.14	0/2680	0.36	0/3646
All	All	0.19	2/10932 (0.0%)	0.36	1/14870 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	44	GLN	C-O	-6.57	1.16	1.24
1	B	87	ALA	C-O	-6.09	1.15	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	89	LYS	CB-CA-C	-5.76	102.41	110.13

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2651	0	2526	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2708	0	2562	9	0
1	C	2664	0	2541	12	0
1	D	2601	0	2481	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
3	C	10	0	4	0	0
3	D	10	0	4	0	0
4	A	12	0	12	0	0
4	B	12	0	12	0	0
4	C	12	0	12	0	0
4	D	12	0	12	1	0
5	D	19	0	26	0	0
6	A	294	0	0	0	0
6	B	338	0	0	0	0
6	C	310	0	0	2	0
6	D	309	0	0	0	0
All	All	11986	0	10200	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:86:THR:HG1	4:D:403:ARG:N	1.90	0.70
1:C:337:GLU:HA	1:C:340:LYS:HD2	1.85	0.59
1:C:271:LYS:NZ	6:C:502:HOH:O	2.26	0.57
1:B:48:THR:O	1:B:52:MET:HG2	2.07	0.55
1:D:8:GLU:HG2	1:D:39:LYS:HE3	1.90	0.54
1:C:193:GLY:HA3	1:C:285:GLU:OE2	2.09	0.53
1:C:333:LEU:HD23	1:C:336:LEU:HD12	1.91	0.52
1:B:191[A]:GLU:HG2	1:B:283:PHE:CE1	2.45	0.52
1:C:191[A]:GLU:HG2	1:C:283:PHE:CE1	2.46	0.51
1:C:337:GLU:HA	1:C:340:LYS:CD	2.42	0.49
1:D:191[A]:GLU:HG2	1:D:283:PHE:CE2	2.48	0.49
1:B:193:GLY:HA3	1:B:285:GLU:OE1	2.12	0.49
1:C:271:LYS:NZ	6:C:513:HOH:O	2.42	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:47:LYS:HD2	1:C:136:GLU:HG3	1.96	0.47
1:A:193:GLY:HA3	1:A:285:GLU:OE1	2.14	0.47
1:A:191[A]:GLU:HG2	1:A:283:PHE:CE1	2.50	0.47
1:B:104:VAL:HA	1:B:109:VAL:HG11	1.97	0.47
1:B:299:SER:CB	1:D:219:ARG:HH12	2.27	0.47
1:D:91:ASP:OD1	1:D:184:ARG:NH2	2.48	0.47
1:A:20:ILE:O	1:A:24:ARG:HG3	2.16	0.46
1:A:192:TYR:O	1:A:254:ILE:HG12	2.17	0.45
1:D:45:ASP:O	1:D:49:GLN:HG2	2.19	0.43
1:D:8:GLU:CG	1:D:39:LYS:HE3	2.49	0.43
1:C:35:ILE:HD11	1:C:248:THR:HB	2.00	0.43
1:C:208:ILE:HG22	1:C:268:HIS:HB3	2.02	0.42
1:D:196:VAL:O	1:D:249:VAL:HA	2.20	0.42
1:B:285:GLU:OE2	1:B:306:TYR:OH	2.28	0.42
1:B:190:THR:HB	1:B:265:SER:OG	2.19	0.41
1:B:318:TYR:HB3	1:B:321:ARG:HB2	2.01	0.41
1:B:137:LEU:HD13	1:B:282:TYR:HB2	2.02	0.41
1:A:35:ILE:HD11	1:A:248:THR:HB	2.02	0.41
1:A:92:PHE:N	1:A:93:PRO:CD	2.84	0.41
1:D:193:GLY:HA3	1:D:285:GLU:OE1	2.21	0.41
1:A:137:LEU:HD13	1:A:282:TYR:HB2	2.03	0.40
1:A:196:VAL:O	1:A:249:VAL:HA	2.21	0.40
1:C:255:LEU:HD23	1:C:255:LEU:HA	1.82	0.40
1:C:175:PHE:HE2	1:C:186:ILE:HG13	1.86	0.40
1:A:193:GLY:HA3	1:A:285:GLU:CD	2.46	0.40
1:D:175:PHE:HE2	1:D:186:ILE:HG13	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/351 (95%)	328 (98%)	4 (1%)	2 (1%)	21	12
1	B	345/351 (98%)	338 (98%)	6 (2%)	1 (0%)	36	27
1	C	336/351 (96%)	330 (98%)	5 (2%)	1 (0%)	36	27
1	D	330/351 (94%)	322 (98%)	6 (2%)	2 (1%)	21	12
All	All	1345/1404 (96%)	1318 (98%)	21 (2%)	6 (0%)	24	20

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	GLY
1	D	205	GLY
1	C	205	GLY
1	B	205	GLY
1	D	92	PHE
1	A	92	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/301 (92%)	276 (99%)	2 (1%)	76	73
1	B	284/301 (94%)	282 (99%)	2 (1%)	76	73
1	C	280/301 (93%)	278 (99%)	2 (1%)	76	73
1	D	272/301 (90%)	270 (99%)	2 (1%)	76	73
All	All	1114/1204 (92%)	1106 (99%)	8 (1%)	84	73

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	191[A]	GLU
1	A	191[B]	GLU
1	B	191[A]	GLU
1	B	191[B]	GLU
1	C	191[A]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	191[B]	GLU
1	D	191[A]	GLU
1	D	191[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	107	GLN
1	A	157	ASN
1	B	37	GLN
1	C	37	GLN
1	D	157	ASN
1	D	233	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AKG	D	402	2	9,9,9	5.21	2 (22%)	11,11,11	1.98	3 (27%)
4	ARG	B	403	-	10,11,11	0.78	1 (10%)	9,13,13	0.94	1 (11%)
4	ARG	A	403	-	10,11,11	0.76	1 (10%)	9,13,13	0.88	1 (11%)
3	AKG	B	402	2	9,9,9	5.28	2 (22%)	11,11,11	1.88	2 (18%)
4	ARG	C	403	-	10,11,11	0.76	1 (10%)	9,13,13	0.94	1 (11%)
5	B3P	D	404	-	18,18,18	0.07	0	23,23,23	0.50	0
4	ARG	D	403	-	10,11,11	0.76	1 (10%)	9,13,13	0.92	1 (11%)
3	AKG	A	402	2	9,9,9	5.24	2 (22%)	11,11,11	2.07	3 (27%)
3	AKG	C	402	2	9,9,9	5.25	2 (22%)	11,11,11	1.88	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AKG	D	402	2	-	2/9/9/9	-
4	ARG	B	403	-	-	0/11/11/11	-
4	ARG	A	403	-	-	0/11/11/11	-
3	AKG	B	402	2	-	2/9/9/9	-
4	ARG	C	403	-	-	0/11/11/11	-
5	B3P	D	404	-	-	0/28/28/28	-
4	ARG	D	403	-	-	0/11/11/11	-
3	AKG	A	402	2	-	4/9/9/9	-
3	AKG	C	402	2	-	1/9/9/9	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	AKG	C2-C1	-15.46	1.30	1.53
3	C	402	AKG	C2-C1	-15.38	1.30	1.53
3	A	402	AKG	C2-C1	-15.35	1.30	1.53
3	D	402	AKG	C2-C1	-15.27	1.30	1.53
3	B	402	AKG	O2-C1	-2.68	1.23	1.30
3	A	402	AKG	O2-C1	-2.66	1.23	1.30
3	D	402	AKG	O2-C1	-2.64	1.23	1.30
3	C	402	AKG	O2-C1	-2.61	1.23	1.30
4	C	403	ARG	OXT-C	-2.22	1.23	1.30
4	A	403	ARG	OXT-C	-2.20	1.23	1.30
4	D	403	ARG	OXT-C	-2.20	1.23	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	403	ARG	OXT-C	-2.17	1.23	1.30

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	AKG	C3-C2-C1	4.95	124.27	115.86
3	B	402	AKG	C3-C2-C1	4.83	124.07	115.86
3	C	402	AKG	C3-C2-C1	4.77	123.95	115.86
3	D	402	AKG	C3-C2-C1	4.69	123.83	115.86
3	A	402	AKG	O1-C1-C2	-3.10	117.96	121.81
3	D	402	AKG	O1-C1-C2	-2.99	118.09	121.81
4	C	403	ARG	OXT-C-O	-2.74	117.85	124.08
4	B	403	ARG	OXT-C-O	-2.71	117.93	124.08
4	D	403	ARG	OXT-C-O	-2.69	117.98	124.08
4	A	403	ARG	OXT-C-O	-2.58	118.23	124.08
3	C	402	AKG	O1-C1-C2	-2.22	119.04	121.81
3	B	402	AKG	O1-C1-C2	-2.13	119.16	121.81
3	A	402	AKG	O2-C1-C2	2.10	119.42	113.59
3	D	402	AKG	O2-C1-C2	2.05	119.28	113.59

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	AKG	C1-C2-C3-C4
3	B	402	AKG	C1-C2-C3-C4
3	C	402	AKG	C1-C2-C3-C4
3	A	402	AKG	C3-C4-C5-O4
3	D	402	AKG	C3-C4-C5-O4
3	B	402	AKG	O5-C2-C3-C4
3	D	402	AKG	C3-C4-C5-O3
3	A	402	AKG	C3-C4-C5-O3
3	A	402	AKG	O5-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	403	ARG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	335/351 (95%)	-0.11	10 (2%)	52 62	7, 16, 36, 74	3 (0%)
1	B	342/351 (97%)	-0.17	15 (4%)	39 49	6, 14, 41, 65	5 (1%)
1	C	337/351 (96%)	-0.08	9 (2%)	56 66	7, 18, 40, 60	3 (0%)
1	D	331/351 (94%)	-0.16	10 (3%)	52 62	6, 15, 36, 55	3 (0%)
All	All	1345/1404 (95%)	-0.13	44 (3%)	49 59	6, 16, 38, 74	14 (1%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	334	ALA	5.7
1	D	334	ALA	5.5
1	A	336	LEU	5.4
1	A	298	PRO	4.8
1	A	335	HIS	4.5
1	D	105	GLY	4.1
1	B	339	LEU	4.1
1	D	298	PRO	4.0
1	A	2	THR	3.8
1	B	341	LYS	3.7
1	B	298	PRO	3.7
1	D	223	PRO	3.7
1	C	342	TYR	3.3
1	C	336	LEU	3.3
1	B	342	TYR	3.2
1	C	339	LEU	3.2
1	D	180	SER	3.1
1	C	333	LEU	3.0
1	B	57	GLN	2.9
1	B	338	ASP	2.9
1	C	301	ASN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	331	ASN	2.8
1	B	299	SER	2.7
1	B	334	ALA	2.6
1	B	331	ASN	2.6
1	D	331	ASN	2.6
1	B	2	THR	2.6
1	A	300	ALA	2.5
1	B	223	PRO	2.5
1	D	333	LEU	2.4
1	B	110	LYS	2.4
1	B	111	ALA	2.4
1	C	334	ALA	2.4
1	B	1	MET	2.4
1	A	301	ASN	2.3
1	C	335	HIS	2.3
1	A	223	PRO	2.2
1	C	341	LYS	2.2
1	A	92	PHE	2.2
1	D	3	ASN	2.1
1	D	301	ASN	2.1
1	D	291	SER	2.1
1	B	336	LEU	2.1
1	A	333	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	B3P	D	404	19/19	0.87	0.14	16,24,39,39	19
3	AKG	C	402	10/10	0.93	0.10	17,20,29,30	0
3	AKG	A	402	10/10	0.94	0.08	17,22,25,26	0
4	ARG	B	403	12/12	0.94	0.07	11,15,17,17	0
4	ARG	C	403	12/12	0.94	0.07	15,17,21,22	0
3	AKG	B	402	10/10	0.94	0.08	14,19,25,26	0
4	ARG	A	403	12/12	0.95	0.07	11,15,19,19	0
3	AKG	D	402	10/10	0.95	0.08	14,20,24,24	0
4	ARG	D	403	12/12	0.97	0.06	10,13,16,16	0
2	FE	B	401	1/1	1.00	0.01	10,10,10,10	0
2	FE	C	401	1/1	1.00	0.01	15,15,15,15	0
2	FE	D	401	1/1	1.00	0.03	9,9,9,9	0
2	FE	A	401	1/1	1.00	0.02	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.