



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 08:50 AM UTC

PDB ID : 8VPK / pdb_00008vpk
EMDB ID : EMD-43409
Title : Structure of Mycobacterium smegmatis 50S ribosomal subunit bound to HflX and erythromycin:50S-HflX-B-Ery
Authors : Majumdar, S.; Koripella, R.K.; Sharma, M.R.; Manjari, S.R.; Banavali, N.K.; Agrawal, R.K.
Deposited on : 2024-01-16
Resolution : 2.63 Å (reported)
Based on initial models : 5O61, 6DZI

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.














Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	8888 (2.13 - 3.13)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	2	61	
2	3	24	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	4	470	
4	A	3120	
5	B	118	
6	C	278	
7	D	217	
8	E	215	
9	F	187	
10	G	179	
11	H	151	
12	I	175	
13	J	142	
14	K	147	
15	L	122	
16	M	147	
17	N	138	
18	O	199	
19	P	127	
20	Q	113	
21	R	129	
22	S	103	
23	T	153	
24	U	100	
25	V	105	
26	W	215	
27	X	88	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
28	Y	64	
29	Z	77	
30	b	57	
31	c	55	
32	d	47	
33	e	64	
34	f	37	
35	g	75	

2 Entry composition

There are 37 unique types of molecules in this entry. The entry contains 96925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	2	59	Total	C	N	O	0	0
			474	292	95	87		

- Molecule 2 is a protein called 50S Ribosomal Protein L37.

Mol	Chain	Residues	Atoms				AltConf	Trace
2	3	23	Total	C	N	O	0	0
			189	111	50	28		

- Molecule 3 is a protein called GTPase HflX.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	4	466	Total	C	N	O	S	0	0
			3509	2165	649	688	7		

- Molecule 4 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	A	2948	Total	C	N	O	P	0	0
			63320	28221	11648	20503	2948		

- Molecule 5 is a RNA chain called 5S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	B	118	Total	C	N	O	P	0	0
			2522	1126	468	810	118		

- Molecule 6 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	C	275	Total	C	N	O	S	0	0
			2110	1298	438	370	4		

- Molecule 7 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	D	214	Total	C	N	O	S	0	0
			1587	982	310	290	5		

- Molecule 8 is a protein called 50S Ribosomal Protein L4.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	E	209	Total	C	N	O	S	0	0
			1569	969	295	303	2		

- Molecule 9 is a protein called 50S Ribosomal Protein L5.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	F	182	Total	C	N	O	S	0	0
			1445	907	271	261	6		

- Molecule 10 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	G	176	Total	C	N	O	S	0	0
			1348	845	249	253	1		

- Molecule 11 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	H	151	Total	C	N	O	S	0	0
			1018	635	188	194	1		

- Molecule 12 is a protein called 50S ribosomal protein L10.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	I	126	Total	C	N	O	S	0	0
			918	580	156	180	2		

- Molecule 13 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	J	132	Total	C	N	O	S	0	0
			981	620	174	184	3		

- Molecule 14 is a protein called 50S Ribosomal Protein L13.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	K	146	Total	C	N	O	S	0	0
			1130	722	207	200	1		

- Molecule 15 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	L	122	Total	C	N	O	S	0	0
			938	586	179	170	3		

- Molecule 16 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	M	145	Total	C	N	O	S	0	0
			1078	676	205	194	3		

- Molecule 17 is a protein called Large ribosomal subunit protein uL16.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	N	136	Total	C	N	O	S	0	0
			1092	690	213	187	2		

- Molecule 18 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	O	118	Total	C	N	O	S	0	0
			928	583	180	163	2		

- Molecule 19 is a protein called 50S Ribosomal Protein L18.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	P	126	Total	C	N	O	S	0	0
			956	586	199	171			

- Molecule 20 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	Q	113	Total	C	N	O	S	0	0
			907	570	171	165	1		

- Molecule 21 is a protein called 50S Ribosomal Protein L20.

Mol	Chain	Residues	Atoms				AltConf	Trace
21	R	124	Total	C	N	O	0	0
			988	613	203	172		

- Molecule 22 is a protein called 50S Ribosomal Protein L21.

Mol	Chain	Residues	Atoms				AltConf	Trace
22	S	100	Total	C	N	O	0	0
			754	478	137	139		

- Molecule 23 is a protein called 50S Ribosomal Protein L22.

Mol	Chain	Residues	Atoms				AltConf	Trace
23	T	114	Total	C	N	O	0	0
			873	543	171	159		

- Molecule 24 is a protein called 50S Ribosomal Protein L23.

Mol	Chain	Residues	Atoms				AltConf	Trace
24	U	97	Total	C	N	O	0	0
			756	479	138	139		

- Molecule 25 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	V	97	Total	C	N	O	S	0	0
			732	456	137	137	2		

- Molecule 26 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms				AltConf	Trace
26	W	99	Total	C	N	O	0	0
			763	469	154	140		

- Molecule 27 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms				AltConf	Trace
27	X	79	Total	C	N	O	0	0
			586	361	123	102		

- Molecule 28 is a protein called 50S Ribosomal Protein L28.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	Y	63	Total	C	N	O	S	0	0
			470	283	103	80	4		

- Molecule 29 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	Z	64	Total	C	N	O	S	0	0
			531	324	103	103	1		

- Molecule 30 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	b	54	Total	C	N	O	S	0	0
			423	260	93	69	1		

- Molecule 31 is a protein called 50S Ribosomal Protein L33.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	c	49	Total	C	N	O	S	0	0
			405	248	82	71	4		

- Molecule 32 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	d	46	Total	C	N	O	S	0	0
			377	225	97	54	1		

- Molecule 33 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms				AltConf	Trace
33	e	63	Total	C	N	O	0	0
			502	302	115	85		


- Molecule 34 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	f	37	Total	C	N	O	S	0	0
			299	181	66	47	5		


- Molecule 35 is a protein called 50S Ribosomal Protein L31.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	g	48	Total	C	N	O	S	0	0
			364	225	63	71	5		



- Molecule 36 is PHOSPHOMETHYLPHOSPHONIC ACID GUANYLATE ESTER (CCD ID: GCP) (formula: $C_{11}H_{18}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
37	A	1	51	37	1	13	0




• Molecule 4: 23S ribosomal RNA













• Molecule 10: 50S ribosomal protein L6




• Molecule 11: 50S ribosomal protein L9




• Molecule 12: 50S ribosomal protein L10




• Molecule 13: 50S ribosomal protein L11




- Molecule 14: 50S Ribosomal Protein L13




- Molecule 15: 50S ribosomal protein L14



- Molecule 16: 50S ribosomal protein L15




- Molecule 17: Large ribosomal subunit protein uL16




- Molecule 18: 50S ribosomal protein L17








• Molecule 19: 50S Ribosomal Protein L18





• Molecule 20: 50S ribosomal protein L19




• Molecule 21: 50S Ribosomal Protein L20



• Molecule 22: 50S Ribosomal Protein L21



• Molecule 23: 50S Ribosomal Protein L22



- Molecule 24: 50S Ribosomal Protein L23

Chain U:  79% 18%






- Molecule 25: 50S ribosomal protein L24

Chain V:  6% 78% 14% 8%



- Molecule 26: 50S ribosomal protein L25

Chain W:  6% 37% 9% 54%




- Molecule 27: 50S ribosomal protein L27

Chain X:  81% 9% 10%



- Molecule 28: 50S Ribosomal Protein L28

Chain Y:  5% 83% 16%




- Molecule 29: 50S ribosomal protein L29

Chain Z:  66% 17% 17%




- Molecule 30: 50S ribosomal protein L32

Chain b:  89% 5% 5%




- Molecule 31: 50S Ribosomal Protein L33

Chain c:  73% 16% 11%



- Molecule 32: 50S ribosomal protein L34

Chain d:  79% 19% .



- Molecule 33: 50S ribosomal protein L35

Chain e:  72% 27% .




- Molecule 34: 50S ribosomal protein L36

Chain f:  84% 16%



- Molecule 35: 50S Ribosomal Protein L31

Chain g:  41% 49% 15% 36%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	113394	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	62.87	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	81000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.749	Depositor
Minimum map value	-0.854	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.059	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	433.152, 433.152, 433.152	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.846, 0.846, 0.846	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ERY, GCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	2	0.18	0/477	0.30	0/640
2	3	0.22	0/191	0.38	0/247
3	4	0.16	0/3550	0.41	0/4811
4	A	0.21	0/70902	0.31	0/110629
5	B	0.16	0/2821	0.29	0/4396
6	C	0.18	0/2153	0.35	0/2895
7	D	0.45	1/1609 (0.1%)	0.53	3/2165 (0.1%)
8	E	0.18	0/1592	0.32	0/2153
9	F	0.14	0/1467	0.35	0/1973
10	G	0.18	0/1369	0.40	0/1848
11	H	0.16	0/1027	0.34	0/1398
12	I	0.18	0/925	0.41	0/1246
13	J	0.15	0/997	0.38	0/1352
14	K	0.19	0/1157	0.32	0/1567
15	L	0.19	0/946	0.31	0/1268
16	M	0.20	0/1091	0.40	0/1457
17	N	0.20	0/1118	0.34	0/1506
18	O	0.19	0/945	0.36	0/1267
19	P	0.17	0/966	0.36	0/1298
20	Q	0.18	0/921	0.35	0/1236
21	R	0.20	0/1000	0.30	0/1341
22	S	0.22	0/764	0.33	0/1030
23	T	0.19	0/887	0.36	0/1204
24	U	0.17	0/766	0.34	0/1030
25	V	0.16	0/738	0.28	0/987
26	W	0.15	0/773	0.28	0/1048
27	X	0.20	0/595	0.35	0/798
28	Y	0.22	0/478	0.37	0/641
29	Z	0.17	0/534	0.28	0/713
30	b	0.19	0/427	0.31	0/572
31	c	0.17	0/413	0.31	0/553
32	d	0.19	0/380	0.28	0/500

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	e	0.18	0/507	0.30	0/672
34	f	0.21	0/303	0.30	0/401
35	g	0.16	0/372	0.33	0/503
All	All	0.20	1/105161 (0.0%)	0.33	3/157345 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	D	158	GLY	C-O	-6.12	1.15	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	156	THR	CA-C-N	-8.37	109.38	119.84
7	D	156	THR	C-N-CA	-8.37	109.38	119.84
7	D	156	THR	N-CA-C	6.29	122.09	113.16

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	474	0	500	10	0
2	3	189	0	205	2	0
3	4	3509	0	3557	126	0
4	A	63320	0	31857	1159	0
5	B	2522	0	1285	49	0
6	C	2110	0	2165	47	0
7	D	1587	0	1630	38	0
8	E	1569	0	1607	31	0
9	F	1445	0	1476	47	0
10	G	1348	0	1399	41	0
11	H	1018	0	988	14	0
12	I	918	0	959	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	J	981	0	1015	36	0
14	K	1130	0	1167	11	0
15	L	938	0	1000	17	0
16	M	1078	0	1151	35	0
17	N	1092	0	1128	17	0
18	O	928	0	972	19	0
19	P	956	0	991	36	0
20	Q	907	0	938	19	0
21	R	988	0	1038	24	0
22	S	754	0	802	11	0
23	T	873	0	909	16	0
24	U	756	0	802	14	0
25	V	732	0	782	9	0
26	W	763	0	784	13	0
27	X	586	0	601	7	0
28	Y	470	0	484	9	0
29	Z	531	0	541	13	0
30	b	423	0	463	5	0
31	c	405	0	409	7	0
32	d	377	0	411	7	0
33	e	502	0	541	15	0
34	f	299	0	324	6	0
35	g	364	0	352	10	0
36	4	32	0	14	4	0
37	A	51	0	67	6	0
All	All	96925	0	65314	1720	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 1720 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2600:A:O2'	4:A:2601:A:O5'	1.74	1.02
4:A:369:G:O2'	4:A:371:G:OP2	1.77	1.00
4:A:1496:G:O2'	4:A:1790:A:N6	1.95	0.98
4:A:2505:C:O2'	4:A:2506:G:O5'	1.82	0.98
4:A:133:G:O2'	4:A:134:A:O5'	1.83	0.97

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2	57/61 (93%)	56 (98%)	1 (2%)	0	100	100
2	3	21/24 (88%)	21 (100%)	0	0	100	100
3	4	464/470 (99%)	395 (85%)	66 (14%)	3 (1%)	21	31
6	C	273/278 (98%)	254 (93%)	19 (7%)	0	100	100
7	D	212/217 (98%)	203 (96%)	9 (4%)	0	100	100
8	E	207/215 (96%)	200 (97%)	7 (3%)	0	100	100
9	F	180/187 (96%)	163 (91%)	17 (9%)	0	100	100
10	G	174/179 (97%)	152 (87%)	22 (13%)	0	100	100
11	H	149/151 (99%)	133 (89%)	16 (11%)	0	100	100
12	I	124/175 (71%)	102 (82%)	22 (18%)	0	100	100
13	J	130/142 (92%)	105 (81%)	25 (19%)	0	100	100
14	K	144/147 (98%)	141 (98%)	3 (2%)	0	100	100
15	L	120/122 (98%)	118 (98%)	2 (2%)	0	100	100
16	M	143/147 (97%)	133 (93%)	10 (7%)	0	100	100
17	N	134/138 (97%)	132 (98%)	2 (2%)	0	100	100
18	O	116/199 (58%)	111 (96%)	5 (4%)	0	100	100
19	P	124/127 (98%)	121 (98%)	3 (2%)	0	100	100
20	Q	111/113 (98%)	105 (95%)	6 (5%)	0	100	100
21	R	122/129 (95%)	121 (99%)	1 (1%)	0	100	100
22	S	98/103 (95%)	95 (97%)	3 (3%)	0	100	100
23	T	112/153 (73%)	109 (97%)	3 (3%)	0	100	100
24	U	95/100 (95%)	90 (95%)	5 (5%)	0	100	100
25	V	93/105 (89%)	88 (95%)	5 (5%)	0	100	100
26	W	97/215 (45%)	91 (94%)	6 (6%)	0	100	100
27	X	77/88 (88%)	73 (95%)	4 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
28	Y	61/64 (95%)	57 (93%)	4 (7%)	0	100	100
29	Z	62/77 (80%)	62 (100%)	0	0	100	100
30	b	52/57 (91%)	52 (100%)	0	0	100	100
31	c	47/55 (86%)	44 (94%)	3 (6%)	0	100	100
32	d	44/47 (94%)	44 (100%)	0	0	100	100
33	e	61/64 (95%)	61 (100%)	0	0	100	100
34	f	35/37 (95%)	35 (100%)	0	0	100	100
35	g	46/75 (61%)	44 (96%)	2 (4%)	0	100	100
All	All	3985/4461 (89%)	3711 (93%)	271 (7%)	3 (0%)	49	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	4	325	ALA
3	4	15	LEU
3	4	17	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	52/54 (96%)	52 (100%)	0	100	100
2	3	18/19 (95%)	18 (100%)	0	100	100
3	4	366/372 (98%)	366 (100%)	0	100	100
6	C	215/218 (99%)	215 (100%)	0	100	100
7	D	160/163 (98%)	159 (99%)	1 (1%)	78	87
8	E	169/173 (98%)	169 (100%)	0	100	100
9	F	151/156 (97%)	151 (100%)	0	100	100
10	G	148/150 (99%)	148 (100%)	0	100	100
11	H	90/116 (78%)	90 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
12	I	89/120 (74%)	89 (100%)	0	100	100
13	J	101/108 (94%)	101 (100%)	0	100	100
14	K	119/120 (99%)	119 (100%)	0	100	100
15	L	100/100 (100%)	100 (100%)	0	100	100
16	M	112/114 (98%)	112 (100%)	0	100	100
17	N	114/116 (98%)	114 (100%)	0	100	100
18	O	97/158 (61%)	97 (100%)	0	100	100
19	P	93/94 (99%)	93 (100%)	0	100	100
20	Q	100/100 (100%)	99 (99%)	1 (1%)	68	80
21	R	97/99 (98%)	97 (100%)	0	100	100
22	S	81/83 (98%)	81 (100%)	0	100	100
23	T	90/117 (77%)	90 (100%)	0	100	100
24	U	83/85 (98%)	83 (100%)	0	100	100
25	V	81/86 (94%)	81 (100%)	0	100	100
26	W	80/168 (48%)	80 (100%)	0	100	100
27	X	58/63 (92%)	58 (100%)	0	100	100
28	Y	50/51 (98%)	50 (100%)	0	100	100
29	Z	58/66 (88%)	58 (100%)	0	100	100
30	b	43/46 (94%)	43 (100%)	0	100	100
31	c	47/52 (90%)	47 (100%)	0	100	100
32	d	35/36 (97%)	35 (100%)	0	100	100
33	e	53/54 (98%)	53 (100%)	0	100	100
34	f	35/35 (100%)	35 (100%)	0	100	100
35	g	43/63 (68%)	43 (100%)	0	100	100
All	All	3228/3555 (91%)	3226 (100%)	2 (0%)	87	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	D	156	THR
20	Q	79	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
24	U	41	GLN
29	Z	43	ASN
34	f	4	ASN
33	e	28	ASN
6	C	143	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	A	2945/3120 (94%)	911 (30%)	54 (1%)
5	B	117/118 (99%)	27 (23%)	1 (0%)
All	All	3062/3238 (94%)	938 (30%)	55 (1%)

5 of 938 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	A	3	A
4	A	7	U
4	A	33	G
4	A	52	G
4	A	60	A

5 of 55 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	A	1208	U
4	A	2168	U
5	B	10	G
4	A	2975	G
4	A	1285	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
37	ERY	A	3201	-	53,53,53	1.42	9 (16%)	82,82,82	2.22	30 (36%)
36	GCP	4	501	-	32,34,34	1.33	5 (15%)	49,54,54	1.83	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
37	ERY	A	3201	-	-	12/72/107/107	0/3/3/3
36	GCP	4	501	-	-	2/19/38/38	0/3/3/3

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
37	A	3201	ERY	O10-C6	-4.05	1.38	1.44
37	A	3201	ERY	O2-C13	-3.32	1.40	1.46
37	A	3201	ERY	O2-C1	3.19	1.41	1.34
36	4	501	GCP	C5-C4	2.89	1.46	1.38
37	A	3201	ERY	C10-C9	-2.84	1.48	1.52

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
37	A	3201	ERY	O5-C16-C17	9.04	116.97	103.86
36	4	501	GCP	C5-C4-N3	-5.90	119.00	128.39
37	A	3201	ERY	O5-C16-C15	-5.49	104.50	112.95
37	A	3201	ERY	O3-C3-C4	5.14	114.30	108.23
36	4	501	GCP	C2-N3-C4	4.92	120.77	112.30

There are no chirality outliers.

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
37	A	3201	ERY	O13-C12-C13-O2
37	A	3201	ERY	O13-C12-C13-C36
37	A	3201	ERY	C17-C16-O5-C20
36	4	501	GCP	O4'-C4'-C5'-O5'
37	A	3201	ERY	C35-C12-C13-O2

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
37	A	3201	ERY	6	0
36	4	501	GCP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-43409. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 252




Y Index: 253




Z Index: 253


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 624 nm³; this corresponds to an approximate mass of 564 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.380 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.380 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.63	-	-
Author-provided FSC curve	2.63	2.96	2.67
Unmasked-calculated*	3.62	6.43	3.75

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.62 differs from the reported value 2.63 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-43409 and PDB model 8VPK. Per-residue inclusion information can be found in section [3](#) on page [12](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8140	 0.5300
2	 0.9300	 0.6410
3	 0.9390	 0.6360
4	 0.5770	 0.4460
A	 0.8370	 0.5260
B	 0.7490	 0.4250
C	 0.9000	 0.6130
D	 0.9200	 0.6250
E	 0.8890	 0.6120
F	 0.5250	 0.4180
G	 0.7030	 0.4970
H	 0.5310	 0.4640
I	 0.1820	 0.2760
J	 0.1650	 0.2540
K	 0.9390	 0.6320
L	 0.9040	 0.6170
M	 0.8830	 0.6010
N	 0.9180	 0.6220
O	 0.9360	 0.6260
P	 0.8010	 0.5330
Q	 0.8300	 0.5890
R	 0.9480	 0.6360
S	 0.9440	 0.6390
T	 0.9300	 0.6300
U	 0.8650	 0.5960
V	 0.8070	 0.5690
W	 0.7450	 0.5410
X	 0.9310	 0.6370
Y	 0.9140	 0.6090
Z	 0.8390	 0.5970
b	 0.9030	 0.6170
c	 0.8730	 0.6020
d	 0.9570	 0.6370
e	 0.9690	 0.6370
f	 0.9440	 0.6310
g	 0.3530	 0.3220

