



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:48 PM UTC

PDB ID : 1VQ0 / pdb_00001vq0
Title : Crystal structure of 33 kDa chaperonin (Heat shock protein 33 homolog) (HSP33) (TM1394) from *Thermotoga maritima* at 2.20 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2004-11-30
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	8% 83% 12% • •
1	B	302	14% 87% 9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNL	A	302	-	-	X	-
4	UNL	B	302	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 33 kDa chaperonin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2279	1458	387	425	9			
1	B	290	Total	C	N	O	S	0	1	0
			2295	1467	388	431	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9X1B4
A	-10	GLY	-	expression tag	UNP Q9X1B4
A	-9	SER	-	expression tag	UNP Q9X1B4
A	-8	ASP	-	expression tag	UNP Q9X1B4
A	-7	LYS	-	expression tag	UNP Q9X1B4
A	-6	ILE	-	expression tag	UNP Q9X1B4
A	-5	HIS	-	expression tag	UNP Q9X1B4
A	-4	HIS	-	expression tag	UNP Q9X1B4
A	-3	HIS	-	expression tag	UNP Q9X1B4
A	-2	HIS	-	expression tag	UNP Q9X1B4
A	-1	HIS	-	expression tag	UNP Q9X1B4
A	0	HIS	-	expression tag	UNP Q9X1B4
B	-11	MET	-	expression tag	UNP Q9X1B4
B	-10	GLY	-	expression tag	UNP Q9X1B4
B	-9	SER	-	expression tag	UNP Q9X1B4
B	-8	ASP	-	expression tag	UNP Q9X1B4
B	-7	LYS	-	expression tag	UNP Q9X1B4
B	-6	ILE	-	expression tag	UNP Q9X1B4
B	-5	HIS	-	expression tag	UNP Q9X1B4
B	-4	HIS	-	expression tag	UNP Q9X1B4
B	-3	HIS	-	expression tag	UNP Q9X1B4
B	-2	HIS	-	expression tag	UNP Q9X1B4
B	-1	HIS	-	expression tag	UNP Q9X1B4
B	0	HIS	-	expression tag	UNP Q9X1B4

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0


- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0
3	B	1	Total 1	Cl 1	0	0

- Molecule 4 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 13	O 13	0	0
4	B	1	Total 13	O 13	0	0

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	130	Total O 130 130	0	0
6	B	143	Total O 143 143	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 33 kDa chaperonin



• Molecule 1: 33 kDa chaperonin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.21Å 101.67Å 113.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.62 – 2.20 49.62 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.0 (49.62-2.20) 97.0 (49.62-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.198 , 0.241 0.208 , 0.250	Depositor DCC
R_{free} test set	2228 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	32.4	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4889	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7024e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, UNL, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	2/2319 (0.1%)	1.06	3/3133 (0.1%)
1	B	1.07	1/2340 (0.0%)	1.01	1/3159 (0.0%)
All	All	1.09	3/4659 (0.1%)	1.03	4/6292 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	7	MET	SD-CE	-7.67	1.60	1.79
1	B	43	ALA	CA-CB	-6.00	1.44	1.53
1	A	150	ILE	C-O	5.51	1.29	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	MET	N-CA-C	7.78	121.90	108.76
1	A	117	ARG	NE-CZ-NH1	6.37	127.86	121.50
1	A	214	ILE	N-CA-C	5.39	115.60	110.42
1	B	146	VAL	N-CA-C	5.13	115.80	110.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2279	0	2291	24	0
1	B	2295	0	2312	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	0	0
4	A	13	0	0	6	0
4	B	13	0	0	2	0
5	A	4	0	6	1	0
5	B	8	0	12	0	0
6	A	130	0	0	2	1
6	B	143	0	0	2	1
All	All	4889	0	4621	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:302:UNL:O8	4:B:302:UNL:O9	1.53	1.26
4:A:302:UNL:O1	4:A:302:UNL:O2	1.52	1.26
4:A:302:UNL:O2	4:A:302:UNL:O3	1.52	1.26
4:A:302:UNL:O8	4:A:302:UNL:O9	1.54	1.25
4:A:302:UNL:O12	4:A:302:UNL:O11	1.54	1.24
4:A:302:UNL:O5	4:A:302:UNL:O6	1.55	1.21
1:B:58:GLU:OE2	1:B:117:ARG:HD2	1.51	1.10
4:A:302:UNL:O2	4:A:302:UNL:O5	1.83	0.97
4:B:302:UNL:O2	4:B:302:UNL:O5	1.83	0.97
1:A:58:GLU:OE2	1:A:117:ARG:HD2	1.68	0.93
1:B:58:GLU:OE2	1:B:117:ARG:CD	2.29	0.81
1:A:58:GLU:OE2	1:A:117:ARG:CD	2.30	0.78
1:B:-2:HIS:CD2	1:B:0:HIS:CD2	2.82	0.67
1:A:71:ARG:HG2	6:A:403:HOH:O	1.95	0.66
1:B:42:ARG:HD3	6:B:388:HOH:O	1.94	0.66
1:A:42:ARG:HD3	6:A:379:HOH:O	1.96	0.65
1:B:241:ALA:O	1:B:244:VAL:HG22	1.98	0.63
1:A:15:SER:HB2	1:A:53:TRP:CH2	2.34	0.63
1:A:241:ALA:O	1:A:244:VAL:HG22	1.99	0.63
1:B:15:SER:HB2	1:B:53:TRP:CH2	2.36	0.61
1:B:78:THR:HB	1:B:80[A]:GLU:OE1	2.01	0.59
1:B:-2:HIS:NE2	1:B:0:HIS:CD2	2.70	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:ARG:HD3	1:A:202:PHE:CD2	2.38	0.58
1:B:-2:HIS:HD2	1:B:0:HIS:H	1.52	0.57
1:B:18:ARG:HD3	1:B:202:PHE:CD2	2.40	0.57
1:A:167:ILE:HD11	1:A:203:GLN:HG3	1.87	0.56
1:B:167:ILE:HD11	1:B:203:GLN:HG3	1.87	0.56
1:A:-2:HIS:HD2	1:A:0:HIS:H	1.58	0.52
1:A:113:LEU:C	1:A:113:LEU:HD23	2.34	0.52
1:B:71:ARG:HG2	6:B:393:HOH:O	2.10	0.51
1:B:-1:HIS:HB3	1:B:20:ARG:HG2	1.94	0.49
1:A:1:MET:HE2	1:A:224:THR:HB	1.94	0.49
1:B:78:THR:CB	1:B:80[A]:GLU:OE1	2.61	0.49
1:A:6:THR:O	1:A:7:MET:HE2	2.13	0.49
1:A:58:GLU:OE2	1:A:117:ARG:HD3	2.12	0.48
1:A:148:GLU:OE2	5:A:303:EDO:O1	2.31	0.47
1:B:-2:HIS:NE2	1:B:0:HIS:NE2	2.62	0.47
1:B:113:LEU:C	1:B:113:LEU:HD23	2.40	0.46
1:A:20:ARG:HA	1:A:168:ALA:HB3	1.98	0.46
1:A:55:ALA:HB3	1:A:119:LEU:CD2	2.46	0.46
1:A:0:HIS:HE1	1:A:226:GLU:OE2	1.99	0.45
1:A:123:THR:HB	1:A:124:PRO:HD2	1.99	0.44
1:A:7:MET:HE1	1:A:220:GLY:H	1.82	0.44
1:B:20:ARG:HA	1:B:168:ALA:HB3	2.01	0.43
1:B:247:LYS:O	1:B:251:GLU:HG2	2.19	0.43
1:A:285:VAL:HG12	1:A:285:VAL:O	2.20	0.42
1:A:60:TRP:HD1	1:A:117:ARG:HD3	1.84	0.42
1:A:0:HIS:HB2	1:A:227:ILE:O	2.19	0.41
1:A:18:ARG:HD2	3:A:301:CL:CL	2.57	0.41
1:B:13:ARG:HD3	1:B:176:ILE:HG21	2.03	0.41
1:A:160:VAL:O	1:A:160:VAL:HG13	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:313:HOH:O	6:B:327:HOH:O[3_545]	0.12	2.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/302 (95%)	278 (96%)	10 (4%)	0	100	100
1	B	289/302 (96%)	278 (96%)	11 (4%)	0	100	100
All	All	577/604 (96%)	556 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/259 (94%)	238 (98%)	5 (2%)	47	63
1	B	247/259 (95%)	244 (99%)	3 (1%)	63	78
All	All	490/518 (95%)	482 (98%)	8 (2%)	55	71

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	91	LYS
1	A	140	LEU
1	A	213	ARG
1	A	228	LYS
1	B	15	SER
1	B	91	LYS
1	B	228	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	0	HIS
1	B	0	HIS
1	B	10	HIS
1	B	110	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic and 2 are unknown - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	304	-	3,3,3	0.59	0	2,2,2	0.10	0
5	EDO	B	303	-	3,3,3	0.34	0	2,2,2	0.42	0
5	EDO	A	303	-	3,3,3	0.50	0	2,2,2	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	304	-	-	0/1/1/1	-
5	EDO	B	303	-	-	0/1/1/1	-
5	EDO	A	303	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	303	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/302 (96%)	0.79	24 (8%) 17 15	25, 35, 52, 65	0
1	B	290/302 (96%)	1.03	41 (14%) 6 4	25, 36, 53, 64	1 (0%)
All	All	580/604 (96%)	0.91	65 (11%) 10 8	25, 36, 53, 65	1 (0%)

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285	VAL	5.6
1	B	285	VAL	5.3
1	B	221	PHE	4.8
1	A	287	ASP	4.8
1	A	1	MET	4.4
1	A	283	PHE	4.4
1	B	216	GLY	4.0
1	B	233	CYS	3.9
1	A	-2	HIS	3.9
1	A	266	CYS	3.8
1	A	233	CYS	3.8
1	B	1	MET	3.6
1	B	266	CYS	3.6
1	B	-2	HIS	3.5
1	A	286	ASP	3.3
1	B	283	PHE	3.2
1	B	119	LEU	3.2
1	B	120	GLY	3.1
1	A	265	TRP	3.1
1	B	268	THR	3.0
1	A	263	CYS	2.9
1	A	237	LYS	2.9
1	B	286	ASP	2.9
1	A	234	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	250	LEU	2.9
1	B	248	LYS	2.8
1	A	183	GLU	2.8
1	B	252	ASP	2.7
1	B	219	VAL	2.6
1	B	0	HIS	2.6
1	B	121	LEU	2.6
1	B	265	TRP	2.5
1	B	263	CYS	2.4
1	A	255	LYS	2.4
1	B	-1	HIS	2.4
1	B	209	ASP	2.4
1	B	182	GLN	2.4
1	A	268	THR	2.4
1	B	234	ASN	2.3
1	B	112	VAL	2.3
1	A	282	LYS	2.3
1	A	-1	HIS	2.3
1	B	203	GLN	2.3
1	B	255	LYS	2.3
1	B	126	VAL	2.3
1	B	236	GLU	2.2
1	B	287	ASP	2.2
1	B	181	GLU	2.2
1	B	199	SER	2.2
1	B	206	GLU	2.1
1	A	269	ARG	2.1
1	B	273	SER	2.1
1	B	122	LYS	2.1
1	B	269	ARG	2.1
1	B	200	LYS	2.0
1	A	199	SER	2.0
1	B	251	GLU	2.0
1	A	243	LEU	2.0
1	A	250	LEU	2.0
1	A	284	LYS	2.0
1	B	31	ALA	2.0
1	A	281	LEU	2.0
1	A	248	LYS	2.0
1	B	258	LYS	2.0
1	B	284	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	B	304	4/4	0.78	0.17	42,42,45,46	0
4	UNL	A	302	13/-	0.82	0.15	24,29,33,51	0
4	UNL	B	302	13/-	0.83	0.14	24,27,32,33	0
5	EDO	B	303	4/4	0.86	0.16	35,36,38,41	0
3	CL	B	301	1/1	0.90	0.17	40,40,40,40	0
3	CL	A	301	1/1	0.92	0.13	41,41,41,41	0
5	EDO	A	303	4/4	0.93	0.10	34,35,36,37	0
2	ZN	B	300	1/1	0.99	0.03	41,41,41,41	0
2	ZN	A	300	1/1	0.99	0.04	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.