



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:26 PM UTC

PDB ID : 1VQ1 / pdb_00001vq1
Title : Crystal structure of N5-glutamine methyltransferase, HemK(EC 2.1.1.-)
(TM0488) from Thermotoga maritima at 2.80 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2004-12-02
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	 3% 69% 18% • 10%
1	B	294	 5% 67% 19% • 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N5-glutamine methyltransferase, HemK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	Se	0	0	0
			2075	1337	344	390	1	3			
1	B	264	Total	C	N	O	S	Se	0	0	0
			2041	1315	336	386	1	3			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MSE	-	expression tag	UNP Q9WYV8
A	-10	GLY	-	expression tag	UNP Q9WYV8
A	-9	SER	-	expression tag	UNP Q9WYV8
A	-8	ASP	-	expression tag	UNP Q9WYV8
A	-7	LYS	-	expression tag	UNP Q9WYV8
A	-6	ILE	-	expression tag	UNP Q9WYV8
A	-5	HIS	-	expression tag	UNP Q9WYV8
A	-4	HIS	-	expression tag	UNP Q9WYV8
A	-3	HIS	-	expression tag	UNP Q9WYV8
A	-2	HIS	-	expression tag	UNP Q9WYV8
A	-1	HIS	-	expression tag	UNP Q9WYV8
A	0	HIS	-	expression tag	UNP Q9WYV8
A	1	MSE	MET	modified residue	UNP Q9WYV8
A	89	MSE	MET	modified residue	UNP Q9WYV8
A	193	MSE	MET	modified residue	UNP Q9WYV8
A	245	MSE	MET	modified residue	UNP Q9WYV8
B	-11	MSE	-	expression tag	UNP Q9WYV8
B	-10	GLY	-	expression tag	UNP Q9WYV8
B	-9	SER	-	expression tag	UNP Q9WYV8
B	-8	ASP	-	expression tag	UNP Q9WYV8
B	-7	LYS	-	expression tag	UNP Q9WYV8
B	-6	ILE	-	expression tag	UNP Q9WYV8
B	-5	HIS	-	expression tag	UNP Q9WYV8
B	-4	HIS	-	expression tag	UNP Q9WYV8
B	-3	HIS	-	expression tag	UNP Q9WYV8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP Q9WYV8
B	-1	HIS	-	expression tag	UNP Q9WYV8
B	0	HIS	-	expression tag	UNP Q9WYV8
B	1	MSE	MET	modified residue	UNP Q9WYV8
B	89	MSE	MET	modified residue	UNP Q9WYV8
B	193	MSE	MET	modified residue	UNP Q9WYV8
B	245	MSE	MET	modified residue	UNP Q9WYV8

- Molecule 2 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		
3	B	5	Total	O	0	0
			5	5		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.97Å 58.61Å 85.67Å 90.00° 109.46° 90.00°	Depositor
Resolution (Å)	36.09 – 2.80 36.09 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.9 (36.09-2.80) 95.9 (36.09-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.208 , 0.273 0.212 , 0.272	Depositor DCC
R_{free} test set	801 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	68.0	Xtriage
Anisotropy	0.727	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 88.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4177	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	1/2109 (0.0%)	1.00	6/2836 (0.2%)
1	B	0.84	0/2073	0.96	5/2790 (0.2%)
All	All	0.88	1/4182 (0.0%)	0.98	11/5626 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	173	PHE	C-O	-5.06	1.18	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	215	PRO	N-CA-C	5.99	118.01	110.70
1	A	213	PHE	N-CA-C	-5.92	106.39	113.97
1	A	117	ILE	CB-CA-C	-5.84	104.16	112.22
1	A	14	ILE	N-CA-C	5.73	116.47	110.62
1	B	56	LEU	N-CA-C	5.59	115.49	108.45
1	B	213	PHE	N-CA-C	-5.37	106.89	113.50
1	A	56	LEU	N-CA-C	5.09	114.86	108.45
1	B	214	GLU	CA-C-N	5.08	125.61	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	214	GLU	C-N-CA	5.08	125.61	120.38
1	A	155	LYS	N-CA-C	-5.03	105.68	111.07
1	A	250	ASP	N-CA-C	5.01	118.66	112.54

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	281	SER	Peptide
1	A	52	PHE	Peptide
1	B	281	SER	Peptide
1	B	52	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2075	0	2077	32	0
1	B	2041	0	2034	35	0
2	A	27	0	22	2	0
2	B	27	0	22	2	0
3	A	2	0	0	0	0
3	B	5	0	0	0	0
All	All	4177	0	4155	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:116:LEU:HB3	1:B:193:MSE:HE1	1.60	0.83
1:A:116:LEU:HD13	1:A:193:MSE:HE1	1.67	0.77
1:A:151:ASP:OD1	2:A:301:SAM:O2'	2.05	0.72
1:B:108:GLU:OE2	1:B:268:SER:OG	2.09	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:193:MSE:HE2	1:B:242:ILE:HG21	1.78	0.64
1:A:218:ALA:O	2:A:301:SAM:HE1	2.00	0.62
1:A:275:LEU:HD12	1:A:275:LEU:C	2.26	0.61
1:A:200:TYR:HB2	1:A:251:GLN:HE22	1.66	0.61
1:A:109:LEU:HD23	1:A:109:LEU:C	2.26	0.60
1:A:14:ILE:HG22	1:A:56:LEU:HA	1.84	0.60
1:B:229:TYR:CD2	1:B:245:MSE:HE3	2.39	0.58
1:A:116:LEU:HD13	1:A:193:MSE:CE	2.33	0.57
1:B:14:ILE:HG22	1:B:56:LEU:HA	1.85	0.57
1:B:197:ASN:O	2:B:301:SAM:HE3	2.06	0.56
1:B:153:SER:O	1:B:157:VAL:HG23	2.07	0.55
1:A:66:ILE:O	1:A:70:VAL:HG23	2.08	0.53
1:B:78:PRO:HD3	1:B:213:PHE:CB	2.39	0.52
1:B:100:PHE:CD2	2:B:301:SAM:HG2	2.44	0.52
1:B:122:ILE:CD1	1:B:193:MSE:HE3	2.40	0.52
1:B:165:GLU:HA	1:B:165:GLU:OE1	2.10	0.51
1:A:78:PRO:HD3	1:A:213:PHE:CB	2.41	0.50
1:A:248:GLY:N	1:A:251:GLN:OE1	2.43	0.50
1:B:15:TRP:HA	1:B:18:ILE:HD12	1.94	0.49
1:B:182:GLU:N	1:B:183:PRO:CD	2.75	0.49
1:B:69:LEU:HD22	1:B:82:ILE:HG23	1.95	0.48
1:B:52:PHE:O	1:B:53:LEU:HD22	2.13	0.48
1:A:125:VAL:O	1:A:147:VAL:HA	2.13	0.47
1:B:46:ILE:HG23	1:B:47:ARG:N	2.30	0.47
1:B:193:MSE:HE2	1:B:242:ILE:CG2	2.45	0.46
1:B:114:LEU:O	1:B:118:ARG:HG3	2.15	0.46
1:A:165:GLU:HA	1:A:165:GLU:OE1	2.16	0.46
1:A:21:CYS:O	1:A:22:SER:C	2.59	0.45
1:A:93:PHE:CZ	1:A:134:ALA:HA	2.50	0.45
1:B:200:TYR:O	1:B:251:GLN:NE2	2.49	0.45
1:A:96:GLU:O	1:A:97:GLU:C	2.59	0.45
1:B:46:ILE:CG2	1:B:47:ARG:N	2.80	0.45
1:A:200:TYR:OH	1:A:246:GLU:OE2	2.26	0.45
1:B:166:ARG:CG	1:B:166:ARG:O	2.64	0.45
1:B:76:GLY:O	1:B:77:TYR:C	2.59	0.44
1:A:25:LEU:HD21	1:A:70:VAL:HG13	1.99	0.44
1:B:96:GLU:O	1:B:97:GLU:C	2.60	0.44
1:B:35:GLU:O	1:B:39:ILE:HG13	2.18	0.43
1:A:106:THR:HG21	1:A:135:ILE:HD11	1.99	0.43
1:A:149:ALA:O	1:A:175:VAL:HA	2.18	0.43
1:A:78:PRO:HD3	1:A:213:PHE:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ASP:C	1:A:52:PHE:N	2.77	0.43
1:B:178:GLY:HA3	1:B:183:PRO:HG3	2.01	0.43
1:A:50:ASP:C	1:A:52:PHE:H	2.26	0.43
1:B:66:ILE:O	1:B:70:VAL:HG23	2.19	0.42
1:A:106:THR:CG2	1:A:135:ILE:HD11	2.49	0.42
1:B:50:ASP:C	1:B:52:PHE:H	2.28	0.42
1:A:166:ARG:HG2	1:A:166:ARG:O	2.20	0.41
1:A:224:ASP:OD2	1:A:230:ARG:NH2	2.53	0.41
1:B:215:PRO:HA	1:B:216:PRO:HD2	1.89	0.41
1:B:229:TYR:CE2	1:B:245:MSE:HE3	2.56	0.41
1:A:117:ILE:O	1:A:121:GLY:N	2.53	0.41
1:A:215:PRO:HA	1:A:216:PRO:HD2	1.87	0.41
1:B:212:LEU:HD13	1:B:216:PRO:HG3	2.02	0.41
1:A:193:MSE:CE	1:A:242:ILE:HG21	2.50	0.41
1:B:200:TYR:HB2	1:B:251:GLN:HE22	1.86	0.41
1:B:87:GLU:OE1	1:B:90:GLY:HA2	2.21	0.41
1:A:130:THR:HB	1:A:149:ALA:HB1	2.04	0.40
1:A:182:GLU:N	1:A:183:PRO:CD	2.84	0.40
1:A:275:LEU:C	1:A:275:LEU:CD1	2.95	0.40
1:B:146:ILE:HG22	1:B:172:ARG:HG2	2.03	0.40
1:B:166:ARG:O	1:B:166:ARG:HG2	2.21	0.40
1:B:238:THR:OG1	1:B:279:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/294 (89%)	241 (92%)	19 (7%)	2 (1%)	16	44
1	B	260/294 (88%)	242 (93%)	16 (6%)	2 (1%)	16	44
All	All	522/588 (89%)	483 (92%)	35 (7%)	4 (1%)	16	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	55	ASP
1	A	55	ASP
1	A	57	GLY
1	B	57	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/250 (88%)	204 (92%)	17 (8%)	12	36
1	B	216/250 (86%)	198 (92%)	18 (8%)	10	32
All	All	437/500 (87%)	402 (92%)	35 (8%)	11	34

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	VAL
1	A	29	THR
1	A	32	SER
1	A	46	ILE
1	A	58	VAL
1	A	72	LYS
1	A	83	LEU
1	A	106	THR
1	A	112	LEU
1	A	159	ILE
1	A	186	GLU
1	A	217	GLU
1	A	224	ASP
1	A	259	VAL
1	A	260	SER
1	A	275	LEU
1	A	281	SER
1	B	28	VAL
1	B	41	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	46	ILE
1	B	58	VAL
1	B	96	GLU
1	B	112	LEU
1	B	122	ILE
1	B	146	ILE
1	B	154	SER
1	B	157	VAL
1	B	159	ILE
1	B	182	GLU
1	B	186	GLU
1	B	217	GLU
1	B	224	ASP
1	B	250	ASP
1	B	259	VAL
1	B	268	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SAM	A	301	-	27,29,29	1.17	4 (14%)	34,42,42	2.32	13 (38%)
2	SAM	B	301	-	27,29,29	1.14	4 (14%)	34,42,42	2.23	10 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAM	A	301	-	-	1/17/33/33	0/3/3/3
2	SAM	B	301	-	-	7/17/33/33	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	SAM	C8-N7	2.70	1.36	1.31
2	A	301	SAM	C2-N1	2.57	1.38	1.33
2	A	301	SAM	C8-N7	2.54	1.36	1.31
2	B	301	SAM	C2-N3	2.43	1.38	1.33
2	B	301	SAM	C2-N1	2.34	1.38	1.33
2	A	301	SAM	C2-N3	2.30	1.38	1.33
2	B	301	SAM	OXT-C	-2.28	1.23	1.30
2	A	301	SAM	OXT-C	-2.03	1.24	1.30

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	SAM	N3-C2-N1	-6.37	118.94	128.58
2	B	301	SAM	N3-C2-N1	-6.00	119.49	128.58
2	B	301	SAM	O4'-C1'-N9	-4.37	99.69	108.09
2	A	301	SAM	C2'-C1'-N9	4.33	124.06	113.30
2	B	301	SAM	C5-C4-N3	-4.26	120.86	126.72
2	A	301	SAM	C5-C4-N3	-4.08	121.11	126.72
2	B	301	SAM	C2'-C1'-N9	3.95	123.12	113.30
2	A	301	SAM	N9-C8-N7	-3.89	108.42	113.94
2	A	301	SAM	C5-N7-C8	3.88	109.54	103.45
2	B	301	SAM	N9-C8-N7	-3.69	108.70	113.94
2	A	301	SAM	C2-N3-C4	3.43	120.21	111.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	SAM	C2-N3-C4	3.40	120.15	111.83
2	B	301	SAM	C5-N7-C8	3.33	108.69	103.45
2	A	301	SAM	C4-C5-N7	-3.28	106.83	110.58
2	A	301	SAM	O4'-C1'-N9	-3.17	102.01	108.09
2	B	301	SAM	N3-C4-N9	2.51	131.43	127.17
2	B	301	SAM	C4-C5-N7	-2.50	107.73	110.58
2	A	301	SAM	OXT-C-O	-2.38	118.67	124.08
2	A	301	SAM	C4'-O4'-C1'	2.37	114.70	109.47
2	A	301	SAM	C5-C4-N9	2.12	108.12	105.81
2	A	301	SAM	N3-C4-N9	2.11	130.76	127.17
2	A	301	SAM	C2-N1-C6	2.07	122.13	118.73
2	B	301	SAM	C2'-C3'-C4'	2.00	106.48	102.61

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	SAM	N-CA-CB-CG
2	B	301	SAM	C-CA-CB-CG
2	B	301	SAM	OXT-C-CA-N
2	B	301	SAM	C4'-C5'-SD-CG
2	B	301	SAM	CB-CG-SD-CE
2	A	301	SAM	OXT-C-CA-N
2	B	301	SAM	CB-CG-SD-C5'
2	B	301	SAM	O-C-CA-N


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SAM	2	0
2	B	301	SAM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/294 (89%)	0.48	9 (3%) 48 39	81, 88, 92, 106	0
1	B	261/294 (88%)	0.82	14 (5%) 31 24	84, 88, 90, 94	0
All	All	524/588 (89%)	0.65	23 (4%) 39 31	81, 88, 91, 106	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	53	LEU	4.1
1	A	57	GLY	3.2
1	A	206	HIS	3.0
1	B	54	LYS	2.9
1	A	53	LEU	2.8
1	A	217	GLU	2.8
1	B	81	TYR	2.8
1	B	204	SER	2.7
1	B	56	LEU	2.6
1	A	220	PHE	2.6
1	B	95	VAL	2.5
1	B	55	ASP	2.3
1	A	205	ALA	2.3
1	B	30	GLU	2.3
1	B	274	PHE	2.2
1	B	277	LEU	2.2
1	A	145	ALA	2.2
1	B	217	GLU	2.2
1	B	146	ILE	2.1
1	A	148	PHE	2.1
1	A	54	LYS	2.1
1	B	148	PHE	2.0
1	B	175	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SAM	B	301	27/27	0.78	0.16	80,96,103,103	1
2	SAM	A	301	27/27	0.81	0.19	80,89,94,95	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SAM B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAM A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.